



US 20080069821A1

(19) **United States**(12) **Patent Application Publication**  
**Yang et al.**(10) **Pub. No.: US 2008/0069821 A1**(43) **Pub. Date: Mar. 20, 2008**(54) **INFLUENZA HEMAGGLUTININ AND  
NEURAMINIDASE VARIANTS****Publication Classification**(75) Inventors: **Chin-Fen Yang**, San Jose, CA (US);  
**George Kemble**, Saratoga, CA (US);  
**Kanta Subbarao**, Washington, DC  
(US); **Brian Murphy**, Bethesda, MD  
(US)

Correspondence Address:

**JOHNATHAN KLEIN-EVANS**  
**ONE MEDIMMUNE WAY**  
**GAITHERSBURG, MD 20878 (US)**(73) Assignees: **MedImmune Vaccines, Inc.**, Gaithers-  
burg, MD (US); **The Government of the**  
**United States of America National**  
**Institutes of Health**, Rockville, MD (US)(21) Appl. No.: **11/836,413**(22) Filed: **Aug. 9, 2007****Related U.S. Application Data**(60) Provisional application No. 60/821,832, filed on Aug.  
9, 2006. Provisional application No. 60/942,804, filed  
on Jun. 8, 2007.(51) **Int. Cl.****A61K 39/145** (2006.01)  
**A61K 35/76** (2006.01)  
**A61K 39/00** (2006.01)  
**A61P 31/16** (2006.01)  
**C12N 1/19** (2006.01)  
**C12N 15/63** (2006.01)  
**C12N 7/01** (2006.01)  
**C12P 21/00** (2006.01)  
**C12N 5/10** (2006.01)  
**C12N 1/21** (2006.01)  
**C07K 14/00** (2006.01)  
**A61K 39/395** (2006.01)  
**A61K 39/12** (2006.01)(52) **U.S. Cl.** ..... **424/139.1**; 424/186.1; 424/206.1;  
424/93.2; 435/235.1; 435/252.33;  
435/254.2; 435/325; 435/349;  
435/350; 435/352; 435/364;  
435/365; 435/69.3; 530/350

(57)

**ABSTRACT**Polypeptides, polynucleotides, methods, compositions, and  
vaccines comprising (avian pandemic) influenza hemagglu-  
tinin and neuraminidase variants are provided.

VN/1203/2004 wildtype HA:

CCT	CAA	AGA	GAG	AGA	AGA	AGA	AAA	AAG	AGA	GGA	TTA	TTT
Pro	Gln	Arg	Glu	Arg	Arg	Arg	Lys	Lys	Arg	Gly	Leu	Phe

Modified HA:

CCT	CAA	AGA	GAG	<u>ACT</u>	<u>CGA</u>	GGA	TTA	TTT
Pro	Gln	Arg	Glu	<u>Thr</u>	Arg	Gly	Leu	Phe

↓ Site of cleavage of into HA1 and HA2 domains.

Residues that were mutagenized are underlined.

Fig 1.

Virus	Mortality (dead/total)	Virus isolation from swabs				Antibody detected/ total
		Oropharyngeal		Cloacal		
		# shedding/ total	Mean log <sub>10</sub> titer (EID <sub>50</sub> )	# shedding/ total	Mean log <sub>10</sub> titer (EID <sub>50</sub> )	
1997, 2003 and 2004 H5N1 wt	8/8	8/8	>6.3	8/8	>4.5	NA
1997, 2003 and 2004 H5N1 ca	0/8	0/8	<0.9	0/8	<0.9	0/8

Chickens were inoculated intranasally with  $10^6$   $\text{TCID}_{50}$  of virus.

**Fig 2.**


<b>LD<sub>50</sub> in mice</b>	
<b>A/AA/6/60 ca</b>	<b>&gt;10<sup>7</sup> TCID<sub>50</sub></b>
<b>A/HK/491/97</b>	<b>10<sup>2</sup> TCID<sub>50</sub></b>
<b>1997 H5N1/AA ca</b>	<b>&gt;10<sup>7</sup> TCID<sub>50</sub></b>
<b>A/HK/213/2003</b>	<b>10<sup>6</sup> TCID<sub>50</sub></b>
<b>2003 H5N1/AA ca</b>	<b>&gt;10<sup>7</sup> TCID<sub>50</sub></b>
<b>A/Vietnam/1203/2004</b>	<b>10<sup>0.4</sup> TCID<sub>50</sub></b>
<b>2004 H5N1/AA ca</b>	<b>&gt;10<sup>7</sup> TCID<sub>50</sub></b>

**Fig 3.**

Tissue	Virus	Average fold difference in titer Over 3 days
LUNGS	A/AA/6/60	93
	1997 H5N1	501
	2003 H5N1	12
	2004 H5N1	430
NASAL TURBINATES	A/AA/6/60	32
	1997 H5N1	185
	2003 H5N1	none
	2004 H5N1	100

$10^6$  TCID<sub>50</sub> of virus was administered intranasally and tissues were harvested on days 2, 3 or 4 post-infection. Virus titers are expressed as log<sub>10</sub> TCID<sub>50</sub>/g of tissue.

Fig 4.



Immunizing virus	Geometric mean serum HAI Ab titers against indicated virus		
	1997 wt	2003 wt	2004 wt
2003 ca	20	213.6	20
2003 wt	20	394	20

An undetectable titer is assigned a value of 20

Fig 6.

Immunizing virus	Geometric mean serum neutralizing Ab titers against indicated virus		
	1997 wt	2003 wt	2004 wt
2003 ca	10	59.2	10
2003 wt	10	93.3	10

An undetectable titer is assigned a value of 10

Fig 7.






Fig 8.

Immunization	Mean reduction in titer in lungs following			
	Homologous challenge	Heterologous H5N1 challenge		
		1997 wt	2003 wt	2004 wt
1997 ca	2.5	NA	3.0	0.7
2003 ca	>5.8	2.3	NA	2.9
2004 ca	2.0	1.4	>5.7	NA

Fig 9.

Immunization	Mean reduction in titer in NT following			
	Homologous challenge	Heterologous H5N1 challenge		
		1997 wt	2003 wt	2004 wt
1997 ca	4.3	NA	>1.2	2.6
2003 ca	>1.2	3.7	NA	>3.3
2004 ca	1.6	4.2	>3.5	NA

Fig 10.




Fig 11.





Fig 12.



Challenge dose  
Fig 13.

A/Netherland/219/03 wildtype HA:

CCA	AAG	AGG	AGG	AGG	AGA	↓	GGC
Pro	Lys	Arg	Arg	Arg	Arg		Gly

Modified HAs:

CCA	AAG	GGG	---	---	AGA		GGC
Pro	Lys	Gly			Arg		Gly

↓ Site of cleavage of into  
HA1 and HA2 domains.

CCA	AAG	ACT	---	---	AGA		GGC
Pro	Lys	Thr			Arg		Gly

CCA	AAG	CCG	---	---	AGA		GGC
Pro	Lys	Pro			Arg		Gly

Fig 14.

Immunizing virus	Doses	Geometric mean serum neutralizing Ab titers against indicated virus		
		1997 wt	2003 wt	2004 wt
<i>A/VN/2004 ca</i>	1	10	10	10
	2	160	528	388
<i>A/HK/2003 ca</i>	1	10	37	10
	2	19	1056	61

Fig 15.






Fig 16.

		<i>Post Vaccination Sera (7 log<sub>10</sub>PFU) HI Titer 32 days after dose 1</i>			
		H6N9 A/Duck <i>ca</i>	H6N1 A/Teal <i>ca</i>	H6N2 A/Mallard <i>ca</i>	H1N1 A/New Cal <i>ca</i>
<i>Test Antigen</i>	Wt A/Duck	13.5	<8	<8	<8
	Wt A/Teal	<8	19.0	<8	<8
	Wt A/Mallard	<8	<8	13.5	<8
	Wt A/New Cal	<8	<8	<8	430.5

Fig 17.




Fig 18.AB




Fig 19.

Virus	Dosing schedule	Geometric mean neutralizing antibody titers achieved at indicated time post-infection <sup>a</sup>		
		d 0	4 weeks	8 weeks
	d 0	<10	80	NA
H7N3 BC 04 ca	d 0	<10	87	403 <sup>b</sup>
	d 0 and d 28	<10	45	470
	d 0	<10	308	NA
A/ck/BC/CN-6/04	d 0	<10	320	941
	d 0 and d 28	<10	154	1701

Fig 20.




Fig 21. A-C




Fig 21. D-F




Fig 21. G-I






Fig 22.A





Fig 22.B



Virus

Fig 23.




Fig 24.AB

---

Vaccine Dose	N	No. Vaccine Virus Culture + (days)	Peak Titer Shed* (log <sub>10</sub> TCID <sub>50</sub> /mL)	No. Vaccine Virus PCR+ (days)
1	26	2 (1)	1.0	8 (1.1)
2	24	0	≤0.6	2 (1.5)

---

\* Among those who shed virus

Fig 25.

Vaccine Dose	Sero- status	N	Mean HI titer, 1/log <sub>2</sub>			No. with 4-fold rise	%
			Days After Dose				
			D0	D28	D42		
1	-	26	1.7	2.6	- -	9	31
2	-	24	3.1	4.6	4.5	12	50
Cumulative		24				22	92

Fig 26.

<u>Vaccine Dose</u>	<u>N</u>	<u>No. Nasal wash Culture + (days)</u>	<u>No. Nasal wash PCR+ (days)</u>
1	21	0	2 (2.0)
2	18	0	3 (1.0)

Vaccine virus was not detected by any method in throat swab specimens.

**Fig 27.**

Vaccine Dose	N	Mean HI titer, 1/log <sub>2</sub>			
		Days after dose		No. with 4-fold rise	%
		D0	D28		
1	20	2.6	2.8	0	0
2	18	2.7	3.1	1	6
Cumulative	18			2	11

Fig 28.



## INFLUENZA HEMAGGLUTININ AND NEURAMINIDASE VARIANTS

### CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit under 35 U.S.C. § 119 (e) of U.S. Provisional Application Nos. 60/821,832 filed Aug. 9, 2006 and 60/942,804, filed Jun. 8, 2007, the disclosures of each of which are incorporated herein in their entirety for all purposes.

### BACKGROUND OF THE INVENTION

[0002] Vaccines against various and evolving strains of influenza are important from a community health stand point, as well as commercially, since each year numerous individuals are infected with different strains and types of influenza virus. Infants, the elderly, those without adequate health care and immuno-compromised persons are at special risk of death from such infections. Compounding the problem of influenza infections is that novel influenza strains evolve readily and can spread amongst various species, thereby necessitating the continuous production of new vaccines.

[0003] Numerous vaccines capable of producing a protective immune response specific for such different and influenza viruses/virus strains have been produced for over 50 years and include whole virus vaccines, split virus vaccines, surface antigen vaccines and live attenuated virus vaccines. However, while appropriate formulations of any of these vaccine types are capable of producing a systemic immune response, live attenuated virus vaccines have the advantage of also being able to stimulate local mucosal immunity in the respiratory tract. Considerable work in the production of influenza viruses, and fragments thereof, for production of vaccines has been done by the present inventors and co-workers; see, e.g., U.S. Application Nos. 60/420,708, filed Oct. 23, 2002; 60/574,117, filed May 24, 2004; Ser. No. 10/423,828, filed Apr. 25, 2003; 60/578,962, filed Jun. 12, 2004; and Ser. No. 10/870,690 filed Jun. 16, 2004, the disclosure of which is incorporated by reference herein.

[0004] Because of the continual emergence (or re-emergence) of different influenza strains, new influenza vaccines are continually desired. Such vaccines typically are created using antigenic moieties of the newly emergent virus strains, thus, polypeptides and polynucleotides of novel, newly emergent, or newly re-emergent virus strains (especially sequences of antigenic genes) are highly desirable.

[0005] The present invention provides new and/or newly isolated influenza hemagglutinin and neuraminidase variants that are capable of use in production of numerous types of vaccines as well as in research, diagnostics, etc. Numerous other benefits will become apparent upon review of the following.

### SUMMARY OF THE INVENTION

[0006] In some aspects herein, the invention comprises an isolated or recombinant polypeptide that is selected from: any one of the polypeptides encoded by SEQ ID NO:1 through SEQ ID NO:10 or SEQ ID NO:21 through SEQ ID NO:26 or SEQ ID NO:33 through SEQ ID NO:38; any one of the polypeptides of SEQ ID NO:11 through SEQ ID

NO:20 or SEQ ID NO:27 through SEQ ID NO:32 or SEQ ID NO:39 through SEQ ID NO:44; only the open reading frame encoding the polypeptides of SEQ ID NO:11 through SEQ ID NO:20 or SEQ ID NO:27 through SEQ ID NO:32 or SEQ ID NO:39 through SEQ ID NO:44; any alternative (e.g., the mature form without the signal peptide, or without the 5' and 3' sequences outside of the open reading frame, or the sequences as expressed on the surface of a virus (e.g., influenza)) form of the polypeptides of SEQ ID NO:11-20 or SEQ ID NO:27-32 or SEQ ID NO:39-44; any polypeptide that is encoded by a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of a polynucleotide sequence of SEQ ID NO:1 through SEQ ID NO:10 or SEQ ID NO:21 through SEQ ID NO:26, SEQ ID NO:33-38, or SEQ ID NO:45; any polypeptide that is encoded by a polynucleotide sequence which hybridizes under highly stringent conditions to a polynucleotide sequence of SEQ ID NO:1 through SEQ ID NO:10 or SEQ ID NO:21 through SEQ ID NO:26 or SEQ ID NO:33 through SEQ ID NO:38, or SEQ ID NO:45; and, a fragment of any of the above wherein the sequence comprises a hemagglutinin or neuraminidase polypeptide, or a fragment of a hemagglutinin or neuraminidase polypeptide, preferably where the fragments generate an antibody that specifically binds a full length polypeptide of the invention. In various embodiments, the isolated or recombinant polypeptides of the invention are substantially identical to about 300 contiguous amino acid residues of any of the above polypeptides. In yet other embodiments, the invention comprises isolated or recombinant polypeptides, that comprise an amino acid sequence that is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at least about 450 amino acids; over at least about 500 amino acids; over at least about 520 amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids contiguous of any of the above polypeptides. In some embodiments, the polypeptide sequence (e.g., as listed in "SEQUENCES" herein) comprises less than 565, 559, etc. amino acids. In such embodiments, the shorter listed polypeptides optionally comprise less than 565, 559, etc. amino acids. In yet other embodiments, the polypeptides of the invention optionally comprise fusion proteins, proteins with a leader sequence, a precursor polypeptide, proteins with a secretion signal or a localization signal, or proteins with an epitope tag, an E-tag, or a His epitope tag. In still other embodiments, the invention comprises a polypeptide comprising a sequence having at least 85%, at least 90%, at least 93%, at least 95%, at least 98%, at least 98.5%, at least 99%, at least 99.2%, at least 99.4%, at least 99.6%, at least 99.8%, or at least 99.9% sequence identity to at least one polypeptide listed above. The hemagglutinin sequences of the invention can comprise both those sequences with unmodified and modified polybasic cleavage sites (thereby allowing growth of the viruses in eggs). The hemagglutinin polypeptide sequences of SEQ ID NOS:11, 13, 15, 17, 19, 27, 29, 31, 39, 41, or 43 comprise the endogenous amino terminal signal peptide sequences, however, the hemagglutinin polypeptide sequences of the invention also include the mature (amino terminal signal peptide cleaved) form of the hemagglutinin polypeptides. The cleavage sites of any hemagglutinin polypeptide sequence of any influenza strain can be routinely measured or predicted using any number of methods in the art.

[0007] In other aspects, the invention comprises a composition with one or more polypeptide listed above, or fragments thereof. The invention also includes polypeptides that are specifically bound by a polyclonal antisera raised against at least 1 antigen that comprises at least one amino acid sequence described above, or a fragment thereof. Such antibodies specific for the polypeptides described above are also features of the invention. The polypeptides of the invention are optionally immunogenic.

[0008] The invention also encompasses immunogenic compositions comprising an immunologically effective amount of one or more of any of the polypeptides described above as well as methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of any of the above polypeptides in a physiologically acceptable carrier.

[0009] Additionally, the invention includes recombinant influenza virus that comprises one or more of the polypeptides or polynucleotides above, in addition to immunogenic compositions comprising an immunologically effective amount of such recombinant influenza virus. Methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus, through administering an immunologically effective amount of such recombinant influenza virus in a physiologically acceptable carrier are also part of the invention.

[0010] In other aspects, the invention comprises an isolated or recombinant nucleic acid that is selected from: any one of the polynucleotide sequences SEQ ID NO:1 through SEQ ID NO:10 or SEQ ID NO:21 through SEQ ID NO:26 or SEQ ID NO:33 through SEQ ID NO:38, or SEQ ID NO:45 (or complementary sequences thereof), any one of the polynucleotide sequences encoding a polypeptide of SEQ ID NO:11 through SEQ ID NO:20 or SEQ ID NO:27 through SEQ ID NO:32 or SEQ ID NO:39 through SEQ ID NO:44 (or complementary polynucleotide sequences thereof), a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of any of the above polynucleotide sequences, and a polynucleotide sequence comprising all or a fragment of any of such polynucleotide sequences wherein the sequence preferably encodes a hemagglutinin or neuraminidase polypeptide or a fragment of a hemagglutinin or neuraminidase polypeptide. The invention also includes an isolated or recombinant nucleic acid that encodes an amino acid sequence which is substantially identical over at least about 300 amino acids of any polypeptide encoded by the above nucleic acids, or over at least about 350 amino acids; over at least about 400 amino acids; over at least about 450 amino acids; over at least about 500 amino acids; over at least about 502 amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids of any polypeptide encoded by the above nucleic acids. Again, in situations wherein the amino acid is less than, e.g., 566, 565, 559, etc. in length (e.g., see, "SEQUENCES") then it should be understood that the length is optionally less than 566, 565, 559, etc. The invention also includes any of the above nucleic acids that comprise a polynucleotide encoding a hemagglutinin or neuraminidase polypeptide, or one or more fragments of one or more hemagglutinin or neuraminidase polypeptide. Other aspects of the invention include isolated

or recombinant nucleic acids that encode a polypeptide (optionally a hemagglutinin or neuraminidase polypeptide) whose sequence has at least 98% identity, at least 98.5% identity, at least 99% identity, at least 99.2% identity, at least 99.4% identity, at least 99.6% identity, at least 99.8% identity, or at least 99.9% identity to at least one of the above described polynucleotides. The invention also includes isolated or recombinant nucleic acids encoding a polypeptide of hemagglutinin or neuraminidase produced by mutating or recombining one or more above described polynucleotide sequences. The polynucleotide sequences of the invention can optionally comprise one or more of, e.g., a leader sequence, a precursor sequence, or an epitope tag sequence or the like, and can optionally encode a fusion protein (e.g., with one or more additional nucleic acid sequences).

[0011] In yet other embodiments, the invention comprises a composition of matter having two or more above described nucleic acids (e.g., a library comprising at least about 2, 5, 10, 50 or more nucleic acids). Such compositions can optionally be produced by cleaving one or more above described nucleic acid (e.g., mechanically, chemically, enzymatically with a restriction endonuclease/RNase/DNase, etc.). Other compositions of the invention include, e.g., compositions produced by incubating one or more above described nucleic acid in the presence of deoxyribonucleotide triphosphates and a thermostable nucleic acid polymerase.

[0012] The invention also encompasses cells comprising at least one of the above described nucleic acids, or a cleaved or amplified fragment or product thereof. Such cells can optionally express a polypeptide encoded by such nucleic acid. Other embodiments of the invention include vectors (e.g., plasmids, cosmids, phage, viruses, virus fragments, etc.) comprising any of above described nucleic acids. Such vectors can optionally comprise an expression vector. Preferred expression vectors of the invention include, but are not limited to, vectors comprising pol I promoter and terminator sequences or vectors using both the pol I and pol II promoters "the polI/polII promoter system" (e.g., Zobel et al., Nucl. Acids Res. 1993, 21:3607; US20020164770; Neumann et al., Proc. Natl. Acad. Sci. USA 1999, 96:9345; Fodor et al., J. Virol. 1999, 73:9679; and US20030035814). Cells transduced by such vectors are also within the current invention.

[0013] In some embodiments, the invention encompasses a virus (e.g., an influenza virus) comprising one or more above described nucleic acids (e.g., encoding hemagglutinin and/or neuraminidase), or one or more fragments thereof. Immunogenic compositions comprising such virus are also part of the current invention. Such viruses can comprises a reassortment virus such as a 6:2 reassortment virus (e.g., comprising 6 genes encoding regions from one or more donor virus and 2 genes encoding regions from one or more above described nucleotide sequence (or one or more fragment thereof) which can optionally comprise hemagglutinin and/or neuraminidase). Reassortment viruses (optionally live viruses) of the invention can include donor viruses that are one or more of, e.g., cold-sensitive, cold-adapted, or an attenuated. For example, reassortment viruses can comprise e.g., A/Ann Arbor/6/60, PR8, etc. Reassortment viruses of the invention may alternatively exclude A/Ann Arbor/6/60. One preferred embodiment of the invention is a reassortant influenza virus, wherein the virus is a 6:2 reassortment

influenza virus and comprises 6 gene encoding regions from A/Ann Arbor/6/60 and 2 gene encoding regions that encode a polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS:11-20, SEQ ID NOS:27-32, and SEQ ID NOS:39-44. In an alternative embodiment, a reassortant influenza virus of the invention includes a 6:2 reassortment influenza virus, wherein said virus comprises 6 gene encoding regions from one or more donor viruses other than A/Ann Arbor/6/60 and 2 gene encoding regions that encode a polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS:11-20, SEQ ID NOS:27-32, and SEQ ID NOS:39-44. In another alternative embodiment, a reassortant influenza virus of the invention includes a 6:2 reassortment influenza virus, wherein said virus comprises 6 gene encoding regions from one or more donor viruses other than A/Ann Arbor/6/60 and 2 gene encoding regions, wherein the 2 gene encoding regions are HA or NA polypeptides from any pandemic influenza strain. Methods of producing recombinant influenza virus through culturing a host cell harboring an influenza virus in a suitable culture medium under conditions permitting expression of nucleic acid and, isolating the recombinant influenza virus from one or more of the host cell or the medium are also part of the invention.

**[0014]** In other embodiments herein, the invention comprises immunogenic compositions having an immunologically effective amount of any of the above described recombinant influenza virus. Other embodiments include methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of any of the recombinant influenza virus described above (optionally in a physiologically effective carrier).

**[0015]** Other aspects of the invention include methods of producing an isolated or recombinant polypeptide by culturing any host cell above, in a suitable culture medium under conditions permitting expression of nucleic acid and, isolating the polypeptide from one or more of the host cells or the medium in which is the cells are grown.

**[0016]** Immunogenic compositions are also features of the invention. For example, immunogenic compositions comprising one or more of any of the polypeptides and/or nucleic acids described above and, optionally, an excipient such as a pharmaceutically acceptable excipient or one or more pharmaceutically acceptable administration component. Immunogenic compositions of the invention can also comprise any one or more above described virus as well (e.g., along with one or more pharmaceutically acceptable administration component).

**[0017]** Methods of producing immunogenic responses in a subject through administration of an effective amount of any of the above viruses (or immunogenic compositions) to a subject are also within the current invention. Additionally, methods of prophylactic or therapeutic treatment of a viral infection (e.g., viral influenza) in a subject through administration of any one or more above described virus (or immunogenic compositions) in an amount effective to produce an immunogenic response against the viral infection are also part of the current invention. Subjects for such treatment can include mammals (e.g., humans). Such methods can also comprise in vivo administration to the subject

as well as in vitro or ex vivo administration to one or more cells of the subject. Additionally, such methods can also comprise administration of a composition of the virus and a pharmaceutically acceptable excipient that are administered to the subject in an amount effective to prophylactically or therapeutically treat the viral infection.

**[0018]** In other aspects the invention includes compositions of matter comprising nucleic acid sequences encoding hemagglutinin and/or neuraminidase polypeptides of one or more pandemic influenza strain and nucleic acid sequences encoding one or more polypeptide of A/Ann Arbor/6/60. Additionally, the invention includes compositions of matter comprising nucleic acid sequences encoding hemagglutinin and/or neuraminidase polypeptides of one or more pandemic influenza strain and nucleic acid sequences encoding one or more polypeptide of PR8 or A/Ann Arbor/6/60. Such sequences can include those listed in the "SEQUENCES" herein. Additionally, preferred embodiments of the invention include compositions of matter comprising sequences encoding hemagglutinin and/or neuraminidase of one or more pandemic influenza strain and nucleic acid sequences encoding a selected backbone strain in a 6:2 reassortment. Such compositions preferably include sequences encoding the hemagglutinin and neuraminidase selected from the "SEQUENCES" herein and a backbone strain, wherein the backbone strain is PR8 or A/Ann Arbor/6/60. The invention also includes such compositions as described above wherein the hemagglutinin comprises a modified polybasic cleavage site. The invention also includes live attenuated influenza vaccine comprising such above compositions.

**[0019]** These and other objects and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the accompanying figures and appendix.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0020]** FIG. 1: Shows modifications engineered into the HA gene of VN/1203/2004 to remove the polybasic cleavage site.

**[0021]** FIG. 2: Displays results showing that intranasally administered H5N1 ca reassortant viruses do not replicate in chickens.

**[0022]** FIG. 3: Illustrates that the H5N1/AA ca vaccine candidates are not lethal to mice.

**[0023]** FIG. 4: Illustrates that the 1997 and 2004 H5N1 ca reassortant viruses are restricted in replication in mice.

**[0024]** FIG. 5: Illustrates that the reassortant H5N1/AA ca influenza viruses are restricted in replication in lungs of mice.

**[0025]** FIG. 6: Shows the serum HAI Ab titers elicited in mice following a single i.n. dose of vaccine.

**[0026]** FIG. 7: Shows serum neutralizing Ab titers elicited in mice following a single i.n. dose of vaccine.

**[0027]** FIG. 8: Illustrates that H5N1 ca reassortant viruses protect mice from lethal challenges with 50, 500 or 5000 LD<sub>50</sub> of wild-type H5N1 viruses.

**[0028]** FIG. 9: Illustrates the efficacy of protection from pulmonary replication of homologous and heterologous H5N1 challenge viruses in mice.

[0029] FIG. 10: Illustrates the efficacy of protection from replication of homologous and heterologous H5N1 challenge viruses in the upper respiratory tract of mice.

[0030] FIG. 11: Illustrates the efficacy of protection conferred by 2004 H5N1 ca vaccine against high dose ( $10^5$ TCID<sub>50</sub>) challenge with homologous or heterologous H5N1 wt viruses in mice.

[0031] FIG. 12: Illustrates the efficacy of protection conferred by 1997 and 2003 H5N1 ca vaccines against high dose ( $10^5$ TCID<sub>50</sub>) challenges with homologous or heterologous H5N1 wild-type viruses in mice.

[0032] FIG. 13: Illustrates the efficacy of protection conferred by 2004 H5N1 ca vaccine against low or high doses of homologous H5N1 wild-type virus challenges in mice.

[0033] FIG. 14: Shows modifications that can be engineered into the HA gene of A/Netherlands/219/03 HA to remove the polybasic cleavage site.

[0034] FIG. 15: H5N1 ca vaccines elicit serum neutralizing antibody titers in mice. Sera were collected before (prebleed) and 28 days following each dose of vaccine; an undetectable titer is assigned a value of 10.

[0035] FIG. 16: H6 ca viruses are attenuated in ferrets. \*EID<sub>50</sub>/g for lungs; PFU/g for nasal turbinates.  $10^7$  TCID<sub>50</sub> inoculated intranasally, tissues were harvested on day 3 post-infection.

[0036] FIG. 17: Immunogenicity of H6 ca vaccines in ferrets.

[0037] FIGS. 18(a) and 18(b): Efficacy of H6 ca vaccines in ferrets. Virus titer was measured in (a) lungs and (b) nasal turbinates. Vaccine: 1 dose of  $7 \log_{10}$  PFU. Challenge:  $7 \log_{10}$  PFU; 3 days post-challenge.

[0038] FIG. 19: H7N3 BC 04 ca is attenuated in ferrets. Inoculum:  $10^7$  TCID<sub>50</sub> in 0.5 mL intranasally. Tissues were harvested on day 3 post-infection.

[0039] FIG. 20: H7N3 BC 2004 ca is immunogenic in mice. a: Reciprocal geometric mean of serum neutralizing antibody titers against ck/BC/CN-6/04 wt. b:  $p < 0.05$  (Mann-Whitney U-test) compared to neutralization titers at 28 days post-infection.

[0040] FIG. 21(a)-21(i): H7N3 BC 04 ca is efficacious against lethal challenge with H7 viruses in mice. Efficacy against a lethal challenge of 50 LD<sub>50</sub> A/ck/BC/CN-7/04: four weeks following immunization with a single dose (a), eight weeks following immunization with a single dose (b), or eight weeks following immunization with 2 doses (2 doses administered at 4 weeks apart) (c). Efficacy against a lethal challenge of 50 LD<sub>50</sub> A/NL/219/03: four weeks following immunization with a single dose (d), eight weeks following immunization with a single dose (e), or eight weeks following immunization with 2 doses (2 doses administered at 4 weeks apart) (f). Efficacy against a lethal challenge of 50 LD<sub>50</sub> A/tk/Eng/63: four weeks following immunization with a single dose (g), eight weeks following immunization with a single dose (h), or eight weeks following immunization with 2 doses (2 doses administered at 4 weeks apart) (i).

[0041] FIGS. 22(a) and (b): H7N3 BC 04 ca vaccine is efficacious in mice. Virus titer was measured at 8 weeks in (a) nasal turbinates and (b) lungs.

[0042] FIGS. 23(a) and (b): H9N2 G9/AA ca is attenuated in ferrets. Virus titer was measured in (a) nasal turbinates and (b) lungs.

[0043] FIGS. 24(a) and (b): Efficacy of the H9N2 ca vaccine in mice.

[0044] FIG. 25: Replication of H9N2 G9/AA ca is highly restricted in healthy adults.

[0045] FIG. 26: HI antibody responses to  $10^{7.0}$  TCID<sub>50</sub> of H9N2 G9/AA ca in healthy adults.

[0046] FIG. 27: Replication of H5N1 VN2004 A/AA ca is highly restricted in healthy adults.

[0047] FIG. 28: HI antibody responses to  $10^{6.7}$  TCID<sub>50</sub> of VN2004 A/AA ca in healthy adults.

## DETAILED DESCRIPTION

[0048] The present invention includes polypeptide and polynucleotide sequences of influenza hemagglutinin and neuraminidase as well as vectors, compositions and the like comprising such sequences and methods of their use. Additional features of the invention are described in more detail herein.

## Definitions

[0049] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. The following definitions supplement those in the art and are directed to the current application and are not necessarily to be imputed to any related or unrelated case, e.g., to any commonly owned patent or application. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein. Accordingly, the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

[0050] As used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a virus” includes a plurality of viruses; reference to a “host cell” includes mixtures of host cells, and the like.

[0051] An “amino acid sequence” is a polymer of amino acid residues (a protein, polypeptide, etc.) or a character string representing an amino acid polymer, depending on context.

[0052] The terms “nucleic acid,” “polynucleotide,” “polynucleotide sequence” and “nucleic acid sequence” refer to single-stranded or double-stranded deoxyribonucleotide or ribonucleotide polymers, chimeras or analogues thereof, or a character string representing such, depending on context. As used herein, the term optionally includes polymers of analogs of naturally occurring nucleotides having the essential nature of natural nucleotides in that they hybridize to single-stranded nucleic acids in a manner similar to naturally occurring nucleotides (e.g., peptide nucleic acids). Unless otherwise indicated, a particular nucleic acid sequence of this invention optionally encompasses complementary sequences in addition to the sequence explicitly indicated.

From any specified polynucleotide sequence, either the given nucleic acid or the complementary polynucleotide sequence (e.g., the complementary nucleic acid) can be determined.

**[0053]** The term “nucleic acid” or “polynucleotide” also encompasses any physical string of monomer units that can be corresponded to a string of nucleotides, including a polymer of nucleotides (e.g., a typical DNA or RNA polymer), PNAs, modified oligonucleotides (e.g., oligonucleotides comprising bases that are not typical to biological RNA or DNA in solution, such as 2'-O-methylated oligonucleotides), and the like. A nucleic acid can be e.g., single-stranded or double-stranded.

**[0054]** A “subsequence” is any portion of an entire sequence, up to and including the complete sequence. Typically, a subsequence comprises less than the full-length sequence. A “unique subsequence” is a subsequence that is not found in any previously determined influenza polynucleotide or polypeptide sequence.

**[0055]** The term “variant” with respect to a polypeptide refers to an amino acid sequence that is altered by one or more amino acids with respect to a reference sequence. The variant can have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. Alternatively, a variant can have “nonconservative” changes, e.g., replacement of a glycine with a tryptophan. Analogous minor variation can also include amino acid deletion or insertion, or both. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without eliminating biological or immunological activity can be found using computer programs well known in the art, for example, DNASTAR software. Examples of conservative substitutions are also described herein.

**[0056]** The term “gene” is used broadly to refer to any nucleic acid associated with a biological function. Thus, genes include coding sequences and/or the regulatory sequences required for their expression. The term “gene” applies to a specific genomic sequence, as well as to a cDNA or an mRNA encoded by that genomic sequence.

**[0057]** Genes also include non-expressed nucleic acid segments that, for example, form recognition sequences for other proteins. Non-expressed regulatory sequences include “promoters” and “enhancers,” to which regulatory proteins such as transcription factors bind, resulting in transcription of adjacent or nearby sequences. A “tissue specific” promoter or enhancer is one that regulates transcription in a specific tissue type or cell type, or types.

**[0058]** “Expression of a gene” or “expression of a nucleic acid” means transcription of DNA into RNA (optionally including modification of the RNA, e.g., splicing), translation of RNA into a polypeptide (possibly including subsequent modification of the polypeptide, e.g., post-translational modification), or both transcription and translation, as indicated by the context.

**[0059]** An “open reading frame” or “ORF” is a possible translational reading frame of DNA or RNA (e.g., of a gene), which is capable of being translated into a polypeptide. That is, the reading frame is not interrupted by stop codons. However, it should be noted that the term ORF does not necessarily indicate that the polynucleotide is, in fact, translated into a polypeptide.

**[0060]** The term “vector” refers to the means by which a nucleic acid can be propagated and/or transferred between organisms, cells, or cellular components. Vectors include plasmids, viruses, bacteriophages, pro-viruses, phagemids, transposons, artificial chromosomes, and the like, that replicate autonomously or can integrate into a chromosome of a host cell. A vector can also be a naked RNA polynucleotide, a naked DNA polynucleotide, a polynucleotide composed of both DNA and RNA within the same strand, a poly-lysine-conjugated DNA or RNA, a peptide-conjugated DNA or RNA, a liposome-conjugated DNA, or the like, that is not autonomously replicating. In many, but not all, common embodiments, the vectors of the present invention are plasmids.

**[0061]** An “expression vector” is a vector, such as a plasmid that is capable of promoting expression, as well as replication of a nucleic acid incorporated therein. Typically, the nucleic acid to be expressed is “operably linked” to a promoter and/or enhancer, and is subject to transcription regulatory control by the promoter and/or enhancer.

**[0062]** A “bi-directional expression vector” is characterized by two alternative promoters oriented in the opposite direction relative to a nucleic acid situated between the two promoters, such that expression can be initiated in both orientations resulting in, e.g., transcription of both plus (+) or sense strand, and negative (–) or antisense strand RNAs.

**[0063]** A “polypeptide” is a polymer comprising two or more amino acid residues (e.g., a peptide or a protein). The polymer can optionally comprise modifications such as glycosylation or the like. The amino acid residues of the polypeptide can be natural or non-natural and can be unsubstituted, unmodified, substituted or modified.

**[0064]** In the context of the invention, the term “isolated” refers to a biological material, such as a virus, a nucleic acid or a protein, which is substantially free from components that normally accompany or interact with it in its naturally occurring environment. The isolated biological material optionally comprises additional material not found with the biological material in its natural environment, e.g., a cell or wild-type virus. For example, if the material is in its natural environment, such as a cell, the material can have been placed at a location in the cell (e.g., genome or genetic element) not native to such material found in that environment. For example, a naturally occurring nucleic acid (e.g., a coding sequence, a promoter, an enhancer, etc.) becomes isolated if it is introduced by non-naturally occurring means to a locus of the genome (e.g., a vector, such as a plasmid or virus vector, or amplicon) not native to that nucleic acid. Such nucleic acids are also referred to as “heterologous” nucleic acids. An isolated virus, for example, is in an environment (e.g., a cell culture system, or purified from cell culture) other than the native environment of wild-type virus (e.g., the nasopharynx of an infected individual).

**[0065]** The term “chimeric” or “chimera,” when referring to a virus, indicates that the virus includes genetic and/or polypeptide components derived from more than one parental viral strain or source. Similarly, the term “chimeric” or “chimera,” when referring to a viral protein, indicates that the protein includes polypeptide components (i.e., amino acid subsequences) derived from more than one parental viral strain or source.

**[0066]** The term “recombinant” indicates that the material (e.g., a nucleic acid or protein) has been artificially or

synthetically (non-naturally) altered by human intervention. The alteration can be performed on the material within, or removed from, its natural environment or state. Specifically, e.g., an influenza virus is recombinant when it is produced by the expression of a recombinant nucleic acid. For example, a “recombinant nucleic acid” is one that is made by recombining nucleic acids, e.g., during cloning, DNA shuffling or other procedures, or by chemical or other mutagenesis; a “recombinant polypeptide” or “recombinant protein” is a polypeptide or protein which is produced by expression of a recombinant nucleic acid; and a “recombinant virus”, e.g., a recombinant influenza virus, is produced by the expression of a recombinant nucleic acid.

[0067] The term “reassortant,” when referring to a virus, indicates that the virus includes genetic and/or polypeptide components derived from more than one parental viral strain or source. For example, a 7:1 reassortant includes 7 viral genomic segments (or gene segments) derived from a first parental virus, and a single complementary viral genomic segment, e.g., encoding a hemagglutinin or neuraminidase of the invention. A 6:2 reassortant includes 6 genomic segments, most commonly the 6 internal genes from a first parental virus, and two complementary segments, e.g., hemagglutinin and neuraminidase, from a different parental virus.

[0068] The term “introduced” when referring to a heterologous or isolated nucleic acid refers to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid can be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA). The term includes such methods as “infection,” “transfection,” “transformation” and “transduction.” In the context of the invention a variety of methods can be employed to introduce nucleic acids into cells, including electroporation, calcium phosphate precipitation, lipid mediated transfection (lipofection), etc.

[0069] The term “host cell” means a cell that contains a heterologous nucleic acid, such as a vector, and supports the replication and/or expression of the nucleic acid. Host cells can be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, avian or mammalian cells, including human cells. Exemplary host cells can include, e.g., Vero (African green monkey kidney) cells, BHK (baby hamster kidney) cells, primary chick kidney (PCK) cells, Madin-Darby Canine Kidney (MDCK) cells, Madin-Darby Bovine Kidney (MDBK) cells, 293 cells (e.g., 293T cells), and COS cells (e.g., COS1, COS7 cells), etc.

[0070] An “immunologically effective amount” of influenza virus is an amount sufficient to enhance an individual’s (e.g., a human’s) own immune response against a subsequent exposure to influenza virus. Levels of induced immunity can be monitored, e.g., by measuring amounts of neutralizing secretory and/or serum antibodies, e.g., by plaque neutralization, complement fixation, enzyme-linked immunosorbent, or microneutralization assay.

[0071] A “protective immune response” against influenza virus refers to an immune response exhibited by an individual (e.g., a human) that is protective against disease when the individual is subsequently exposed to and/or infected with such influenza virus. In some instances, the influenza virus (e.g., naturally circulating) can still cause infection, but

it cannot cause a serious infection. Typically, the protective immune response results in detectable levels of host engendered serum and secretory antibodies that are capable of neutralizing virus of the same strain and/or subgroup (and possibly also of a different, non-vaccine strain and/or subgroup) in vitro and in vivo.

[0072] As used herein, an “antibody” is a protein comprising one or more polypeptides substantially or partially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. A typical immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kD) and one “heavy” chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively. Antibodies exist as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab)<sub>2</sub>, a dimer of Fab which itself is a light chain joined to VH-CH1 by a disulfide bond. The F(ab)<sub>2</sub> may be reduced under mild conditions to break the disulfide linkage in the hinge region thereby converting the (Fab)<sub>2</sub> dimer into a Fab’ monomer. The Fab’ monomer is essentially a Fab with part of the hinge region (see, Fundamental Immunology, W. E. Paul, ed., Raven Press, N.Y. (1999), for a more detailed description of other antibody fragments). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such Fab’ fragments may be synthesized de novo either chemically or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein, includes antibodies or fragments either produced by the modification of whole antibodies or synthesized de novo using recombinant DNA methodologies. Antibodies include, e.g., polyclonal antibodies, monoclonal antibodies, multiple or single chain antibodies, including single chain Fv (sFv or scFv) antibodies in which a variable heavy and a variable light chain are joined together (directly or through a peptide linker) to form a continuous polypeptide, and humanized or chimeric antibodies.

#### Influenza Virus

[0073] The polypeptides and polynucleotides of the invention, e.g., SEQ ID NO:1-45, are variants of influenza HA and NA sequences. In general, influenza viruses are made up of an internal ribonucleoprotein core containing a segmented single-stranded RNA genome and an outer lipoprotein envelope lined by a matrix protein. The genome of influenza viruses is composed of eight segments of linear (–) strand ribonucleic acid (RNA), encoding the immunogenic hemagglutinin (HA) and neuraminidase (NA) proteins, and six internal core polypeptides: the nucleocapsid nucleoprotein (NP); matrix proteins (M); non-structural proteins (NS); and

3 RNA polymerase (PA, PB1, PB2) proteins. During replication, the genomic viral RNA is transcribed into (+) strand messenger RNA and (-) strand genomic cRNA in the nucleus of the host cell. Each of the eight genomic segments is packaged into ribonucleoprotein complexes that contain, in addition to the RNA, NP and a polymerase complex (PB1, PB2, and PA).

[0074] Influenza is commonly grouped into influenza A and influenza B categories. Influenza A and influenza B viruses each contain eight segments of single stranded RNA with negative polarity. The influenza A genome encodes eleven polypeptides. Segments 1-3 encode three polypeptides, making up a RNA-dependent RNA polymerase. Segment 1 encodes the polymerase complex protein PB2. The remaining polymerase proteins PB1 and PA are encoded by segment 2 and segment 3, respectively. In addition, segment 1 of some influenza strains encodes a small protein, PB1-F2, produced from an alternative reading frame within the PB1 coding region. Segment 4 encodes the hemagglutinin (HA) surface glycoprotein involved in cell attachment and entry during infection. Segment 5 encodes the nucleocapsid nucleoprotein (NP) polypeptide, the major structural component associated with viral RNA. Segment 6 encodes a neuraminidase (NA) envelope glycoprotein. Segment 7 encodes two matrix proteins, designated M1 and M2, which are translated from differentially spliced mRNAs. Segment 8 encodes NS1 and NS2, two nonstructural proteins, which are translated from alternatively spliced mRNA variants. The eight genome segments of influenza B encode 11 proteins. The three largest genes code for components of the RNA polymerase, PB1, PB2 and PA. Segment 4 encodes the HA protein. Segment 5 encodes NP. Segment 6 encodes the NA protein and the NB protein. Both proteins, NB and NA, are translated from overlapping reading frames of a bicistronic mRNA. Segment 7 of influenza B also encodes two proteins: M1 and BM2. The smallest segment encodes two products: NS1 is translated from the full length RNA, while NS2 is translated from a spliced mRNA variant.

#### Influenza Virus Vaccines

[0075] The sequences, compositions and methods herein are primarily, but not solely, concerned with production of influenza viruses for vaccines. Historically, influenza virus vaccines have primarily been produced in embryonated hen eggs using strains of virus selected or based on empirical predictions of relevant strains. More recently, reassortant viruses have been produced that incorporate selected hemagglutinin and/or neuraminidase antigens in the context of an approved attenuated, temperature sensitive master strain. Following culture of the virus through multiple passages in hen eggs, influenza viruses are recovered and, optionally, inactivated, e.g., using formaldehyde and/or  $\beta$ -propiolactone (or alternatively used in live attenuated vaccines). Thus, it will be appreciated that HA and NA sequences (e.g., SEQ ID NO:1-45) are quite useful in constructing influenza vaccines. The current invention includes viruses/vaccines comprising HA and/or NA sequences of pandemic influenza strains (including wherein the HA sequences comprise modified polybasic cleavage sites such as the modifications described herein); and including wherein the viruses/vaccines comprise a ca backbone such as A/AA/6/60 or the backbone of PR8.

[0076] Attempts at producing recombinant and reassortant vaccines in cell culture have been hampered by the inability

of some of the strains approved for vaccine production to grow efficiently under standard cell culture conditions. However, prior work by the inventors and their coworkers provided a vector system, and methods for producing recombinant and reassortant viruses in culture, thus, making it possible to rapidly produce vaccines corresponding to one or many selected antigenic strains of virus, e.g., either A or B strains, various subtypes or substrains, etc., e.g., comprising the HA and/or NA sequences herein. See, Multi-Plasmid System for the production of Influenza virus, U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003 and U.S. Application 60/574,117 filed May 24, 2004. Typically, the cultures are maintained in a system, such as a cell culture incubator, under controlled humidity and CO<sub>2</sub>, at constant temperature using a temperature regulator, such as a thermostat to insure that the temperature does not exceed 35° C. Reassortant influenza viruses can be readily obtained by introducing a subset of vectors corresponding to genomic segments of a master influenza virus, in combination with complementary segments derived from strains of interest (e.g., HA and/or NA antigenic variants herein). Typically, the master strains are selected on the basis of desirable properties relevant to vaccine administration. For example, for vaccine production, e.g., for production of a live attenuated vaccine, the master donor virus strain may be selected for an attenuated phenotype, cold adaptation and/or temperature sensitivity. As explained elsewhere herein and, e.g., in U.S. patent application Ser. No. 10/423,828, etc., various embodiments of the invention utilize A/Ann Arbor (AA)/6/60 influenza strain as a "backbone" upon which to add HA and/or NA genes (e.g., such as those sequences listed herein, etc.) to create desired reassortant viruses. Thus, for example, in a 6:2 reassortant, 2 genes (i.e., NA and HA) would be from the influenza strain(s) against which an immunogenic reaction is desired, while the other 6 genes would be from the Ann Arbor strain, or other backbone strain, etc. The Ann Arbor virus is useful for its cold adapted, attenuated, temperature sensitive attributes. Of course, it will be appreciated that the HA and NA sequences herein are capable of reassortment with a number of other virus genes or virus types (e.g., a number of different "backbones" such as PR8, etc., containing the other influenza genes present in a reassortant, namely, the non-HA and non-NA genes

[0077] Various embodiments herein can comprise live attenuated vaccines, having the HA and/or NA sequences herein, for pandemic influenza. Such vaccines typically comprise, e.g., the HA and/or NA sequences of SEQ ID NO:11-20, 27-32, or 39-44, or their corresponding encoding nucleotides of SEQ ID NO:1-10, 21-26, 33-38, or 45. One problem arising from growth of vaccine virus strains (e.g., reassortants) in eggs is that avian strains (which can be involved in pandemics) can kill the eggs in which the vaccines are to be produced and are, thus, hard to manipulate, produce, etc. through use of traditional (non-plasmid rescue) reassortant production. Such avian strains are of interest since evidence indicates they can result in influenza in humans and possible pandemics. Thus, use of plasmid-rescue systems to create/manipulate influenza reassortants with pandemic strains such as various avian sequences (e.g., the HA and NA sequences herein) are quite desirable and are features of the invention. It will be appreciated, however, that the current sequences are also capable of use with non-plasmid or traditional systems.

[0078] Aquatic birds (among others) can be infected by influenza A viruses of 15 hemagglutinin (HA) and 9 neuraminidase (NA) subtypes. Such birds can serve as a reservoir from which novel influenza subtypes can be introduced into human populations and cause pandemics. The observation that avian H7N7 influenza A viruses infected humans in The Netherlands in 2003 and avian H5N1 and H9N2 viruses infected humans in Hong Kong and China earlier, raise concerns that these (and other) subtypes have the potential to cause pandemics. Thus, vaccines are needed to prevent human infections with avian influenza A viruses. Live, attenuated influenza A virus vaccines against human influenza viruses were recently licensed in the United States. See above. Such vaccines are reassortant H1N1 and H3N2 viruses in which the internal protein genes of A/Ann Arbor (AA)/6/60 (H2N2) cold adapted (ca) virus confer the cold adapted, attenuation and temperature sensitive phenotypes of the AA ca virus on the reassortant viruses (i.e., the ones having the hemagglutinin and neuraminidase genes from the non-Ann Arbor strain). Classical genetic reassortment and plasmid-based reverse genetics techniques have been applied to generate reassortant viruses that contain the hemagglutinin and neuraminidase genes from avian influenza A viruses (H4-H14 subtypes) and six internal gene segments from the AA ca virus. Such reassortant viruses are features of the invention. See the HA and NA gene sequences below. These viruses bear biological properties that are desirable in candidate vaccines because the phenotypes associated with the AA ca virus are present in the reassortant viruses. The generation and evaluation of these reassortant viruses as seed viruses for vaccines are important steps in pandemic preparedness. It is contemplated that clinical trials can establish the safety, infectivity and immunogenicity of such live attenuated pandemic vaccines. Other embodiments of the invention include reassortant viruses (e.g., those used in vaccines) comprising pandemic antigenic genes HA and/or NA from, e.g., avian, porcine, etc., pandemic virus strains in addition to those listed herein, to produce pandemic vaccines which are created through plasmid-rescue reassortment (e.g., reassortment with A/Ann Arbor 6/60 (i.e., A/AA/6/60), PR8, etc. Methods of construction and use of such viruses and vaccines are also included. "Pandemic virus strains" as used herein is defined as an influenza strain A virus subtype that it is not circulating in the human population, that is declared to be a pandemic strain by the Centers for Disease Control or the World Health Organization or generally acknowledged as such within the scientific community.

[0079] In various embodiments herein, the antigenic sequences (e.g., the HA sequences) as well as viruses and vaccines from such viruses comprise modified polybasic cleavage sites. Some highly pathogenic avian pandemic influenza strains comprise multiple basic amino acid cleavage sites within hemagglutinin sequences. See, e.g., Li et al., *J. of Infectious Diseases*, 179:1132-8, 1999. Such cleavage sites, in typical embodiments herein, are, e.g., modified or altered in their sequences in comparison to the wild-type sequences from which the current sequences are derived (e.g., to disable the cleavage or reduce the cleavage there, etc.). Such modifications/alterations can be different in different strains due to the various sequences of the cleavage sites in the wild-type sequences. For example, 4 polybasic residues (RRKK) at 326-329 of mature H5 are typically removed in sequences herein (as compared to wt). See

"SEQUENCES." In various embodiments, the polybasic cleavage sites can be modified in a number of ways (all of which are contained within the invention). For example, the polybasic cleavage site can be removed one amino acid at a time (e.g., one R removed, two Rs removed, RRR removed, or RRRK removed). Additionally, the amino acid residue directly upstream of the cleavage site can also be removed or altered (e.g., from an R to a T, etc.); also, the nucleotides encoding the amino acid residue directly after the cleavage site can also be modified. See, e.g., FIG. 1 for an illustration of cleavage site modification. In addition, hemagglutinin polypeptide sequences of influenza virus comprise amino terminal signal peptide sequences, thus, the hemagglutinin polypeptide sequences of the invention include both the mature (amino terminal signal peptide cleaved) form of the hemagglutinin polypeptides and the pre-cleaved form of hemagglutinin. The cleavage sites of any hemagglutinin polypeptide sequence of any influenza strain can be routinely measured or predicted using any number of methods in the art.

[0080] The terms "temperature sensitive," "cold adapted" and "attenuated" as applied to viruses (typically used as vaccines or for vaccine production) which optionally encompass the current sequences, are well known in the art. For example, the term "temperature sensitive" (ts) indicates, e.g., that the virus exhibits a 100 fold or greater reduction in titer at 39° C. relative to 33° C. for influenza A strains, or that the virus exhibits a 100 fold or greater reduction in titer at 37° C. relative to 33° C. for influenza B strains. The term "cold adapted" (ca) indicates that the virus exhibits growth at 25° C. within 100 fold of its growth at 33° C., while the term "attenuated" (att) indicates that the virus replicates in the upper airways of ferrets but is not detectable in their lung tissues, and does not cause influenza-like illness in the animal. It will be understood that viruses with intermediate phenotypes, i.e., viruses exhibiting titer reductions less than 100 fold at 39° C. (for A strain viruses) or 37° C. (for B strain viruses), or exhibiting growth at 25° C. that is more than 100 fold than its growth at 33° C. (e.g., within 200 fold, 500 fold, 1000 fold, 10,000 fold less), and/or exhibit reduced growth in the lungs relative to growth in the upper airways of ferrets (i.e., partially attenuated) and/or reduced influenza like illness in the animal, are also useful viruses and can be used in conjunction with the HA and NA sequences herein.

[0081] Again, the HA and NA sequences of the current invention are optionally utilized in the production of or in reassortment vaccines (and/or in other ts, cs, ca, and/or att viruses and vaccines). However, it should be noted that the HA and NA sequences, etc. of the invention are not limited to specific vaccine compositions or production methods, and can, thus, be utilized in substantially any vaccine type or vaccine production method which utilizes strain specific HA and NA antigens (e.g., any of SEQ ID NO:11-20, 27-32, or 39-44 or the corresponding nucleotides encoding the specific HA and NA antigens, e.g., SEQ ID NO:1-10, 21-26, 33-38, or 45).

#### FLUMIST™

[0082] As mentioned previously, numerous examples and types of influenza vaccine exist. An exemplary influenza vaccine is FluMist™ which is a live, attenuated vaccine that protects children and adults from influenza illness (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted,*



trivalent, intranasal influenza virus vaccine in children *N Engl J Med* 338:1405-12; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial JAMA* 282:137-44). In typical, and preferred, embodiments, the methods and compositions of the current invention are preferably adapted to/used with production of FluMist™ vaccine. However, it will be appreciated by those skilled in the art that the sequences, methods, compositions, etc. herein are also adaptable to production of similar or even different viral vaccines.

[0083] FluMist™ vaccine strains contain, e.g., HA and NA gene segments derived from the strains (e.g., wild-type strains) to which the vaccine is addressed along with six gene segments, PB1, PB2, PA, NP, M and NS, from a common master donor virus (MDV). The HA sequences herein, thus, are part of various FluMist™ formulations. The MDV for influenza A strains of FluMist™ (MDV-A), was created by serial passage of the wild-type A/Ann Arbor/6/60 (A/AA/6/60) strain in primary chicken kidney tissue culture at successively lower temperatures (Maassab (1967) *Adaptation and growth characteristics of influenza virus at 25 degrees C. Nature* 213:612-4). MDV-A replicates efficiently at 25° C. (ca. cold adapted), but its growth is restricted at 38 and 39° C. (ts, temperature sensitive). Additionally, this virus does not replicate in the lungs of infected ferrets (att, attenuation). The ts phenotype is believed to contribute to the attenuation of the vaccine in humans by restricting its replication in all but the coolest regions of the respiratory tract. The stability of this property has been demonstrated in animal models and clinical studies. In contrast to the ts phenotype of influenza strains created by chemical mutagenesis, the ts property of MDV-A does not revert following passage through infected hamsters or in shed isolates from children (for a recent review, see Murphy & Coelingh (2002) *Principles underlying the development and use of live attenuated cold-adapted influenza A and B virus vaccines Viral Immunol* 15:295-323).

[0084] Clinical studies in over 20,000 adults and children involving 12 separate 6:2 reassortant strains have shown that these vaccines are attenuated, safe and efficacious (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children N Engl J Med* 338:1405-12; Boyce et al. (2000) *Safety and immunogenicity of adjuvanted and unadjuvanted subunit influenza vaccines administered intranasally to healthy adults Vaccine* 19:217-26; Edwards et al. (1994) *A randomized controlled trial of cold adapted and inactivated vaccines for the prevention of influenza A disease J Infect Dis* 169:68-76; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial JAMA* 282: 137-44). Reassortants carrying the six internal genes of MDV-A and the two HA and NA gene segments of a wild-type virus (i.e., a 6:2 reassortant) consistently maintain ca, ts and att phenotypes (Maassab et al. (1982) *Evaluation of a cold-recombinant influenza virus vaccine in ferrets J. Infect. Dis.* 146:780-900).

[0085] Production of such reassorted virus using B strains of influenza is more difficult, however, recent work (see, e.g., Multi-Plasmid System for the Production of Influenza Virus, U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003,

and U.S. Application No. 60/574,117, filed May 24, 2004) has shown an eight plasmid system for the generation of influenza B virus entirely from cloned cDNA. Methods for the production of attenuated live influenza A and B virus suitable for vaccine formulations, such as live virus vaccine formulations useful for intranasal administration were also shown.

[0086] The system and methods described previously are useful for the rapid production in cell culture of recombinant and reassortant influenza A and B viruses, including viruses suitable for use as vaccines, including live attenuated vaccines, such as vaccines suitable for intranasal administration. The sequences (e.g., nucleotide sequences SEQ ID NO:1-10, 21-26, 33-38, or 45 and the corresponding amino acids encoded by the nucleotide sequences in SEQ ID NO:11-20, 27-32, or 39-44), methods, etc. of the current invention, are optionally used in conjunction with, or in combination with, such previous work involving, e.g., reassorted influenza viruses for vaccine production to produce viruses for vaccines.

#### Methods and Compositions for Prophylactic Administration of Vaccines

[0087] As stated above, alternatively, or in addition to, use in production of FluMist™ vaccine, the current invention can be used in other vaccine formulations. In general, recombinant and reassortant viruses of the invention (e.g., those comprising polynucleotides of SEQ ID NO:1-10, 21, 23-26, 33-38, or 45 or polypeptides of SEQ ID NO:11-20, 27-32, or 39-44, or fragments thereof) can be administered prophylactically in an immunologically effective amount and in an appropriate carrier or excipient to stimulate an immune response specific for one or more strains of influenza virus as determined by the HA and/or NA sequence. Typically, the carrier or excipient is a pharmaceutically acceptable carrier or excipient, such as sterile water, aqueous saline solution, aqueous buffered saline solutions, aqueous dextrose solutions, aqueous glycerol solutions, ethanol, or combinations thereof. The preparation of such solutions insuring sterility, pH, isotonicity, and stability is effected according to protocols established in the art. Generally, a carrier or excipient is selected to minimize allergic and other undesirable effects, and to suit the particular route of administration, e.g., subcutaneous, intramuscular, intranasal, etc.

[0088] A related aspect of the invention provides methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus. In the methods, an immunologically effective amount of a recombinant influenza virus (e.g., comprising an HA and/or NA molecule of the invention), an immunologically effective amount of a polypeptide of the invention, and/or an immunologically effective amount of a nucleic acid of the invention is administered to the individual in a physiologically acceptable carrier.

[0089] Generally, the influenza viruses of the invention are administered in a quantity sufficient to stimulate an immune response specific for one or more strains of influenza virus (i.e., against the HA and/or NA strains of the invention). Preferably, administration of the influenza viruses elicits a protective immune response to such strains. Dosages and methods for eliciting a protective immune response against one or more influenza strains are known to those of skill in the art. See, e.g., U.S. Pat. No. 5,922,326; Wright et al.,

*Infect. Immun.* 37:397-400 (1982); Kim et al., *Pediatrics* 52:56-63 (1973); and Wright et al., *J. Pediatr.* 88:931-936 (1976). For example, influenza viruses are provided in the range of about 1-1000  $\text{HID}_{50}$  (human infectious dose), i.e., about  $10^5$ - $10^8$  pfu (plaque forming units) per dose administered. Typically, the dose will be adjusted within this range based on, e.g., age, physical condition, body weight, sex, diet, time of administration, and other clinical factors. The prophylactic vaccine formulation is systemically administered, e.g., by subcutaneous or intramuscular injection using a needle and syringe, or a needle-less injection device. Alternatively, the vaccine formulation is administered intranasally, either by drops, large particle aerosol (greater than about 10 microns), or spray into the upper respiratory tract. While any of the above routes of delivery results in a protective systemic immune response, intranasal administration confers the added benefit of eliciting mucosal immunity at the site of entry of the influenza virus. For intranasal administration, attenuated live virus vaccines are often preferred, e.g., an attenuated, cold adapted and/or temperature sensitive recombinant or reassortant influenza virus. See above. While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect.

[0090] Typically, the attenuated recombinant influenza of this invention as used in a vaccine is sufficiently attenuated such that symptoms of infection, or at least symptoms of serious infection, will not occur in most individuals immunized (or otherwise infected) with the attenuated influenza virus. In some instances, the attenuated influenza virus can still be capable of producing symptoms of mild illness (e.g., mild upper respiratory illness) and/or of dissemination to unvaccinated individuals. However, its virulence is sufficiently abrogated such that severe lower respiratory tract infections do not occur in the vaccinated or incidental host.

[0091] Alternatively, an immune response can be stimulated by ex vivo or in vivo targeting of dendritic cells with influenza viruses comprising the sequences herein. For example, proliferating dendritic cells are exposed to viruses in a sufficient amount and for a sufficient period of time to permit capture of the influenza antigens by the dendritic cells. The cells are then transferred into a subject to be vaccinated by standard intravenous transplantation methods.

[0092] While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect. In neonates and infants, for example, multiple administrations may be required to elicit sufficient levels of immunity. Administration can continue at intervals throughout childhood, as necessary to maintain sufficient levels of protection against wild-type influenza infection. Similarly, adults who are particularly susceptible to repeated or serious influenza infection, such as, for example, health care workers, day care workers, family members of young children, the elderly, and individuals with compromised cardiopulmonary function may require multiple immunizations to establish and/or maintain protective immune responses. Levels of induced immunity can be monitored, for example, by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to elicit and maintain desired levels of protection.

[0093] Optionally, the formulation for prophylactic administration of the influenza viruses also contains one or more adjuvants for enhancing the immune response to the influenza antigens. Suitable adjuvants include: complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, bacille Calmette-Guerin (BCG), *Corynebacterium parvum*, and the synthetic adjuvant QS-21.

[0094] If desired, prophylactic vaccine administration of influenza viruses can be performed in conjunction with administration of one or more immunostimulatory molecules. Immunostimulatory molecules include various cytokines, lymphokines and chemokines with immunostimulatory, immunopotentiating, and pro-inflammatory activities, such as interleukins (e.g., IL-1, IL-2, IL-3, IL-4, IL-12, IL-13); growth factors (e.g., granulocyte-macrophage (GM)-colony stimulating factor (CSF)); and other immunostimulatory molecules, such as macrophage inflammatory factor, Flt3 ligand, B7.1; B7.2, etc. The immunostimulatory molecules can be administered in the same formulation as the influenza viruses, or can be administered separately. Either the protein (e.g., an HA and/or NA polypeptide of the invention, e.g., any of SEQ ID NO:11-20, 27-32, or 39-44) or an expression vector comprising a nucleic acid (e.g., any of SEQ ID NO:1-10, 21-26, 33-38, or 45) encoding the protein can be administered to produce an immunostimulatory effect.

[0095] The above described methods are useful for therapeutically and/or prophylactically treating a disease or disorder, typically influenza, by introducing a vector of the invention comprising a heterologous polynucleotide encoding a therapeutically or prophylactically effective HA and/or NA polypeptide (or peptide) or HA and/or NA RNA (e.g., an antisense RNA or ribozyme) into a population of target cells in vitro, ex vivo or in vivo. Typically, the polynucleotide encoding the polypeptide (or peptide), or RNA, of interest is operably linked to appropriate regulatory sequences as described above in the sections entitled "Expression Vectors" and "Additional Expression Elements." Optionally, more than one heterologous coding sequence is incorporated into a single vector or virus. For example, in addition to a polynucleotide encoding a therapeutically or prophylactically active HA and/or NA polypeptide or RNA, the vector can also include additional therapeutic or prophylactic polypeptides, e.g., antigens, co-stimulatory molecules, cytokines, antibodies, etc., and/or markers, and the like.

[0096] Although vaccination of an individual with an attenuated influenza virus of a particular strain of a particular subgroup can induce cross-protection against influenza virus of different strains and/or subgroups, cross-protection can be enhanced, if desired, by vaccinating the individual with attenuated influenza virus from at least two strains, e.g., each of which represents a different subgroup. Additionally, vaccine combinations can optionally include mixes of pandemic vaccines (e.g., those against pandemic influenza strains such as various avian strains, see, e.g., the sequences herein, or other pandemic strains) and non-pandemic strains. Vaccine mixtures (or multiple vaccinations) can comprise components from human strains and/or non-human influenza strains (e.g., avian and human, etc.). Similarly, the attenuated influenza virus vaccines of this invention can

optionally be combined with vaccines that induce protective immune responses against other infectious agents.

#### Polynucleotides of the Invention

**[0097]** It is well known in the art that the HA and NA polynucleotide segments of influenza viruses comprise both a coding region (encoding the ORF) and noncoding regions (NCRs), both 5' and 3' of the HA and NA coding sequence. An example of these NCRs are shown in SEQ ID NOS:1-9 (outside of the ORFs). It is also known that primers can be made to these NCRs to facilitate amplification of the entire HA and NA segments of influenza virus. (see, e.g., Hoffmann et al. Arch Virol. 2001 December; 146(12):2275-89). Further, it is known that the NCRs of the HA and NA of influenza may increase the efficiency of achieving reassortants. Therefore, the polynucleotide sequences of these NCRs (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9% identity) thereof) are within the scope of this invention. When amplifying the HA and NA segments of any pandemic strain, one could make and use polynucleotide primers to bind conserved (e.g., among related strains) regions of the HA and NA NCRs for amplification (e.g., by RT-PCR). In one embodiment, HA and NA polynucleotides of the invention include both the NCR and ORF of the HA and NA sequences (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9%) thereof) of pandemic virus strains. In alternative embodiments, the HA and NA polynucleotides of the invention exclude the NCR, but include the ORF (including fragments and variants (e.g., at least about 60%, or at least 70%, or at least 80%, or at least 90%, or at least about 91% or at least about 92%, or at least about 93%, or at least about 94%, or at least about 95%, or at least about 96%, or at least about 97%, or at least about 98%, or at least about 98.5%, or at least about 98.7%, or at least about 99%, or at least about 99.1%, or at least about 99.2%, or at least about 99.3%, or at least about 99.4%, or at least about 99.5%, or at least about 99.6% or at least about 99.7%, or at least about 99.8%, or at least about 99.9% thereof)) of the HA and NA sequences of pandemic virus strains (e.g., SEQ ID NOS:1-9).

**[0098]** The HA and NA polynucleotides of the invention, e.g., SEQ ID NO:1 through SEQ ID NO:10, SEQ ID NO:21 through SEQ ID NO:26, SEQ ID NO:33 through SEQ ID NO:38, SEQ ID NO:45, and fragments thereof, are optionally used in a number of different capacities alternative to, or in addition to, the vaccines described above. Other exemplary uses are described herein for illustrative purpose

and not as limitations on the actual range of uses. Different methods of construction, purification, and characterization of the nucleotide sequences of the invention are also described herein. In some embodiments, nucleic acids including one or more polynucleotide sequence of the invention are favorably used as probes for the detection of corresponding or related nucleic acids in a variety of contexts, such as in nucleic hybridization experiments, e.g., to find and/or characterize homologous influenza variants (e.g., homologues to the sequences herein, etc.) infecting other species or in different influenza outbreaks, etc. The probes can be either DNA or RNA molecules, such as restriction fragments of genomic or cloned DNA, cDNAs, PCR amplification products, transcripts, and oligonucleotides, and can vary in length from oligonucleotides as short as about 10 nucleotides in length to full length sequences or cDNAs in excess of 1 kb or more. For example, in some embodiments, a probe of the invention includes a polynucleotide sequence or subsequence selected, e.g., from among SEQ ID NO:1 through SEQ ID NO:10, SEQ ID NO:21 through SEQ ID NO:26, SEQ ID NO:33 through SEQ ID NO:38, SEQ ID NO:45, or sequences complementary thereto. Alternatively, polynucleotide sequences that are variants of one of the above-designated sequences are used as probes. Most typically, such variants include one or a few conservative nucleotide variations. For example, pairs (or sets) of oligonucleotides can be selected, in which the two (or more) polynucleotide sequences are conservative variations of each other, wherein one polynucleotide sequence corresponds identically to a first variant or and the other(s) corresponds identically to additional variants. Such pairs of oligonucleotide probes are particularly useful, e.g., for specific hybridization experiments to detect polymorphic nucleotides or to, e.g., detect homologous influenza HA and NA variants, e.g., homologous to the current HA and NA sequences, infecting other species or present in different (e.g., either temporally and/or geographically different) influenza outbreaks. In other applications, probes are selected that are more divergent, that is probes that are at least about 91% (or about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 98.5%, about 98.7%, about 99%, about 99.1%, about 99.2%, about 99.3%, about 99.4%, about 99.5%, or about 99.6% or more about 99.7%, about 99.8%, about 99.9% or more) identical are selected.

**[0099]** The probes of the invention, e.g., as exemplified by sequences derived from the sequences herein, can also be used to identify additional useful polynucleotide sequences according to procedures routine in the art. In one set of embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or chromosomal segments (e.g., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence similarity to, e.g., one or more probe of the sequences herein, i.e., variants, homologues, etc. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences. Polynucleotide sequences identified in this manner are also a feature of the invention.

**[0100]** Oligonucleotide probes are optionally produced via a variety of methods well known to those skilled in the art.

Most typically, they are produced by well known synthetic methods, such as the solid phase phosphoramidite triester method described by Beaucage and Caruthers (1981) *Tetrahedron Letts* 22(20):1859-1862, e.g., using an automated synthesizer, or as described in Needham-Van Devanter et al. (1984) *Nucl Acids Res*, 12:6159-6168. Oligonucleotides can also be custom made and ordered from a variety of commercial sources known to persons of skill. Purification of oligonucleotides, where necessary, is typically performed by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) *J Chrom* 255:137-149. The sequence of the synthetic oligonucleotides can be verified using the chemical degradation method of Maxam and Gilbert (1980) in Grossman and Moldave (eds.) Academic Press, New York, *Methods in Enzymology* 65:499-560. Custom oligos can also easily be ordered from a variety of commercial sources known to persons of skill.

[0101] In other circumstances, e.g., relating to attributes of cells or organisms expressing the polynucleotides and polypeptides of the invention (e.g., those harboring virus comprising the sequences of the invention), probes that are polypeptides, peptides or antibodies are favorably utilized. For example, isolated or recombinant polypeptides, polypeptide fragments and peptides derived from any of the amino acid sequences of the invention (e.g., SEQ ID NO:11-20, SEQ ID NO: 27-32, SEQ ID NO:39-44) and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1 through SEQ ID NO:10, SEQ ID NO: 21 through SEQ ID NO: 26, SEQ ID NO:33 through SEQ ID NO:38, and SEQ ID NO:45 are favorably used to identify and isolate antibodies, e.g., from phage display libraries, combinatorial libraries, polyclonal sera, and the like. Polypeptide fragments of the inventions include a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, or at least 10 contiguous amino acid residues, or at least 15 contiguous amino acid residues, or at least 20 contiguous amino acid residues, or at least 25 contiguous amino acid residues, or at least 40 contiguous amino acid residues, or at least 50 contiguous amino acid residues, or at least 60 contiguous amino residues, or at least 70 contiguous amino acid residues, or at least contiguous 80 amino acid residues, or at least contiguous 90 amino acid residues, or at least contiguous 100 amino acid residues, or at least contiguous 125 amino acid residues, or at least 150 contiguous amino acid residues, or at least contiguous 175 amino acid residues, or at least contiguous 200 amino acid residues, or at least contiguous 250 amino acid residues, or at least contiguous 350, or at least contiguous 400, or at least contiguous 450, or at least contiguous 500, or at least contiguous 550 amino acid residues of the amino acid sequence an HA or NA polypeptide of the invention (e.g., SEQ ID NOS:11-20, SEQ ID NOS: 27-32, and SEQ ID NOS: 39-44). Polynucleotides encoding said polypeptide fragments and antibodies that specifically bind said polypeptides are also preferred embodiments of the invention.

[0102] Antibodies specific for any polypeptide sequence or subsequence, e.g., of SEQ ID NO:11 through SEQ ID NO: 20, SEQ ID NO: 27 through SEQ ID NO: 32, and/or SEQ ID NO: 39 through SEQ ID NO: 44, and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1 through SEQ ID NO:10, SEQ ID NO: 21 through SEQ ID NO: 26, SEQ ID NO: 33 through SEQ

ID NO: 38, and SEQ ID NO:45 are likewise valuable as probes for evaluating expression products, e.g., from cells or tissues. In addition, antibodies are particularly suitable for evaluating expression of proteins comprising amino acid subsequences, e.g., of those given herein, or encoded by polynucleotides sequences of the invention, e.g., selected from those shown herein, in situ, in a tissue array, in a cell, tissue or organism, e.g., an organism infected by an unidentified influenza virus or the like. Antibodies can be directly labeled with a detectable reagent, or detected indirectly by labeling of a secondary antibody specific for the heavy chain constant region (i.e., isotype) of the specific antibody. Additional details regarding production of specific antibodies are provided below.

#### [0103] Diagnostic Assays

[0104] The nucleic acid sequences of the present invention can be used in diagnostic assays to detect influenza (and/or hemagglutinin and/or neuraminidase) in a sample, to detect hemagglutinin-like and/or neuraminidase-like sequences, and to detect strain differences in clinical isolates of influenza using either chemically synthesized or recombinant polynucleotide fragments, e.g., selected from the sequences herein. For example, fragments of the hemagglutinin and/or neuraminidase sequences comprising at least between 10 and 20 nucleotides can be used as primers to amplify nucleic acids using polymerase chain reaction (PCR) methods well known in the art (e.g., reverse transcription-PCR) and as probes in nucleic acid hybridization assays to detect target genetic material such as influenza RNA in clinical specimens.

[0105] The probes of the invention, e.g., as exemplified by unique subsequences selected from those given herein, can also be used to identify additional useful polynucleotide sequences (such as to characterize additional strains of influenza) according to procedures routine in the art. In one set of preferred embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or cloned viral nucleic acids (i.e., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence identity to the sequences herein. In turn, each of these identified sequences can be used to make probes, including pairs or sets of variant probes as described above. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences.

[0106] The probes of the invention are particularly useful for detecting the presence and for determining the identity of influenza nucleic acids in cells, tissues or other biological samples (e.g., a nasal wash or bronchial lavage). For example, the probes of the invention are favorably utilized to determine whether a biological sample, such as a subject (e.g., a human subject) or model system (such as a cultured cell sample) has been exposed to, or become infected with influenza, or particular strain(s) of influenza. Detection of hybridization of the selected probe to nucleic acids originating in (e.g., isolated from) the biological sample or model system is indicative of exposure to or infection with the virus (or a related virus) from which the probe polynucleotide is selected.

[0107] It will be appreciated that probe design is influenced by the intended application. For example, where several allele-specific probe-target interactions are to be detected in a single assay, e.g., on a single DNA chip, it is desirable to have similar melting temperatures for all of the probes. Accordingly, the lengths of the probes are adjusted so that the melting temperatures for all of the probes on the array are closely similar (it will be appreciated that different lengths for different probes may be needed to achieve a particular  $T_m$  where different probes have different GC contents). Although melting temperature is a primary consideration in probe design, other factors are optionally used to further adjust probe construction, such as selecting against primer self-complementarity and the like.

[0108] Vectors, Promoters and Expression Systems

[0109] The present invention includes recombinant constructs incorporating one or more of the nucleic acid sequences described herein. Such constructs optionally include a vector, for example, a plasmid, a cosmid, a phage, a virus, a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), etc., into which one or more of the polynucleotide sequences of the invention, e.g., comprising any of SEQ ID NO:1 through SEQ ID NO:10, SEQ ID NO:21 through SEQ ID NO:26, SEQ ID NO:33 through SEQ ID NO:38, SEQ ID NO:45 or a subsequence thereof etc., has been inserted, in a forward or reverse orientation. For example, the inserted nucleic acid can include a viral chromosomal sequence or cDNA including all or part of at least one of the polynucleotide sequences of the invention. In one embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

[0110] The polynucleotides of the present invention can be included in any one of a variety of vectors suitable for generating sense or antisense RNA, and optionally, polypeptide (or peptide) expression products (e.g., a hemagglutinin and/or neuraminidase molecule of the invention, or fragments thereof). Such vectors include chromosomal, non-chromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, pseudorabies, adenovirus, adeno-associated virus, retroviruses and many others (e.g., pCDL). Any vector that is capable of introducing genetic material into a cell, and, if replication is desired, which is replicable in the relevant host can be used.

[0111] In an expression vector, the HA and/or NA polynucleotide sequence of interest is physically arranged in proximity and orientation to an appropriate transcription control sequence (e.g., promoter, and optionally, one or more enhancers) to direct mRNA synthesis. That is, the polynucleotide sequence of interest is operably linked to an appropriate transcription control sequence. Examples of such promoters include: LTR or SV40 promoter, *E. coli* lac or trp promoter, phage lambda  $P_L$  promoter, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

[0112] A variety of promoters are suitable for use in expression vectors for regulating transcription of influenza

virus genome segments. In certain embodiments, the cytomegalovirus (CMV) DNA dependent RNA Polymerase II (Pol II) promoter is utilized. If desired, e.g., for regulating conditional expression, other promoters can be substituted which induce RNA transcription under the specified conditions, or in the specified tissues or cells. Numerous viral and mammalian, e.g., human promoters are available, or can be isolated according to the specific application contemplated. For example, alternative promoters obtained from the genomes of animal and human viruses include such promoters as the adenovirus (such as Adenovirus 2), papilloma virus, hepatitis-B virus, polyoma virus, and Simian Virus 40 (SV40), and various retroviral promoters. Mammalian promoters include, among many others, the actin promoter, immunoglobulin promoters, heat-shock promoters, and the like.

[0113] Transcription is optionally increased by including an enhancer sequence. Enhancers are typically short, e.g., 10-500 bp, cis-acting DNA elements that act in concert with a promoter to increase transcription. Many enhancer sequences have been isolated from mammalian genes (hemoglobin, elastase, albumin, alpha-fetoprotein, and insulin), and eukaryotic cell viruses. The enhancer can be spliced into the vector at a position 5' or 3' to the heterologous coding sequence, but is typically inserted at a site 5' to the promoter. Typically, the promoter, and if desired, additional transcription enhancing sequences are chosen to optimize expression in the host cell type into which the heterologous DNA is to be introduced (Scharf et al. (1994) *Heat stress promoters and transcription factors Results Probl Cell Differ* 20:125-62; Kriegler et al. (1990) *Assembly of enhancers, promoters, and splice signals to control expression of transferred genes Methods in Enzymol* 185: 512-27). Optionally, the amplicon can also contain a ribosome binding site or an internal ribosome entry site (IRES) for translation initiation.

[0114] The vectors of the invention also favorably include sequences necessary for the termination of transcription and for stabilizing the mRNA, such as a polyadenylation site or a terminator sequence. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. In one embodiment, the SV40 polyadenylation signal sequences can provide a bi-directional polyadenylation site that insulates transcription of (+) strand mRNA molecules from the Poll promoter initiating replication of the (-) strand viral genome.

[0115] In addition, as described above, the expression vectors optionally include one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells, in addition to genes previously listed, markers such as dihydrofolate reductase or neomycin resistance are suitable for selection in eukaryotic cell culture.

[0116] The vector containing the appropriate nucleic acid sequence as described above, as well as an appropriate promoter or control sequence, can be employed to transform a host cell permitting expression of the protein. While the vectors of the invention can be replicated in bacterial cells, most frequently it will be desirable to introduce them into mammalian cells, e.g., Vero cells, BHK cells, MDCK cell, 293 cells, COS cells, or the like, for the purpose of expression.

[0117] As described elsewhere, the HA and NA sequences herein, in various embodiments, can be comprised within

plasmids involved in plasmid-rescue reassortment. See, e.g., U.S. Application Nos. 60/420,708, filed Oct. 23, 2002; 60/574,117, filed May 24, 2004; Ser. No. 10/423,828, filed Apr. 25, 2003; 60/578,962, filed Jun. 12, 2004; and Ser. No. 10/870,690 filed Jun. 16, 2004; and US20020164770, which are incorporated by reference herein. For example, preferred expression vectors of the invention include, but are not limited to, vectors comprising pol I promoter and terminator sequences or vectors using both the pol I and pol II promoters "the poll/pollII promoter system" (e.g., Zobel et al., Nucl. Acids Res. 1993, 21:3607; US20020164770; Neumann et al., Proc. Natl. Acad. Sci. USA 1999, 96:9345; Fodor et al., J. Virol. 1999, 73:9679; and US20030035814). The reassortants produced can include the HA and NA genes arranged with the 6 other influenza genes from the A/Ann Arbor/6/60 donor strain (and/or derivatives and modifications thereof), the PR8 donor strain backbone, the A/Leningrad/17 donor strain backbone, etc. Other backbone strains are described, for example, in 20040137013 and 20030147916, which are incorporated by reference herein.

#### [0118] Additional Expression Elements

[0119] Most commonly, the genome segment encoding the influenza virus HA and/or NA protein includes any additional sequences necessary for its expression, including translation into a functional viral protein. In other situations, a minigene, or other artificial construct encoding the viral proteins, e.g., an HA and/or NA protein, can be employed. Again, in such case, it is often desirable to include specific initiation signals that aid in the efficient translation of the heterologous coding sequence. These signals can include, e.g., the ATG initiation codon and adjacent sequences. To insure translation of the entire insert, the initiation codon is inserted in the correct reading frame relative to the viral protein. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

[0120] If desired, polynucleotide sequences encoding additional expressed elements, such as signal sequences, secretion or localization sequences, and the like can be incorporated into the vector, usually, in-frame with the polynucleotide sequence of interest, e.g., to target polypeptide expression to a desired cellular compartment, membrane, or organelle, or to direct polypeptide secretion to the periplasmic space or into the cell culture media. Such sequences are known to those of skill, and include secretion leader peptides, organelle targeting sequences (e.g., nuclear localization sequences, ER retention signals, mitochondrial transit sequences), membrane localization/anchor sequences (e.g., stop transfer sequences, GPI anchor sequences), and the like.

[0121] Where translation of a polypeptide encoded by a nucleic acid sequence of the invention is desired, additional translation specific initiation signals can improve the efficiency of translation. These signals can include, e.g., an ATG initiation codon and adjacent sequences, an IRES region, etc. In some cases, for example, full-length cDNA molecules or chromosomal segments including a coding sequence incorporating, e.g., a polynucleotide sequence of the invention (e.g., as in the sequences herein), a translation initiation codon and associated sequence elements are inserted into the appropriate expression vector simultaneously with the poly-

nucleotide sequence of interest. In such cases, additional translational control signals frequently are not required. However, in cases where only a polypeptide coding sequence, or a portion thereof, is inserted, exogenous translational control signals, including, e.g., an ATG initiation codon is often provided for expression of the relevant sequence. The initiation codon is put in the correct reading frame to ensure transcription of the polynucleotide sequence of interest. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, e.g., Scharf D. et al. (1994) *Results Probl Cell Differ* 20:125-62; Bittner et al. (1987) *Methods in Enzymol* 153:516-544).

#### [0122] Production of Recombinant Virus

[0123] Negative strand RNA viruses can be genetically engineered and recovered using a recombinant reverse genetics approach (see, e.g., U.S. Pat. No. 5,166,057 to Palese et al.). Such method was originally applied to engineer influenza viral genomes (Luytjes et al. (1989) *Cell* 59:1107-1113; Enami et al. (1990) *Proc. Natl. Acad. Sci. USA* 92:11563-11567), and has been successfully applied to a wide variety of segmented and nonsegmented negative strand RNA viruses, e.g., rabies (Schnell et al. (1994) *EMBO J.* 13: 4195-4203); VSV (Lawson et al. (1995) *Proc. Natl. Acad. Sci. USA* 92: 4477-4481); measles virus (Radecke et al. (1995) *EMBO J.* 14:5773-5784); rinderpest virus (Baron & Barrett (1997) *J. Virol.* 71: 1265-1271); human parainfluenza virus (Hoffman & Banerjee (1997) *J. Virol.* 71: 3272-3277; Dubin et al. (1997) *Virology* 235:323-332); SV5 (He et al. (1997) *Virology* 237:249-260); canine distemper virus (Gassen et al. (2000) *J. Virol.* 74:10737-44); and Sendai virus (Park et al. (1991) *Proc. Natl. Acad. Sci. USA* 88: 5537-5541; Kato et al. (1996) *Genes to Cells* 1:569-579). Those of skill in the art will be familiar with these and similar techniques to produce influenza virus comprising the HA and NA sequences of the invention. Recombinant influenza viruses produced according to such methods are also a feature of the invention, as are recombinant influenza virus comprising one or more nucleic acids and/or polypeptides of the invention.

#### [0124] Cell Culture and Expression Hosts

[0125] The present invention also relates to host cells that are introduced (transduced, transformed or transfected) with vectors of the invention, and the production of polypeptides of the invention by recombinant techniques. Host cells are genetically engineered (i.e., transduced, transformed or transfected) with a vector, such as an expression vector, of this invention. As described above, the vector can be in the form of a plasmid, a viral particle, a phage, etc. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells, such as *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Neurospora crassa*; or insect cells such as *Drosophila* and *Spodoptera frugiperda*.

[0126] Most commonly, mammalian cells are used to culture the HA and NA molecules of the invention. Suitable host cells for the replication of influenza virus include, e.g., Vero cells, BHK cells, MDCK cells, 293 cells and COS cells, including 293T cells, COS7 cells or the like. Commonly, co-cultures including two of the above cell lines, e.g.,

MDCK cells and either 293T or COS cells are employed at a ratio, e.g., of 1:1, to improve replication efficiency. Typically, cells are cultured in a standard commercial culture medium, such as Dulbecco's modified Eagle's medium supplemented with serum (e.g., 10% fetal bovine serum), or in serum free medium, under controlled humidity and CO<sub>2</sub> concentration suitable for maintaining neutral buffered pH (e.g., at pH between 7.0 and 7.2). Optionally, the medium contains antibiotics to prevent bacterial growth, e.g., penicillin, streptomycin, etc., and/or additional nutrients, such as L-glutamine, sodium pyruvate, non-essential amino acids, additional supplements to promote favorable growth characteristics, e.g., trypsin,  $\beta$ -mercaptoethanol, and the like.

[0127] The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the inserted polynucleotide sequences. The culture conditions, such as temperature, pH and the like, are typically those previously used with the particular host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, e.g., Freshney (1994) *Culture of Animal Cells, a Manual of Basic Technique*, 3<sup>rd</sup> edition, Wiley-Liss, New York and the references cited therein. Other helpful references include, e.g., Paul (1975) *Cell and Tissue Culture*, 5<sup>th</sup> ed., Livingston, Edinburgh; Adams (1980) *Laboratory Techniques in Biochemistry and Molecular Biology-Cell Culture for Biochemists*, Work and Burdon (eds.) Elsevier, Amsterdam. Additional details regarding tissue culture procedures of particular interest in the production of influenza virus in vitro include, e.g., Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*, in Cohen and Shaf-ferman (eds.) *Novel Strategies in Design and Production of Vaccines*, which is incorporated herein in its entirety for all purposes. Additionally, variations in such procedures adapted to the present invention are readily determined through routine experimentation and will be familiar to those skilled in the art.

[0128] Cells for production of influenza virus (e.g., having the HA and/or NA sequences of the invention) can be cultured in serum-containing or serum free medium. In some cases, e.g., for the preparation of purified viruses, it is typically desirable to grow the host cells in serum free conditions. Cells can be cultured in small scale, e.g., less than 25 ml medium, culture tubes or flasks or in large flasks with agitation, in rotator bottles, or on microcarrier beads (e.g., DEAE-Dextran microcarrier beads, such as Dorma-cell, Pfeifer & Langen; Superbead, Flow Laboratories; styrene copolymer-tri-methylamine beads, such as Hillex, SoloHill, Ann Arbor) in flasks, bottles or reactor cultures. Microcarrier beads are small spheres (in the range of 100-200 microns in diameter) that provide a large surface area for adherent cell growth per volume of cell culture. For example a single liter of medium can include more than 20 million microcarrier beads providing greater than 8000 square centimeters of growth surface. For commercial production of viruses, e.g., for vaccine production, it is often desirable to culture the cells in a bioreactor or fermenter. Bioreactors are available in volumes from under 1 liter to in excess of 100 liters, e.g., Cyto3 Bioreactor (Osmonics, Minnetonka, Minn.); NBS bioreactors (New Brunswick Scientific, Edison, N.J.); laboratory and commercial scale bioreactors from B. Braun Biotech International (B. Braun Biotech, Melsungen, Germany).

[0129] Regardless of the culture volume, in many desired aspects of the current invention, it is important that the cultures be maintained at an appropriate temperature, to insure efficient recovery of recombinant and/or reassortant influenza virus using temperature dependent multi plasmid systems (see, e.g., Multi-Plasmid System for the Production of Influenza Virus, U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 24, 2004), heating of virus solutions for filtration, etc. Typically, a regulator, e.g., a thermostat, or other device for sensing and maintaining the temperature of the cell culture system and/or other solution, is employed to insure that the temperature is at the correct level during the appropriate period (e.g., virus replication, etc.).

[0130] In some embodiments herein (e.g., wherein reassorted viruses are to be produced from segments on vectors) vectors comprising influenza genome segments are introduced (e.g., transfected) into host cells according to methods well known in the art for introducing heterologous nucleic acids into eukaryotic cells, including, e.g., calcium phosphate co-precipitation, electroporation, microinjection, lipofection, and transfection employing polyamine transfection reagents. For example, vectors, e.g., plasmids, can be transfected into host cells, such as COS cells, 293T cells or combinations of COS or 293T cells and MDCK cells, using the polyamine transfection reagent TransIT-LT1 (Mirus) according to the manufacturer's instructions in order to produce reassorted viruses, etc. Thus, in one example, approximately 1  $\mu$ g of each vector is introduced into a population of host cells with approximately 2  $\mu$ l of TransIT-LT1 diluted in 160  $\mu$ l medium, preferably serum-free medium, in a total volume of 200  $\mu$ l. The DNA:transfection reagent mixtures are incubated at room temperature for 45 minutes followed by addition of 800  $\mu$ l of medium. The transfection mixture is added to the host cells, and the cells are cultured as described via other methods well known to those skilled in the art. Accordingly, for the production of recombinant or reassortant viruses in cell culture, vectors incorporating each of the 8 genome segments, (PB2, PB1, PA, NP, M, NS, HA and NA, e.g., of the invention) are mixed with approximately 20  $\mu$ l TransIT-LT1 and transfected into host cells. Optionally, serum-containing medium is replaced prior to transfection with serum-free medium, e.g., Opti-MEM I, and incubated for 4-6 hours.

[0131] Alternatively, electroporation can be employed to introduce such vectors incorporating influenza genome segments into host cells. For example, plasmid vectors incorporating an influenza A or influenza B virus are favorably introduced into Vero cells using electroporation according to the following procedure. In brief, approximately  $5 \times 10^6$  Vero cells, e.g., grown in Modified Eagle's Medium (MEM) supplemented with 10% Fetal Bovine Serum (FBS) are resuspended in 0.4 ml OptiMEM and placed in an electroporation cuvette. Twenty micrograms of DNA in a volume of up to 25  $\mu$ l is added to the cells in the cuvette, which is then mixed gently by tapping. Electroporation is performed according to the manufacturer's instructions (e.g., BioRad Gene Pulser II with Capacitance Extender Plus connected) at 300 volts, 950 microFarads with a time constant of between 28-33 msec. The cells are remixed by gently tapping and approximately 1-2 minutes following electroporation 0.7 ml MEM with 10% FBS is added directly to the cuvette. The cells are then transferred to two wells of a standard 6 well



tissue culture dish containing 2 ml MEM, 10% FBS. The cuvette is washed to recover any remaining cells and the wash suspension is divided between the two wells. Final volume is approximately 3.5 mL. The cells are then incubated under conditions permissive for viral growth, e.g., at approximately 33° C. for cold adapted strains.

**[0132]** In mammalian host cells, a number of expression systems, such as viral-based systems, can be utilized. In cases where an adenovirus is used as an expression vector, a coding sequence is optionally ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome will result in a viable virus capable of expressing the polypeptides of interest in infected host cells (Logan and Shenk (1984) *Proc Natl Acad Sci* 81:3655-3659). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

**[0133]** A host cell strain is optionally chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the protein include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation. Post-translational processing, which cleaves a precursor form into a mature form, of the protein is sometimes important for correct insertion, folding and/or function. Additionally proper location within a host cell (e.g., on the cell surface) is also important. Different host cells such as COS, CHO, BHK, MDCK, 293, 293T, COS7, etc. have specific cellular machinery and characteristic mechanisms for such post-translational activities and can be chosen to ensure the correct modification and processing of the current introduced, foreign protein.

**[0134]** For long-term, high-yield production of recombinant proteins encoded by, or having subsequences encoded by, the polynucleotides of the invention, stable expression systems are optionally used. For example, cell lines, stably expressing a polypeptide of the invention, are transfected using expression vectors that contain viral origins of replication or endogenous expression elements and a selectable marker gene. For example, following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Thus, resistant clumps of stably transformed cells, e.g., derived from single cell type, can be proliferated using tissue culture techniques appropriate to the cell type.

**[0135]** Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The cells expressing said protein can be sorted, isolated and/or purified. The protein or fragment thereof produced by a recombinant cell can be secreted, membrane-bound, or retained intracellularly, depending on the sequence (e.g., depending upon fusion proteins encoding a membrane retention signal or the like) and/or the vector used.

**[0136]** Expression products corresponding to the nucleic acids of the invention can also be produced in non-animal cells such as plants, yeast, fungi, bacteria and the like. In

addition to Sambrook, Berger and Ausubel, all infra, details regarding cell culture can be found in Payne et al. (1992) *Plant Cell and Tissue Culture in Liquid Systems* John Wiley & Sons, Inc. New York, N.Y.; Gamborg and Phillips (eds.) (1995) *Plant Cell, Tissue and Organ Culture*; Fundamental Methods Springer Lab Manual, Springer-Verlag (Berlin Heidelberg New York) and Atlas and Parks (eds.) *The Handbook of Microbiological Media* (1993) CRC Press, Boca Raton, Fla.

**[0137]** In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the expressed product. For example, when large quantities of a polypeptide or fragments thereof are needed for the production of antibodies, vectors that direct high-level expression of fusion proteins that are readily purified are favorably employed. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the coding sequence of interest, e.g., sequences comprising those found herein, etc., can be ligated into the vector in-frame with sequences for the amino-terminal translation initiating methionine and the subsequent 7 residues of beta-galactosidase producing a catalytically active beta galactosidase fusion protein; pIN vectors (Van Heeke & Schuster (1989) *J Biol Chem* 264:5503-5509); pET vectors (Novagen, Madison Wis.); and the like. Similarly, in the yeast *Saccharomyces cerevisiae* a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH can be used for production of the desired expression products. For reviews, see Ausubel, infra, and Grant et al., (1987); *Methods in Enzymology* 153:516-544.

#### **[0138] Nucleic Acid Hybridization**

**[0139]** Comparative hybridization can be used to identify nucleic acids (e.g., SEQ ID NO:1-10, SEQ ID NO: 21-26, SEQ ID NO:33-38, SEQ ID NO:45) of the invention, including conservative variations of nucleic acids of the invention. This comparative hybridization method is a preferred method of distinguishing nucleic acids of the invention. In addition, target nucleic acids which hybridize to the nucleic acids represented by, e.g., those shown herein under high, ultra-high and ultra-ultra-high stringency conditions are features of the invention. Examples of such nucleic acids include those with one or a few silent or conservative nucleic acid substitutions as compared to a given nucleic acid sequence.

**[0140]** A test target nucleic acid is said to specifically hybridize to a probe nucleic acid when it hybridizes at least one-half as well to the probe as to the perfectly matched complementary target, i.e., with a signal to noise ratio at least one-half as high as hybridization of the probe and target under conditions in which a perfectly matched probe binds to a perfectly matched complementary target with a signal to noise ratio that is at least about 5×-10× as high as that observed for hybridization to any of the unmatched target nucleic acids.

**[0141]** Nucleic acids "hybridize" when they associate, typically in solution. Nucleic acids hybridize due to a variety of well-characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. Numerous protocols for nucleic acid hybridization are well known in the art. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory*



*Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes* part I chapter 2, “Overview of principles of hybridization and the strategy of nucleic acid probe assays,” (Elsevier, New York), as well as in Ausubel, Sambrook, and Berger and Kimmel, all below. Hames and Higgins (1995) *Gene Probes* 1 IRL Press at Oxford University Press, Oxford, England, (Hames and Higgins 1) and Hames and Higgins (1995) *Gene Probes* 2 IRL Press at Oxford University Press, Oxford, England (Hames and Higgins 2) provide details on the synthesis, labeling, detection and quantification of DNA and RNA, including oligonucleotides.

[0142] An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formalin with 1 mg of heparin at 42° C., with the hybridization being carried out overnight. An example of stringent wash conditions comprises a 0.2×SSC wash at 65° C. for 15 minutes (see, Sambrook, *infra* for a description of SSC buffer and other nucleic acid hybridization parameters). Often the high stringency wash is preceded by a low stringency wash to remove background probe signal. An example low stringency wash is 2×SSC at 40° C. for 15 minutes. In general, a signal to noise ratio of 5× (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

[0143] After hybridization, unhybridized nucleic acids can be removed by a series of washes, the stringency of which can be adjusted depending upon the desired results. Low stringency washing conditions (e.g., using higher salt and lower temperature) increase sensitivity, but can produce nonspecific hybridization signals and high background signals. Higher stringency conditions (e.g., using lower salt and higher temperature that is closer to the  $T_m$ ) lower the background signal, typically with primarily the specific signal remaining. See, also, Rapley, R. and Walker, J. M. eds., *Molecular Biomethods Handbook* (Humana Press, Inc. 1998).

[0144] “Stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993), *supra*, and in Hames and Higgins, 1 and 2. Stringent hybridization and wash conditions can easily be determined empirically for any test nucleic acid. For example, in determining highly stringent hybridization and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents such as formalin in the hybridization or wash), until a selected set of criteria is met. For example, the hybridization and wash conditions are gradually increased until a probe binds to a perfectly matched complementary target with a signal to noise ratio that is at least 5× as high as that observed for hybridization of the probe to an unmatched target.

[0145] In general, a signal to noise ratio of at least 2× (or higher, e.g., at least 5×, 10×, 20×, 50×, 100×, or more) than that observed for an unrelated probe in the particular hybrid-

ization assay indicates detection of a specific hybridization. Detection of at least stringent hybridization between two sequences in the context of the present invention indicates relatively strong structural similarity to, e.g., the nucleic acids of the present invention provided in the sequence listings herein.

[0146] “Very stringent” conditions are selected to be equal to the thermal melting point ( $T_m$ ) for a particular probe. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the test sequence hybridizes to a perfectly matched probe. For the purposes of the present invention, generally, “highly stringent” hybridization and wash conditions are selected to be about 5° C. lower than the  $T_m$  for the specific sequence at a defined ionic strength and pH (as noted below, highly stringent conditions can also be referred to in comparative terms). Target sequences that are closely related or identical to the nucleotide sequence of interest (e.g., “probe”) can be identified under stringent or highly stringent conditions. Lower stringency conditions are appropriate for sequences that are less complementary.

[0147] “Ultra high-stringency” hybridization and wash conditions are those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio for binding of a probe to a perfectly matched complementary target nucleic acid is at least 10× as high as that observed for hybridization to any unmatched target nucleic acids. A target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid is said to bind to the probe under ultra-high stringency conditions.

[0148] In determining stringent or highly stringent hybridization (or even more stringent hybridization) and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents, such as formamide, in the hybridization or wash), until a selected set of criteria are met. For example, the hybridization and wash conditions are gradually increased until a probe comprising one or more polynucleotide sequences of the invention, e.g., sequences or unique subsequences selected from those given herein (e.g., SEQ ID NO:1-10, 21-26, 33-38, SEQ ID NO:45) and/or complementary polynucleotide sequences, binds to a perfectly matched complementary target (again, a nucleic acid comprising one or more nucleic acid sequences or subsequences selected from those given herein and/or complementary polynucleotide sequences thereof), with a signal to noise ratio that is at least 2× (and optionally 5×, 10×, or 100× or more) as high as that observed for hybridization of the probe to an unmatched target (e.g., a polynucleotide sequence comprising one or more sequences or subsequences selected from known influenza sequences present in public databases such as GenBank at the time of filing, and/or complementary polynucleotide sequences thereof), as desired.

[0149] Using the polynucleotides of the invention, or subsequences thereof, novel target nucleic acids can be obtained; such target nucleic acids are also a feature of the invention. For example, such target nucleic acids include sequences that hybridize under stringent conditions to a

unique oligonucleotide probe corresponding to any of the polynucleotides of the invention, e.g., SEQ ID NO:1-10, 21-26, 33-38, 45).

[0150] Similarly, even higher levels of stringency can be determined by gradually increasing the hybridization and/or wash conditions of the relevant hybridization assay. For example, those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio for binding of the probe to the perfectly matched complementary target nucleic acid is at least 10×, 20×, 50×, 100×, or 500× or more as high as that observed for hybridization to any unmatched target nucleic acids. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a calorimetric label, a radioactive label, or the like. A target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid is said to bind to the probe under ultra-ultra-high stringency conditions and are also features of the invention.

[0151] Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

[0152] Cloning, Mutagenesis and Expression of Biomolecules of Interest

[0153] General texts which describe molecular biological techniques, which are applicable to the present invention, such as cloning, mutation, cell culture and the like, include Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, Calif. (Berger); Sambrook et al., *Molecular Cloning—A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 2000 ("Sambrook") and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2002) ("Ausubel"). These texts describe mutagenesis, the use of vectors, promoters and many other relevant topics related to, e.g., the generation of HA and/or NA molecules, etc.

[0154] Various types of mutagenesis are optionally used in the present invention, e.g., to produce and/or isolate, e.g., novel or newly isolated HA and/or NA molecules and/or to further modify/mutate the polypeptides (e.g., HA and NA molecules as in SEQ ID NO: 11-20 or 27-32 or 39-44) of the invention. They include but are not limited to site-directed, random point mutagenesis, homologous recombination (DNA shuffling), mutagenesis using uracil containing templates, oligonucleotide-directed mutagenesis, phosphorothioate-modified DNA mutagenesis, mutagenesis using gapped duplex DNA or the like. Additional suitable methods include point mismatch repair, mutagenesis using repair-deficient host strains, restriction-selection and restriction-purification, deletion mutagenesis, mutagenesis by total gene synthesis, double-strand break repair, and the like. Mutagenesis, e.g., involving chimeric constructs, is also included in the present invention. In one embodiment, mutagenesis can be guided by known information of the naturally occurring molecule or altered or mutated naturally

occurring molecule, e.g., sequence, sequence comparisons, physical properties, crystal structure or the like.

[0155] The above texts and examples found herein describe these procedures as well as the following publications (and references cited within): Sieber, et al., *Nature Biotechnology*, 19:456-460 (2001); Ling et al., *Approaches to DNA mutagenesis: an overview, Anal Biochem* 254(2): 157-178 (1997); Dale et al., *Oligonucleotide-directed random mutagenesis using the phosphorothioate method, Methods Mol Biol* 57:369-374 (1996); I. A. Lorimer, I. Pastan, *Nucleic Acids Res* 23, 3067-8 (1995); W. P. C. Stemmer, *Nature* 370, 389-91 (1994); Arnold, *Protein engineering for unusual environments, Current Opinion in Biotechnology* 4:450-455 (1993); Bass et al., *Mutant Trp repressors with new DNA-binding specificities, Science* 242:240-245 (1988); Fritz et al., *Oligonucleotide-directed construction of mutations: a gapped duplex DNA procedure without enzymatic reactions in vitro, Nucl Acids Res* 16: 6987-6999 (1988); Kramer et al., *Improved enzymatic in vitro reactions in the gapped duplex DNA approach to oligonucleotide-directed construction of mutations, Nucl Acids Res* 16: 7207 (1988); Sakamar and Khorana, *Total synthesis and expression of a gene for the  $\alpha$ -subunit of bovine rod outer segment guanine nucleotide-binding protein (transducin), Nucl Acids Res* 14: 6361-6372 (1988); Sayers et al., *Y-T Exonucleases in phosphorothioate-based oligonucleotide-directed mutagenesis, Nucl Acids Res* 16:791-802 (1988); Sayers et al., *Strand specific cleavage of phosphorothioate-containing DNA by reaction with restriction endonucleases in the presence of ethidium bromide, (1988) Nucl Acids Res* 16: 803-814; Carter, *Improved oligonucleotide-directed mutagenesis using M13 vectors, Methods in Enzymol* 154: 382-403 (1987); Kramer & Fritz *Oligonucleotide-directed construction of mutations via gapped duplex DNA, Methods in Enzymol* 154:350-367 (1987); Kunkel, *The efficiency of oligonucleotide directed mutagenesis, in Nucleic Acids & Molecular Biology* (Eckstein, F. and Lilley, D. M. J. eds., Springer Verlag, Berlin)) (1987); Kunkel et al., *Rapid and efficient site-specific mutagenesis without phenotypic selection, Methods in Enzymol* 154, 367-382 (1987); Zoller & Smith, *Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template, Methods in Enzymol* 154:329-350 (1987); Carter, *Site-directed mutagenesis, Biochem J* 237:1-7 (1986); Eghtedarzadeh & Henikoff, *Use of oligonucleotides to generate large deletions, Nucl Acids Res* 14: 5115 (1986); Mandecki, *Oligonucleotide-directed double-strand break repair in plasmids of Escherichia coli: a method for site-specific mutagenesis, Proc Natl Acad Sci USA*, 83:7177-7181 (1986); Nakamaye & Eckstein, *Inhibition of restriction endonuclease Nci I cleavage by phosphorothioate groups and its application to oligonucleotide-directed mutagenesis, Nucl Acids Res* 14: 9679-9698 (1986); Wells et al., *Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin, Phil Trans R Soc Lond A* 317: 415-423 (1986); Botstein & Shortle, *Strategies and applications of in vitro mutagenesis, Science* 229:1193-1201 (1985); Carter et al., *Improved oligonucleotide site-directed mutagenesis using M13 vectors, Nucl Acids Res* 13: 4431-4443 (1985); Grundström et al., *Oligonucleotide-directed mutagenesis by microscale 'shot-gun' gene synthesis, Nucl Acids Res* 13:3305-3316 (1985); Kunkel, *Rapid and efficient site-specific mutagenesis without phenotypic selection, Proc Natl Acad Sci USA* 82:488-492 (1985); Smith, In

*vitro* mutagenesis, *Ann Rev Genet.* 19:423-462 (1985); Taylor et al., *The use of phosphorothioate-modified DNA in restriction enzyme reactions to prepare nicked DNA*, *Nucl Acids Res* 13: 8749-8764 (1985); Taylor et al., *The rapid generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA*, *Nucl Acids Res* 13: 8765-8787 (1985); Wells et al., *Cassette mutagenesis: an efficient method for generation of multiple mutations at defined sites*, *Gene* 34:315-323 (1985); Kramer et al., *The gapped duplex DNA approach to oligonucleotide-directed mutation construction*, *Nucl Acids Res* 12: 9441-9456 (1984); Kramer et al., *Point Mismatch Repair*, *Cell* 38:879-887 (1984); Nambiar et al., *Total synthesis and cloning of a gene coding for the ribonuclease S protein*, *Science* 223: 1299-1301 (1984); Zoller & Smith, *Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors*, *Methods in Enzymol* 100:468-500 (1983); and Zoller & Smith, *Oligonucleotide-directed mutagenesis using M13-derived vectors: an efficient and general procedure for the production of point mutations in any DNA fragment*, *Nucl Acids Res* 10:6487-6500 (1982). Additional details on many of the above methods can be found in *Methods in Enzymol* Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis, gene isolation, expression, and other methods.

[0156] Oligonucleotides, e.g., for use in mutagenesis of the present invention, e.g., mutating libraries of the HA and/or NA molecules of the invention, or altering such, are typically synthesized chemically according to the solid phase phosphoramidite triester method described by Beaucage and Caruthers, *Tetrahedron Letts* 22(20):1859-1862, (1981) e.g., using an automated synthesizer, as described in Needham-Van Devanter et al., *Nucleic Acids Res.* 12:6159-6168 (1984).

[0157] In addition, essentially any nucleic acid can be custom or standard ordered from any of a variety of commercial sources, such as The Midland Certified Reagent Company (mcr@oligos.com), The Great American Gene Company (www.genco.com), ExpressGen Inc. (www.expressgen.com), Operon Technologies Inc. (Alameda, Calif.) and many others. Similarly, peptides and antibodies can be custom ordered from any of a variety of sources, such as PeptideGenic (available at pkim@ccnet.com), HTI Bio-products, Inc. (www.htibio.com), BMA Biomedicals Ltd. (U.K.), Bio.Synthesis, Inc., and many others.

[0158] The present invention also relates to host cells and organisms comprising a HA and/or NA molecule or other polypeptide and/or nucleic acid of the invention, e.g., SEQ ID NOS:1-45. Host cells are genetically engineered (e.g., transformed, transduced or transfected) with the vectors of this invention, which can be, for example, a cloning vector or an expression vector. The vector can be, for example, in the form of a plasmid, a bacterium, a virus, a naked polynucleotide, or a conjugated polynucleotide. The vectors are introduced into cells and/or microorganisms by standard methods including electroporation (see, From et al., *Proc Natl Acad Sci USA* 82, 5824 (1985), infection by viral vectors, high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., *Nature* 327, 70-73 (1987)). Berger, Sambrook, and Ausubel provide a variety of appropriate transformation methods. See, above.

[0159] Several well-known methods of introducing target nucleic acids into bacterial cells are available, any of which can be used in the present invention. These include: fusion of the recipient cells with bacterial protoplasts containing the DNA, electroporation, projectile bombardment, and infection with viral vectors, etc. Bacterial cells can be used to amplify the number of plasmids containing DNA constructs of this invention. The bacteria are grown to log phase and the plasmids within the bacteria can be isolated by a variety of methods known in the art (see, for instance, Sambrook). In addition, a plethora of kits are commercially available for the purification of plasmids from bacteria, (see, e.g., EasyPrep™, FlexiPrep™, both from Pharmacia Biotech; StrataClean™, from Stratagene; and, QIAprep™ from Qiagen). The isolated and purified plasmids are then further manipulated to produce other plasmids, used to transfect cells or incorporated into related vectors to infect organisms. Typical vectors contain transcription and translation terminators, transcription and translation initiation sequences, and promoters useful for regulation of the expression of the particular target nucleic acid. The vectors optionally comprise generic expression cassettes containing at least one independent terminator sequence, sequences permitting replication of the cassette in eukaryotes, or prokaryotes, or both, (e.g., shuttle vectors) and selection markers for both prokaryotic and eukaryotic systems. Vectors are suitable for replication and integration in prokaryotes, eukaryotes, or optionally both. See, Gilman & Smith, *Gene* 8:81 (1979); Roberts, et al., *Nature*, 328:731 (1987); Schneider, B., et al., *Protein Expr Purif* 6435:10 (1995); Ausubel, Sambrook, Berger (all supra). A catalogue of Bacteria and Bacteriophages useful for cloning is provided, e.g., by the ATCC, e.g., *The ATCC Catalogue of Bacteria and Bacteriophage* (1992) Gherna et al. (eds.) published by the ATCC. Additional basic procedures for sequencing, cloning and other aspects of molecular biology and underlying theoretical considerations are also found in Watson et al. (1992) *Recombinant DNA Second Edition* Scientific American Books, NY. See, above. Further vectors useful with the sequences herein are illustrated above in the section concerning production of influenza virus for vaccines and the references cited therein.

#### Polypeptide Production and Recovery

[0160] Following transduction of a suitable host cell line or strain and growth of the host cells to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. In some embodiments, a secreted polypeptide product, e.g., a HA and/or NA polypeptide as in a secreted fusion protein form, etc., is then recovered from the culture medium. In other embodiments, a virus particle containing a HA and/or a NA polypeptide of the invention is produced from the cell. Alternatively, cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Eukaryotic or microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods, which are well known to those skilled in the art. Additionally, cells expressing a HA and/or a NA polypeptide product of the invention can be utilized without separating the polypeptide from the cell. In such situations, the polypeptide of the invention is optionally expressed on the cell surface and is examined thus (e.g., by having HA and/or

NA molecules (or fragments thereof, e.g., comprising fusion proteins or the like) on the cell surface bind antibodies, etc. Such cells are also features of the invention.

**[0161]** Expressed polypeptides can be recovered and purified from recombinant cell cultures by any of a number of methods well known in the art, including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography (e.g., using any of the tagging systems known to those skilled in the art), hydroxylapatite chromatography, and lectin chromatography. Protein refolding steps can be used, as desired, in completing configuration of the mature protein. Also, high performance liquid chromatography (HPLC) can be employed in the final purification steps. In addition to the references noted herein, a variety of purification methods are well known in the art, including, e.g., those set forth in Sandana (1997) *Bioseparation of Proteins*, Academic Press, Inc.; and Bollag et al. (1996) *Protein Methods*, 2<sup>nd</sup> Edition Wiley-Liss, NY; Walker (1996) *The Protein Protocols Handbook* Humana Press, NJ; Harris and Angal (1990) *Protein Purification Applications: A Practical Approach* IRL Press at Oxford, Oxford, England; Harris and Angal *Protein Purification Methods: A Practical Approach* IRL Press at Oxford, Oxford, England; Scopes (1993) *Protein Purification: Principles and Practice* 3<sup>rd</sup> Edition Springer Verlag, NY; Janson and Ryden (1998) *Protein Purification: Principles, High Resolution Methods and Applications*, Second Edition Wiley-VCH, NY; and Walker (1998) *Protein Protocols on CD-ROM* Humana Press, NJ.

**[0162]** When the expressed polypeptides of the invention are produced in viruses, the viruses are typically recovered from the culture medium, in which infected (transfected) cells have been grown. Typically, crude medium is clarified prior to concentration of influenza viruses. Common methods include ultrafiltration, adsorption on barium sulfate and elution, and centrifugation. For example, crude medium from infected cultures can first be clarified by centrifugation at, e.g., 1000-2000×g for a time sufficient to remove cell debris and other large particulate matter, e.g., between 10 and 30 minutes. Optionally, the clarified medium supernatant is then centrifuged to pellet the influenza viruses, e.g., at 15,000×g, for approximately 3-5 hours. Following resuspension of the virus pellet in an appropriate buffer, such as STE (0.01 M Tris-HCl; 0.15 M NaCl; 0.0001 M EDTA) or phosphate buffered saline (PBS) at pH 7.4, the virus is concentrated by density gradient centrifugation on sucrose (60%-12%) or potassium tartrate (50%-10%). Either continuous or step gradients, e.g., a sucrose gradient between 12% and 60% in four 12% steps, are suitable. The gradients are centrifuged at a speed, and for a time, sufficient for the viruses to concentrate into a visible band for recovery. Alternatively, and for most large-scale commercial applications, virus is elutriated from density gradients using a zonal-centrifuge rotor operating in continuous mode. Additional details sufficient to guide one of skill through the preparation of influenza viruses from tissue culture are provided, e.g., in Furminger. *Vaccine Production*, in Nicholson et al. (eds.) *Textbook of Influenza* pp. 324-332; Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*, in Cohen & Shafferman (eds.) *Novel Strategies in Design and Production of Vaccines* pp. 141-151, and U.S. Pat. No. 5,690,937. If desired, the recovered

viruses can be stored at -80° C. in the presence of sucrose-phosphate-glutamate (SPG) as a stabilizer

**[0163]** Alternatively, cell-free transcription/translation systems can be employed to produce polypeptides comprising an amino acid sequence or subsequence of, e.g., the sequences given herein such as SEQ ID NOS:11-20 or 27-32 or 39-44, or encoded by the polynucleotide sequences of the invention, e.g., SEQ ID NOS:1-10 or 21-26 or 33-38 or 45. A number of suitable in vitro transcription and translation systems are commercially available. A general guide to in vitro transcription and translation protocols is found in Tymms (1995) *In vitro Transcription and Translation Protocols: Methods in Molecular Biology* Volume 37, Garland Publishing, NY.

**[0164]** In addition, the polypeptides, or subsequences thereof, e.g., subsequences comprising antigenic peptides, can be produced manually or by using an automated system, by direct peptide synthesis using solid-phase techniques (see, Stewart et al. (1969) *Solid-Phase Peptide Synthesis*, WH Freeman Co, San Francisco; Merrifield J (1963) *J Am Chem Soc* 85:2149-2154). Exemplary automated systems include the Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City, Calif.). If desired, subsequences can be chemically synthesized separately, and combined using chemical methods to provide full-length polypeptides.

**[0165]** Modified Amino Acids

**[0166]** Expressed polypeptides of the invention can contain one or more modified amino acids. The presence of modified amino acids can be advantageous in, for example, (a) increasing polypeptide serum half-life, (b) reducing/increasing polypeptide antigenicity, (c) increasing polypeptide storage stability, etc. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production (e.g., N-linked glycosylation at N-X-S/T motifs during expression in mammalian cells) or modified by synthetic means (e.g., via PEGylation).

**[0167]** Non-limiting examples of a modified amino acid include a glycosylated amino acid, a sulfated amino acid, a prenylated (e.g., farnesylated, geranylgeranylated) amino acid, an acetylated amino acid, an acylated amino acid, a PEG-ylated amino acid, a biotinylated amino acid, a carboxylated amino acid, a phosphorylated amino acid, and the like, as well as amino acids modified by conjugation to, e.g., lipid moieties or other organic derivatizing agents. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature. Example protocols are found in Walker (1998) *Protein Protocols on CD-ROM* Human Press, Towata, N.J.

**[0168]** Fusion Proteins

**[0169]** The present invention also provides fusion proteins comprising fusions of the sequences of the invention (e.g., encoding HA and/or NA polypeptides as exemplified by SEQ ID NOS:11-20, 27-32, and 39-44) or fragments thereof with, e.g., immunoglobulins (or portions thereof), sequences encoding, e.g., GFP (green fluorescent protein), or other similar markers, etc. Nucleotide sequences encoding such fusion proteins are another aspect of the invention. Fusion proteins of the invention are optionally used for, e.g., similar applications (including, e.g., therapeutic, prophylactic, diagnostic, experimental, etc. applications as described herein) as the non-fusion proteins of the invention. In addition to

fusion with immunoglobulin sequences and marker sequences, the proteins of the invention are also optionally fused with, e.g., sequences which allow sorting of the fusion proteins and/or targeting of the fusion proteins to specific cell types, regions, etc.

#### [0170] Antibodies

[0171] The polypeptides of the invention can be used to produce antibodies specific for the polypeptides given herein and/or polypeptides encoded by the polynucleotides of the invention, e.g., those shown herein, and conservative variants thereof. Antibodies specific for the above mentioned polypeptides are useful, e.g., for diagnostic and therapeutic purposes, e.g., related to the activity, distribution, and expression of target polypeptides.

[0172] Antibodies specific for the polypeptides of the invention can be generated by methods well known in the art. Such antibodies can include, but are not limited to, polyclonal, monoclonal, chimeric, humanized, single chain, Fab fragments and fragments produced by an Fab expression library.

[0173] Polypeptides do not require biological activity for antibody production (e.g., full length functional hemagglutinin or neuraminidase is not required). However, the polypeptide or oligopeptide must be antigenic. Peptides used to induce specific antibodies typically have an amino acid sequence of at least about 4 amino acids, and often at least 5 or 10 amino acids. Short stretches of a polypeptide can be fused with another protein, such as keyhole limpet hemocyanin, and antibody produced against the chimeric molecule.

[0174] Numerous methods for producing polyclonal and monoclonal antibodies are known to those of skill in the art, and can be adapted to produce antibodies specific for the polypeptides of the invention, and/or encoded by the polynucleotide sequences of the invention, etc. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, NY; Paul (ed.) (1998) *Fundamental Immunology Fourth Edition*, Lippincott-Raven, Lippincott Williams & Wilkins; Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, NY; Stites et al. (eds.) *Basic and Clinical Immunology* (4th ed.) Lange Medical Publications, Los Altos, Calif., and references cited therein; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed.) Academic Press, New York, N.Y.; and Kohler and Milstein (1975) *Nature* 256: 495-497. Other suitable techniques for antibody preparation include selection of libraries of recombinant antibodies in phage or similar vectors. See, Huse et al. (1989) *Science* 246: 1275-1281; and Ward, et al. (1989) *Nature* 341: 544-546. Specific monoclonal and polyclonal antibodies and antisera will usually bind with a  $K_D$  of, e.g., at least about 0.1  $\mu$ M, at least about 0.01  $\mu$ M or better, and, typically and at least about 0.001  $\mu$ M or better.

[0175] For certain therapeutic applications, humanized antibodies are desirable. Detailed methods for preparation of chimeric (humanized) antibodies can be found in U.S. Pat. No. 5,482,856. Additional details on humanization and other antibody production and engineering techniques can be found in Borrebaeck (ed.) (1995) *Antibody Engineering*, 2<sup>nd</sup> Edition Freeman and Company, NY (Borrebaeck); McCafferty et al. (1996) *Antibody Engineering, A Practical Approach* IRL at Oxford Press, Oxford, England (McCafferty), and Paul (1995) *Antibody Engineering Protocols*

Humana Press, Towata, N.J. (Paul). Additional details regarding specific procedures can be found, e.g., in Ostberg et al. (1983), *Hybridoma* 2: 361-367, Ostberg, U.S. Pat. No. 4,634,664, and Engelman et al., U.S. Pat. No. 4,634,666.

#### [0176] Defining Polypeptides by Immunoreactivity

[0177] Because the polypeptides of the invention provide a variety of new polypeptide sequences (e.g., comprising HA and NA molecules), the polypeptides also provide new structural features which can be recognized, e.g., in immunological assays. The generation of antisera which specifically bind the polypeptides of the invention, as well as the polypeptides which are bound by such antisera, are features of the invention.

[0178] For example, the invention includes polypeptides (e.g., HA and NA molecules) that specifically bind to or that are specifically immunoreactive with an antibody or antisera generated against an immunogen comprising an amino acid sequence selected from one or more of the sequences given herein (e.g., SEQ ID NOS:11-20, 27-32, and 39-44), etc. To eliminate cross-reactivity with other homologues, the antibody or antisera is subtracted with the HA and/or NA molecules found in public databases at the time of filing, e.g., the "control" polypeptide(s). Where the other control sequences correspond to a nucleic acid, a polypeptide encoded by the nucleic acid is generated and used for antibody/antisera subtraction purposes.

[0179] In one typical format, the immunoassay uses a polyclonal antiserum which was raised against one or more polypeptide comprising one or more of the sequences corresponding to the sequences herein (e.g., SEQ ID NOS:11-20, 27-32, and 39-44), etc. or a substantial subsequence thereof (i.e., at least about 30% of the full length sequence provided). The set of potential polypeptide immunogens derived from the present sequences are collectively referred to below as "the immunogenic polypeptides." The resulting antisera is optionally selected to have low cross-reactivity against the control hemagglutinin and/or neuraminidase homologues and any such cross-reactivity is removed, e.g., by immunoabsorption, with one or more of the control hemagglutinin and neuraminidase homologues, prior to use of the polyclonal antiserum in the immunoassay.

[0180] In order to produce antisera for use in an immunoassay, one or more of the immunogenic polypeptides is produced and purified as described herein. For example, recombinant protein can be produced in a recombinant cell. An inbred strain of mice (used in this assay because results are more reproducible due to the virtual genetic identity of the mice) is immunized with the immunogenic protein(s) in combination with a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see, e.g., Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a standard description of antibody generation, immunoassay formats and conditions that can be used to determine specific immunoreactivity). Additional references and discussion of antibodies is also found herein and can be applied here to defining polypeptides by immunoreactivity. Alternatively, one or more synthetic or recombinant polypeptide derived from the sequences disclosed herein is conjugated to a carrier protein and used as an immunogen.

[0181] Polyclonal sera are collected and titered against the immunogenic polypeptide in an immunoassay, for example,

a solid phase immunoassay with one or more of the immunogenic proteins immobilized on a solid support. Polyclonal antisera with a titer of  $10^6$  or greater are selected, pooled and subtracted with the control hemagglutinin and/or neuraminidase polypeptide(s) to produce subtracted pooled titrated polyclonal antisera.

[0182] The subtracted pooled titrated polyclonal antisera are tested for cross reactivity against the control homologue(s) in a comparative immunoassay. In this comparative assay, discriminatory binding conditions are determined for the subtracted titrated polyclonal antisera which result in at least about a 5-10 fold higher signal to noise ratio for binding of the titrated polyclonal antisera to the immunogenic polypeptides as compared to binding to the control homologues. That is, the stringency of the binding reaction is adjusted by the addition of non-specific competitors such as albumin or non-fat dry milk, and/or by adjusting salt conditions, temperature, and/or the like. These binding conditions are used in subsequent assays for determining whether a test polypeptide (a polypeptide being compared to the immunogenic polypeptides and/or the control polypeptides) is specifically bound by the pooled subtracted polyclonal antisera. In particular, test polypeptides which show at least a 2-5 $\times$  higher signal to noise ratio than the control receptor homologues under discriminatory binding conditions, and at least about a  $\frac{1}{2}$  signal to noise ratio as compared to the immunogenic polypeptide(s), shares substantial structural similarity with the immunogenic polypeptide as compared to the known receptor, etc., and is, therefore a polypeptide of the invention.

[0183] In another example, immunoassays in the competitive binding format are used for detection of a test polypeptide. For example, as noted, cross-reacting antibodies are removed from the pooled antisera mixture by immunoabsorption with the control polypeptides. The immunogenic polypeptide(s) are then immobilized to a solid support which is exposed to the subtracted pooled antisera. Test proteins are added to the assay to compete for binding to the pooled subtracted antisera. The ability of the test protein(s) to compete for binding to the pooled subtracted antisera as compared to the immobilized protein(s) is compared to the ability of the immunogenic polypeptide(s) added to the assay to compete for binding (the immunogenic polypeptides compete effectively with the immobilized immunogenic polypeptides for binding to the pooled antisera). The percent cross-reactivity for the test proteins is calculated, using standard calculations.

[0184] In a parallel assay, the ability of the control protein(s) to compete for binding to the pooled subtracted antisera is optionally determined as compared to the ability of the immunogenic polypeptide(s) to compete for binding to the antisera. Again, the percent cross-reactivity for the control polypeptide(s) is calculated, using standard calculations. Where the percent cross-reactivity is at least 5-10 $\times$  as high for the test polypeptides as compared to the control polypeptide(s) and/or where the binding of the test polypeptides is approximately in the range of the binding of the immunogenic polypeptides, the test polypeptides are said to specifically bind the pooled subtracted antisera.

[0185] In general, the immunoabsorbed and pooled antisera can be used in a competitive binding immunoassay as described herein to compare any test polypeptide to the

immunogenic and/or control polypeptide(s). In order to make this comparison, the immunogenic, test and control polypeptides are each assayed at a wide range of concentrations and the amount of each polypeptide required to inhibit 50% of the binding of the subtracted antisera to, e.g., an immobilized control, test or immunogenic protein is determined using standard techniques. If the amount of the test polypeptide required for binding in the competitive assay is less than twice the amount of the immunogenic polypeptide that is required, then the test polypeptide is said to specifically bind to an antibody generated to the immunogenic protein, provided the amount is at least about 5-10 $\times$  as high as for the control polypeptide.

[0186] As an additional determination of specificity, the pooled antisera is optionally fully immunosorbed with the immunogenic polypeptide(s) (rather than the control polypeptide(s)) until little or no binding of the resulting immunogenic polypeptide subtracted pooled antisera to the immunogenic polypeptide(s) used in the immunosorption is detectable. This fully immunosorbed antisera is then tested for reactivity with the test polypeptide. If little or no reactivity is observed (i.e., no more than 2 $\times$  the signal to noise ratio observed for binding of the fully immunosorbed antisera to the immunogenic polypeptide), then the test polypeptide is specifically bound by the antisera elicited by the immunogenic protein.

#### Nucleic Acid and Polypeptide Sequence Variants

[0187] As described herein, the invention provides for nucleic acid polynucleotide sequences and polypeptide amino acid sequences, e.g., hemagglutinin and neuraminidase sequences, and, e.g., compositions and methods comprising said sequences. Examples of said sequences are disclosed herein (e.g., SEQ ID NOS:1-45). However, one of skill in the art will appreciate that the invention is not necessarily limited to those sequences disclosed herein and that the present invention also provides many related and unrelated sequences with the functions described herein, e.g., encoding a HA and/or a NA molecule.

[0188] One of skill will also appreciate that many variants of the disclosed sequences are included in the invention. For example, conservative variations of the disclosed sequences that yield a functionally identical sequence are included in the invention. Variants of the nucleic acid polynucleotide sequences, wherein the variants hybridize to at least one disclosed sequence, are considered to be included in the invention. Unique subsequences of the sequences disclosed herein, as determined by, e.g., standard sequence comparison techniques, are also included in the invention.

#### [0189] Silent Variations

[0190] Due to the degeneracy of the genetic code, any of a variety of nucleic acid sequences encoding polypeptides and/or viruses of the invention are optionally produced, some which can bear lower levels of sequence identity to the HA and NA nucleic acid and polypeptide sequences herein. The following provides a typical codon table specifying the genetic code, found in many biology and biochemistry texts.

TABLE 1

Codon Table			
Amino acids			Codon
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU
Serine	Ser	S	AGC AGU UCA UCC UCG UCU
Threonine	Thr	T	ACA ACC ACG ACU
Valine	Val	V	GUA GUC GUG GUU
Tryptophan	Trp	W	UGG
Tyrosine	Tyr	Y	UAC UAU

[0191] The codon table shows that many amino acids are encoded by more than one codon. For example, the codons AGA, AGG, CGA, CGC, CGG, and CGU all encode the amino acid arginine. Thus, at every position in the nucleic acids of the invention where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described above without altering the encoded polypeptide. It is understood that U in an RNA sequence corresponds to T in a DNA sequence.

[0192] Such “silent variations” are one species of “conservatively modified variations,” discussed below. One of skill will recognize that each codon in a nucleic acid (except ATG, which is ordinarily the only codon for methionine, and TTG, which is ordinarily the only codon for tryptophan) can be modified by standard techniques to encode a functionally identical polypeptide. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in any described sequence. The invention, therefore, explicitly provides each and every possible variation of a nucleic acid sequence encoding a polypeptide of the invention that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code (e.g., as set forth in Table 1, or as is commonly available in the art) as applied to the nucleic acid sequence encoding a hemagglutinin or a

neuraminidase polypeptide of the invention. All such variations of every nucleic acid herein are specifically provided and described by consideration of the sequence in combination with the genetic code. One of skill is fully able to make these silent substitutions using the methods herein.

#### [0193] Conservative Variations

[0194] Owing to the degeneracy of the genetic code, “silent substitutions” (i.e., substitutions in a nucleic acid sequence which do not result in an alteration in an encoded polypeptide) are an implied feature of every nucleic acid sequence of the invention which encodes an amino acid. Similarly, “conservative amino acid substitutions,” in one or a few amino acids in an amino acid sequence are substituted with different amino acids with highly similar properties, are also readily identified as being highly similar to a disclosed construct such as those herein. Such conservative variations of each disclosed sequence are a feature of the present invention.

[0195] “Conservative variations” of a particular nucleic acid sequence refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or, where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences, see, Table 2 below. One of skill will recognize that individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 4%, 3%, 2% or 1%) in an encoded sequence are “conservatively modified variations” where the alterations result in the deletion of an amino acid, addition of an amino acid, or substitution of an amino acid with a chemically similar amino acid. Thus, “conservative variations” of a listed polypeptide sequence of the present invention include substitutions of a small percentage, typically less than 5%, more typically less than 4%, 3%, 2% or 1%, of the amino acids of the polypeptide sequence, with a conservatively selected amino acid of the same conservative substitution group. Finally, the addition of sequences which do not alter the encoded activity of a nucleic acid molecule, such as the addition of a non-functional sequence, is a conservative variation of the basic nucleic acid.

TABLE 2

Conservative Substitution Groups			
1 Alanine (A)	Serine (S)	Threonine (T)	
2 Aspartic acid (D)	Glutamic acid (E)		
3 Asparagine (N)	Glutamine (Q)		
4 Arginine (R)	Lysine (K)		
5 Isoleucine (I)	Leucine (L)	Methionine (M)	Valine (V)
6 Phenylalanine (F)	Tyrosine (Y)	Tryptophan (W)	

#### [0196] Unique Polypeptide and Polynucleotide Subsequences

[0197] In one aspect, the invention provides a nucleic acid which comprises a unique subsequence in a nucleic acid selected from the sequence of HA and NA molecules disclosed herein, e.g., SEQ ID NOS:1-10, 21-26, 33-38, and 45. The unique subsequence is unique as compared to a nucleic acids corresponding to nucleic acids such as, e.g., those found in GenBank or other similar public databases at the time of filing. Alignment can be performed using, e.g.,

BLAST set to default parameters. Any unique subsequence is useful, e.g., as a probe to identify the nucleic acids of the invention. See, above.

[0198] Similarly, the invention includes a polypeptide which comprises a unique subsequence in a polypeptide selected from the sequence of HA and NA molecules disclosed herein, e.g., SEQ ID NOS:11-20, 27-32, and 39-44. Here, the unique subsequence is unique as compared to a polypeptide corresponding to, e.g., the amino acid corresponding to polynucleotide sequences found in, e.g., GenBank or other similar public databases at the time of filing.

[0199] The invention also provides for target nucleic acids which hybridize under stringent conditions to a unique coding oligonucleotide which encodes a unique subsequence in a polypeptide selected from the sequences of HA and NA molecules of the invention wherein the unique subsequence is unique as compared to a polypeptide corresponding to any of the control polypeptides (sequences of, e.g., the nucleic acids corresponding to those found in, e.g., GenBank or other similar public databases at the time of filing). Unique sequences are determined as noted above.

[0200] Sequence Comparison, Identity, and Homology

[0201] The terms "identical" or percent "identity," in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (or other algorithms available to persons of skill) or by visual inspection.

[0202] The phrase "substantially identical," in the context of two nucleic acids or polypeptides (e.g., DNAs encoding a HA or NA molecule, or the amino acid sequence of a HA or NA molecule) refers to two or more sequences or subsequences that have at least about 90%, preferably 91%, most preferably 92%, 93%, 94%, 95%, 96%, 97%, 98%, 98.5%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection. Such "substantially identical" sequences are typically considered to be "homologous," without reference to actual ancestry. Preferably, "substantial identity" exists over a region of the amino acid sequences that is at least about 200 residues in length, at least about 250 residues, at least about 300 residues, 350 residues, 400 residues, 425 residues, 450 residues, 475 residues, 480 residues, 490 residues, 495 residues, 499 residues, 500 residues, 502 residues, 559 residues, 565 residues, or 566 residues, or over the full length of the two sequences to be compared.

[0203] For sequence comparison and homology determination, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0204] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv Appl Math* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J Mol Biol* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc Natl Acad Sci USA* 85:2444 (1988), by computerized implementations of algorithms such as GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis., or by visual inspection (see generally, Ausubel et al., *supra*).

[0205] One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J Mol Biol* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information ([www.ncbi.nlm.nih.gov/](http://www.ncbi.nlm.nih.gov/)). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (see, Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see, Henikoff & Henikoff (1989) *Proc Natl Acad Sci USA* 89:10915).

[0206] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc Natl Acad Sci USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0207] Another example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progres-



sive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle (1987) *J. Mol. Evol.* 35:351-360. The method used is similar to the method described by Higgins & Sharp (1989) *CABIOS* 5:151-153. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster can then be aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences can be aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program can also be used to plot a dendrogram or tree representation of clustering relationships. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison.

[0208] An additional example of an algorithm that is suitable for multiple DNA, or amino acid, sequence alignments is the CLUSTALW program (Thompson, J. D. et al. (1994) *Nucl. Acids. Res.* 22: 4673-4680). CLUSTALW performs multiple pairwise comparisons between groups of sequences and assembles them into a multiple alignment based on homology. Gap open and Gap extension penalties can be, e.g., 10 and 0.05 respectively. For amino acid alignments, the BLOSUM algorithm can be used as a protein weight matrix. See, e.g., Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919.

#### Digital Systems

[0209] The present invention provides digital systems, e.g., computers, computer readable media and integrated systems comprising character strings corresponding to the sequence information herein for the nucleic acids and isolated or recombinant polypeptides herein, including, e.g., the sequences shown herein, and the various silent substitutions and conservative substitutions thereof. Integrated systems can further include, e.g., gene synthesis equipment for making genes corresponding to the character strings.

[0210] Various methods known in the art can be used to detect homology or similarity between different character strings (see, above), or can be used to perform other desirable functions such as to control output files, provide the basis for making presentations of information including the sequences and the like. Examples include BLAST, discussed supra. Computer systems of the invention can include such programs, e.g., in conjunction with one or more data file or data base comprising a sequence as noted herein.

[0211] Thus, different types of homology and similarity of various stringency and length between various HA or NA sequences or fragments, etc. can be detected and recognized in the integrated systems herein. For example, many homology determination methods have been designed for comparative analysis of sequences of biopolymers, for spell-checking in word processing, and for data retrieval from various databases. With an understanding of double-helix pair-wise complement interactions among 4 principal nucleobases in natural polynucleotides, models that simulate annealing of complementary homologous polynucleotide strings can also be used as a foundation of sequence alignment or other operations typically performed on the char-

acter strings corresponding to the sequences herein (e.g., word-processing manipulations, construction of figures comprising sequence or subsequence character strings, output tables, etc.).

[0212] Thus, standard desktop applications such as word processing software (e.g., Microsoft Word™ or Corel Word-Perfect™) and database software (e.g., spreadsheet software such as Microsoft Excel™, Corel Quattro Pro™, or database programs such as Microsoft Access™, Paradox™, Gene-Works™, or MacVector™ or other similar programs) can be adapted to the present invention by inputting a character string corresponding to one or more polynucleotides and polypeptides of the invention (either nucleic acids or proteins, or both). For example, a system of the invention can include the foregoing software having the appropriate character string information, e.g., used in conjunction with a user interface (e.g., a GUI in a standard operating system such as a Windows, Macintosh or LINUX system) to manipulate strings of characters corresponding to the sequences herein. As noted, specialized alignment programs such as BLAST can also be incorporated into the systems of the invention for alignment of nucleic acids or proteins (or corresponding character strings).

[0213] Systems in the present invention typically include a digital computer with data sets entered into the software system comprising any of the sequences herein. The computer can be, e.g., a PC (Intel x86 or Pentium chip-compatible DOS™, OS2™ WINDOWS™ WINDOWSNT™, WINDOWS95™, WINDOWS2000™, WINDOWS98™, LINUX based machine, a MACINTOSH™, Power PC, or a UNIX based (e.g., SUN™ work station) machine) or other commercially available computer that is known to one of skill. Software for aligning or otherwise manipulating sequences is available, or can easily be constructed by one of skill using a standard programming language such as Visualbasic, PERL, Fortran, Basic, Java, or the like.

[0214] Any controller or computer optionally includes a monitor which is often a cathode ray tube ("CRT") display, a flat panel display (e.g., active matrix liquid crystal display, liquid crystal display), or others. Computer circuitry is often placed in a box which includes numerous integrated circuit chips, such as a microprocessor, memory, interface circuits, and others. The box also optionally includes a hard disk drive, a floppy disk drive, a high capacity removable drive such as a writeable CD-ROM, and other common peripheral elements. Inputting devices such as a keyboard or mouse optionally provide for input from a user and for user selection of sequences to be compared or otherwise manipulated in the relevant computer system.

[0215] The computer typically includes appropriate software for receiving user instructions, either in the form of user input into a set parameter fields, e.g., in a GUI, or in the form of preprogrammed instructions, e.g., preprogrammed for a variety of different specific operations. The software then converts these instructions to appropriate language for instructing the operation, e.g., of appropriate mechanisms or transport controllers to carry out the desired operation. The software can also include output elements for controlling nucleic acid synthesis (e.g., based upon a sequence or an alignment of sequences herein), comparisons of samples for differential gene expression, or other operations.

### Kits and Reagents

[0216] The present invention is optionally provided to a user as a kit. For example, a kit of the invention contains one or more nucleic acid, polypeptide, antibody, or cell line described herein (e.g., comprising, or with, a HA and/or NA molecule of the invention). The kit can contain a diagnostic nucleic acid or polypeptide, e.g., antibody, probe set, e.g., as a cDNA micro-array packaged in a suitable container, or other nucleic acid such as one or more expression vector. The kit can also further comprise, one or more additional reagents, e.g., substrates, labels, primers, for labeling expression products, tubes and/or other accessories, reagents for collecting samples, buffers, hybridization chambers, cover slips, etc. The kit optionally further comprises an instruction set or user manual detailing preferred methods of using the kit components for discovery or application of diagnostic sets, etc.

[0217] When used according to the instructions, the kit can be used, e.g., for evaluating a disease state or condition, for evaluating effects of a pharmaceutical agent or other treatment intervention on progression of a disease state or condition in a cell or organism, or for use as a vaccine, etc.

[0218] In an additional aspect, the present invention provides system kits embodying the methods, composition, systems and apparatus herein. System kits of the invention optionally comprise one or more of the following: (1) an apparatus, system, system component or apparatus component; (2) instructions for practicing methods described herein, and/or for operating the apparatus or apparatus components herein and/or for using the compositions herein. In a further aspect, the present invention provides for the use of any apparatus, apparatus component, composition or kit herein, for the practice of any method or assay herein, and/or for the use of any apparatus or kit to practice any assay or method herein.

[0219] Additionally, the kits can include one or more translation system as noted above (e.g., a cell) with appropriate packaging material, containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like. Similarly, products of the translation systems (e.g., proteins such as HA and/or NA molecules) can be provided in kit form, e.g., with containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like.

[0220] To facilitate use of the methods and compositions of the invention, any of the vaccine components and/or compositions, e.g., reassorted virus in allantoic fluid, etc., and additional components, such as, buffer, cells, culture medium, useful for packaging and infection of influenza viruses for experimental or therapeutic vaccine purposes, can be packaged in the form of a kit. Typically, the kit contains, in addition to the above components, additional materials which can include, e.g., instructions for performing the methods of the invention, packaging material, and a container.

### EXAMPLES

#### Example 1

#### Construction and Analysis of H5N1 ca Viruses and Vaccines

[0221] Various sequences herein comprising H5N1 HA/NA sequences were used to create influenza viruses and

vaccines. The HA sequences in such vaccines were altered from wild-type by removal of the polybasic cleavage site within the HA. The HA/NA sequences were reassorted (in a 6:2 reassortment) with ca A/AA/6/60 (a ts, att, ca virus, see above).

[0222] Three strains of H5N1 influenza were used in this example: A/VN/1203/2004, A/HK/491/1997, and A/HK/213/2003. Such strains are also referred to within this example as the '97, '03, and '04 strains based on their year designations. The HA sequence homology of these three strains is 95-96%. FIG. 1 illustrates modification of the polybasic cleavage site of an exemplary HA sequence, the '04 HA sequences, used to construct the viruses/vaccines. As stated previously, various embodiments of the invention comprise sequences which have differing regions of the polybasic cleavage site removed. See above.

[0223] As stated, the modified H5N1 sequences (i.e., the modified '97, '03, and '04 genes) were used to construct 6:2 reassortant viruses with ca A/AA/6/60. It will be appreciated, and is pointed out elsewhere herein, that other desirable backbones could also have been used (e.g., PR8, etc.).

[0224] In the 6:2 reassortants of this example, the HA and NA gene sequences were derived from one or more wild type parent virus, i.e., the HA and NA gene sequences of the '03 virus were derived from A/HK/213/2003, the HA and NA gene sequences of the '04 virus were derived from A/VN/1203/2004, and the HA gene sequence of the '97 virus was derived from A/HK/491/1997 while the NA gene sequence was derived from A/HK/486/1997. The remaining genes of the 6:2 reassortants were characterized by sequence analysis as derived from the A/AA/6/60 ca parent virus. The reassorted viruses replicated to 8.0-8.5 log<sub>10</sub> TCID<sub>50</sub> in eggs. However, it will be appreciated that other embodiments wherein the log<sub>10</sub>TCID<sub>50</sub> comprises from about 7.0 to about 9.0, from about 7.5-8.5, or from about 8.0-8.5 are also within the scope of the invention. The cleavability of the modified HA in the constructed viruses by endogenous proteases was restricted in vitro and the viruses were dependent on trypsin (e.g., from about 0.1 ug/ml to about 1.0 ug/ml) for growth. The constructed viruses were temperature sensitive as assayed by an in vitro assay.

[0225] The H5N1 ca reassortant viruses (having the modified '97, '03, or '04 HA genes) were not lethal for chickens. For example, when 4-week-old SPF white Plymouth Rock chickens were inoculated intravenously with a 1:10 dilution of stock virus (10<sup>8-8.75</sup> TCID<sub>50</sub>/ml) and observed for 10 days, it was observed that 8 out of 8 chickens died within 1-2 days when wild-type '97, '03, and '04 H5N1 were used, while 0 of 8 chickens died when the H5N1 ca reassortant viruses were used. As can be seen in FIG. 2, the intranasally administered H5N1 ca reassortant viruses did not replicate in chickens.

[0226] The H5N1/AA ca reassortants were also not lethal for mice. See FIG. 3, which also shows the TCID<sub>50</sub> for the H5N1 wild-type strains. FIG. 4 shows that the 1997 and 2004 H5N1 ca reassortant viruses were restricted in replication in mice. FIG. 5, shows that the H5N1 ca reassorted viruses are restricted in replication in lungs of mice.

[0227] A comparison of the serum HAI antibody titers elicited in mice following a single intranasal dose of vaccine (2003 ca as compared against 2003 wild-type), is shown in

FIG. 6. FIGS. 7 and 15 show similar measurements, but using serum neutralizing antibody titers.

[0228] FIG. 8 displays that the H5N1 ca reassortant viruses protect mice from lethal challenge with 50, 500, or 5,000 LD<sub>50</sub> of wild-type H5N1 virus. FIG. 9 shows the efficacy of protection from pulmonary replication of homologous and heterologous H5N1 challenge viruses in mice. As can be seen, the ca reassortants replicated less well than the wild-type viruses did. FIG. 10 shows related data using upper respiratory tracts of mice. Those of skill in the art will be familiar with homologous and heterologous challenges (e.g., testing whether 2003 vaccine protects against a 2003 wild-type challenge (homologous) or whether a 2003 vaccine protects against a 1997 wild-type challenge (heterologous), etc.).

[0229] FIG. 11 shows efficacy of protection conferred by 2004 H5N1 ca vaccine against high dose (10<sup>5</sup>TCID<sub>50</sub>) challenge with homologous or heterologous H5N1 wild-type viruses in mice. FIG. 12 shows efficacy of protection conferred by 1997 and 2003 H5N1 ca vaccines against high dose (10<sup>5</sup>TCID<sub>50</sub>) challenge with homologous or heterologous H5N1 wild-type viruses in mice. FIG. 13 shows efficacy of protection conferred by 2004 H5N1 ca vaccine against low or high doses of homologous H5N1 wild-type virus challenge in mice. FIGS. 11-13 demonstrate that the tested vaccines could protect against other related viruses.

[0230] In healthy human adults nasal spray administration the '04 vaccine was well tolerated and its replication was highly restricted. See FIG. 27 for replication restriction of the vaccine in healthy adults. HI antibody responses to 10<sup>6.7</sup> TCID<sub>50</sub> of the '04 vaccine were also observed in some of the healthy adults. See FIG. 28.

[0231] The current example demonstrates several points concerning exemplary H5N1 ca reassortant viruses/vaccines of the invention. The modified ca reassortant '97, '03, and '04 viruses were shown to have in vitro ts phenotype, loss of pathogenicity in chickens and attenuation in mice. It is expected that attenuation is also present in ferrets. Efficacy of protection and cross-protection against lethal challenge and systemic spread with wild-type viruses in mice was also shown. Efficacy of protection and cross-protections against replication of wild-type challenge viruses in the respiratory tract of mice is also expected.

[0232] It is contemplated to use these (and similar) viruses/vaccines to determine whether immunogenicity and efficacy is improved following 2 doses of vaccine; to assess immunogenicity in non-human primates; to assess attenuation and vaccine efficacy in ferrets; to determine the contribution of humoral and cellular immunity to observed efficacy of the produced vaccines in mice; to determine which residues of the 2003 HA contribute to enhanced immunogenicity and introduce them into 1997 and 2004 HAs; and to determine the effects of deleting the multibasic amino acid cleavage site and of the gene constellation.

#### Example 2

##### Construction and Analysis of H6 ca Viruses and Vaccines

[0233] A set of three recombinant influenza viruses and vaccines comprising H6 HA sequences were prepared: (a)

A/Duck, which comprised the H6 HA and N9 NA of A/Duck77; (b) A/Teal, which comprised the H6 HA and N1 NA of A/Teal97; and (c) A/Mallard, which comprised the H6 HA and N2 NA of A/Mallard85. The six internal genome segments of each recombinant virus were those of ca A/AA/6/60.

[0234] Each of the A/Duck, A/Teal, and A/Mallard recombinant viruses was attenuated in nasal turbinates and lungs of ferrets. Ferrets were intranasally inoculated with 10<sup>7</sup> TCID<sub>50</sub> recombinant (ca; see paragraph immediately above) or wild-type (wt) H6 influenza virus. Nasal turbinate and lung tissue was harvested from the ferrets three days post-infection for examination. FIG. 16 shows that the nasal turbinate and lung tissue of ferrets inoculated with recombinant virus (ca) exhibited lower virus titers than did the nasal turbinate and lung tissue of ferrets inoculated with the respective counterpart wt virus.

[0235] Each of the A/Duck, A/Teal, and A/Mallard recombinant (ca) viruses was also immunogenic in the ferrets. See FIG. 17.

[0236] FIG. 18 shows the efficacy of protection conferred by the A/Duck, A/Teal, and A/Mallard vaccines. Ferrets were vaccinated with a single dose of 7 log<sub>10</sub> PFU recombinant A/Duck, A/Teal, or A/Mallard vaccine. The ferrets were then challenged with 7 log<sub>10</sub> PFU wt A/Duck, A/Teal, or A/Mallard virus. Three days post challenge lungs and nasal turbinates of the ferrets were harvested and virus titer in the tissues was determined.

[0237] FIG. 18 shows efficacy of protection conferred by the recombinant (ca) H6 vaccines against homologous and heterologous wild-type H6 viruses in ferrets.

#### Example 3

##### Construction and Analysis of an H7N3 BC 04 ca, Virus and Vaccine

[0238] A further recombinant influenza virus and vaccine was prepared using the HA H7 and NA N3 sequences of A/cK/BC/CN-6/04 (BC 04 ca). These HA and NA sequences were combined with the six internal genome segments of ca A/AA/6/60.

[0239] The BC 04 ca vaccine was attenuated in the ferrets. Ferrets were intranasally inoculated with 10<sup>7</sup> TCID<sub>50</sub> vaccine in 0.5 mL. Three days following inoculation, ferret nasal turbinates, lungs, brain, and olfactory bulb were harvested. Virus titer in each of these tissues was diminished in ferrets inoculated with the vaccine virus relative to ferrets inoculated with wt viruses A/BC/CN-6/04 or A/BC/CN-7/04. See FIG. 19.

[0240] The BC 04 ca vaccine was immunogenic in mice. In mice receiving the BC 04 ca vaccine, neutralizing antibodies were detected at 4 weeks and these titers rose over 8 weeks. A second dose of vaccine boosted antibody titer but final titer achieved was similar to that following a single dose. See FIG. 20.

[0241] FIGS. 21 and 22 show the efficacy of protection conferred by the BC 04 ca vaccine against both homologous and heterologous H7 wt viruses. For FIG. 21, mice were intranasally inoculated with 1 dose vaccine four weeks before challenge, 1 dose vaccine 8 weeks before challenge,

or 2 doses vaccine (administered 4 weeks apart) before lethal challenge with 50 LD<sub>50</sub> homologous (A/ck/BC/CN-7/04) and heterologous (A/NL/219/03 or A/tk/Eng/63) H7 wt viruses. Weight change of the mice following lethal challenge was monitored each day for fourteen days, to monitor morbidity associated with the wt influenza virus challenge.

[0242] For each of the mice lethally challenged with the homologous A/ck/BC/CN-7/04 virus little or no weight change was observed regardless of whether 1 dose of vaccine was administered 4 weeks prior to challenge (a), 1 dose of vaccine was administered 8 weeks prior to challenge (b) or 2 doses of vaccine were administered prior to challenge (c). Likewise, little to no weight loss occurred following challenge of the mice with either heterologous influenza virus, A/NL/2109/03 (d, e, f) or A/tk/Eng/63 (g, h, i). Again, the lack of weight loss was observed regardless of whether 1 dose of vaccine was administered 4 weeks prior to challenge (d or g), 1 dose of vaccine was administered 8 weeks prior to challenge (e, or h), or 2 doses of vaccine were administered prior to challenge (f or i).

[0243] FIG. 22 provides further evidence of the efficacy of the H7N3 BC 04 ca vaccine. In both nasal turbinates (a) and lungs (b) of mice receiving the H7N3 BC 04 ca vaccine, protection was observed against challenge using ck/BC/CN-6/04 (H7N3), ck/BC/CN-7/04 (H7N3), NL/219/03 (H7N7), tk/Eng/63 (H7N3), tk/UT/95 (H7N3), and tk/VA/02 (H7N2) viruses.

#### Example 4

##### Construction and Analysis of an H9N2 G9/AA ca, Virus and Vaccine

[0244] A further recombinant influenza virus and vaccine was prepared using the HA H9 and NA N2 sequences of A/ck/Hong Kong/G9/97 (G9/AA ca). These HA and NA sequences were combined with the six internal genome segments of ca A/AA/6/60.

[0245] The H9N2 G9/AA ca vaccine was attenuated in the ferrets. See FIG. 23, which shows reduced virus titers in nasal turbinates (a) and lungs (b) of ferrets following administration of the H9N2 G9/AA ca virus relative to the H9N2 G9 wt virus.

[0246] FIG. 24 provides evidence of the efficacy of the H9N2 G9 ca vaccine in mice. In the mice receiving the H9N2 G9 ca vaccine, protection was observed against challenge using H9N2 G9 wt and A/HK/1073/99 viruses.

[0247] The H9N2 G9/AA ca vaccine was also well tolerated in healthy adults in a clinical trial setting. Healthy adults were administered the H9N2 G9/AA ca vaccine by nose drop. In the healthy adults, the H9N2 G9/AA ca vaccine was highly restricted in replication. See FIG. 25. Furthermore, administration of 10<sup>7.0</sup> TCID<sub>50</sub> of H9N2 G9/AA ca vaccine induced  $\geq 4$ -fold increases in HI titer in 92% of healthy volunteers. See FIG. 26.

[0248] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques and apparatus described above may be used in various combinations. All publications, patents, patent applications, or other documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, or other document were individually indicated to be incorporated by reference for all purposes. In particular, U.S. provisional application Nos. 60/821,832 filed Aug. 9, 2006 and 60/942,804, filed Jun. 8, 2007, are incorporated herein in their entirety for all purposes.

#### SPECIFIC EMBODIMENTS

[0249] Additional embodiments of the present invention are presented in Table 3 and 4.

TABLE 3

##### Specific embodiments

- 1 An isolated polypeptide, wherein said polypeptide is selected from the group consisting of:
  - a) a polypeptide encoded by a polynucleotide sequence as shown in any one of SEQ ID NOS: 21-26 or 33-38 or 45;
  - b) a polypeptide as shown in any one of SEQ ID NOS: 27-32 or 39-44;
  - c) the mature form of the polypeptide as shown in any one of SEQ ID NOS: 27-32 or 39-44;
  - d) a polypeptide encoded by a polynucleotide sequence which hybridizes under highly stringent conditions to a polynucleotide sequence encoding (a) (b) or (c); and
  - e) a polypeptide having at least 90% sequence identity to the polypeptide of (b).
- 2 An immunogenic composition comprising an immunologically effective amount of at least one polypeptide of embodiment 1.
- 3 An isolated antibody that specifically binds the polypeptide of embodiment 1.
- 4 A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the polypeptide of embodiment 1 in a physiologically acceptable carrier.
- 5 A recombinant influenza virus comprising the polypeptide of embodiment 1.
- 6 An immunogenic composition comprising an immunologically effective amount of the recombinant influenza virus of embodiment 5.
- 7 A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the recombinant influenza virus of embodiment 5 in a physiologically acceptable carrier.

TABLE 3-continued

Specific embodiments
<p>8 An isolated nucleic acid, wherein said nucleic acid is selected from the group consisting of:</p> <p>a) a polynucleotide sequence as shown in any one of SEQ ID NOS: 21-26 or 33-38 or 45, or a complementary sequence thereof;</p> <p>b) a polynucleotide sequence encoding a polypeptide as shown in any one of SEQ ID NOS: 27-32 or 39-44, or a complementary polynucleotide sequence thereof;</p> <p>c) a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of polynucleotide sequence (a); and</p> <p>d) a polynucleotide sequence having at least 98% sequence identity to the polynucleotide sequence of (a).</p> <p>9 An immunogenic composition comprising at least one of the nucleic acids of embodiment 8.</p> <p>10 A cell comprising at least one nucleic acid of embodiment 8.</p> <p>11 A vector comprising the nucleic acid of embodiment 8.</p> <p>12 The vector of embodiment 12, wherein the vector is a plasmid, a cosmid, a phage, a virus, or a fragment of a virus.</p> <p>13 The vector of embodiment 12, wherein the vector is an expression vector.</p> <p>14 A cell comprising the vector of embodiment 13.</p> <p>15 An influenza virus comprising one or more nucleic acids of embodiment 8.</p> <p>16 The virus of embodiment 15, wherein the virus is a reassortment virus.</p> <p>17 A 6:2 reassortment influenza virus, wherein said virus comprises 6 gene encoding regions from A/Ann Arbor/6/60 and 2 gene encoding regions that encode polypeptides selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44.</p> <p>18 A method of producing a recombinant influenza virus, the method comprising: culturing the cell of embodiment 14 in a suitable culture medium under conditions permitting expression of nucleic acid; and, isolating the recombinant influenza virus from a cell population comprising said cell or the medium.</p> <p>19 An immunogenic composition comprising an immunologically effective amount of the recombinant influenza virus of embodiment 17.</p> <p>20 A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the recombinant influenza virus of embodiment 17 in a physiologically effective carrier.</p> <p>21 A method of producing an isolated or recombinant polypeptide, the method comprising: culturing the host cell of embodiment 10 in a suitable culture medium under conditions permitting expression of said nucleic acid; and, isolating the polypeptide from one or more of the host cells or the medium.</p> <p>22 A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, a virus of embodiment 17 in an amount effective to produce an immunogenic response against the viral infection.</p> <p>23 The method of embodiment 22, wherein the subject is a human.</p> <p>24 The immunogenic composition of embodiment 19, wherein the hemagglutinin comprises a modified polybasic cleavage site.</p> <p>25 A live attenuated influenza vaccine comprising the composition of embodiment 19.</p> <p>26 A split virus or killed virus vaccine comprising the composition of embodiment 19.</p> <p>27 A live attenuated influenza vaccine comprising the composition of embodiment 24.</p> <p>28 A split virus or killed virus vaccine comprising the composition of embodiment 24.</p> <p>29 A method for producing influenza viruses in cell culture, the method comprising:</p> <p>i) introducing into a population of host cells, which population of host cells is capable of supporting replication of influenza virus, a plurality of vectors comprising nucleic acid encoding at least 6 internal genome segments of a first influenza strain, wherein the first influenza strain is A/Ann Arbor/6/60; and, at least one genome segment encoding an immunogenic influenza surface antigen of a second influenza strain, wherein said second strain is a pandemic influenza strain,</p> <p>ii) culturing the population of host cells at a temperature less than or equal to 35° C.; and,</p> <p>iii) recovering a plurality of influenza viruses.</p> <p>30 The method of embodiment 29, wherein the plurality of vectors comprise at least one isolated nucleic acid, wherein said nucleic acid is selected from the group consisting of:</p> <p>a) a polynucleotide sequence of one of SEQ ID NOS: 21-26 or 33-38, or 45, or a complementary sequence thereof;</p> <p>b) a polynucleotide sequence encoding a polypeptide of one of SEQ ID NOS: 27-32 or 39-44, or a complementary polynucleotide sequence thereof;</p> <p>c) a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of polynucleotide sequence (a); and</p> <p>d) a polynucleotide sequence having at least 98% sequence identity to the polynucleotide sequence of (a).</p> <p>31 An immunogenic composition comprising an immunologically effective amount of the influenza virus of embodiment 29.</p> <p>32 An immunogenic composition comprising an immunologically effective amount of the influenza virus of embodiment 30.</p> <p>33 A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus of embodiment 29 in a physiologically effective carrier.</p>

TABLE 3-continued

Specific embodiments	
34	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus of embodiment 30 in a physiologically effective carrier.
35	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of embodiment 31.
36	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of embodiment 32.
37	A live attenuated influenza vaccine comprising the immunogenic composition of embodiment 31.
38	A split virus or killed virus vaccine comprising the immunogenic composition of embodiment 32.

[0250]

TABLE 4

Specific embodiments.	
1	An isolated polypeptide, wherein said polypeptide is selected from the group consisting of: a) a polypeptide comprising the amino acid sequence encoded by the nucleotide sequence as shown in any one of SEQ ID NOS: 21-26 or 33-38 or 45; b) a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44; c) the mature form of a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44; d) a polypeptide comprising an amino acid sequence encoded by a polynucleotide which hybridizes under highly stringent conditions to a polynucleotide comprising a nucleotide sequence encoding (a) (b) or (c); and e) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the polypeptide of (b).
2	An immunogenic composition comprising an immunologically effective amount of at least one polypeptide of embodiment 1.
3	An isolated antibody that specifically binds the polypeptide of embodiment 1.
4	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the polypeptide of embodiment 1 in a physiologically acceptable carrier.
5	A recombinant influenza virus comprising the polypeptide of embodiment 1.
6	An immunogenic composition comprising an immunologically effective amount of the recombinant influenza virus of embodiment 5.
7	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the recombinant influenza virus of embodiment 5 in a physiologically acceptable carrier.
8	An isolated polynucleotide, wherein said polynucleotide is selected from the group consisting of: a) a polynucleotide comprising the nucleotide sequence as shown in any one of SEQ ID NOS: 21-26 or 33-38 or 45, or a complementary sequence thereof; b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44, or a complementary nucleotide sequence thereof; c) a polynucleotide which hybridizes under highly stringent conditions over substantially the entire length of the polynucleotide of (a); and d) a polynucleotide comprising a nucleotide sequence having at least 98% sequence identity to the polynucleotide of (a).
9	An immunogenic composition comprising at least one polynucleotide of embodiment 8.
10	A cell comprising at least one polynucleotide of embodiment 8.
11	A vector comprising the polynucleotide of embodiment 8.
12	The vector of embodiment 11, wherein the vector is a plasmid, a cosmid, a phage, a virus, or a fragment of a virus.
13	The vector of embodiment 12, wherein the vector is an expression vector.
14	A cell comprising the vector of embodiment 13.
15	An influenza virus comprising one or more polynucleotides of embodiment 8.
16	The virus of embodiment 15, wherein the virus is a reassortant virus.

TABLE 4-continued

	Specific embodiments.
17	A 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from A/Ann Arbor/6/60 and 2 genome segments that encode an HA and/or a NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44.
18	A method of producing a reassortant influenza virus, the method comprising: culturing the cell of embodiment 14 in a suitable culture medium under conditions permitting expression of said polynucleotide; and, isolating the reassortant influenza virus from a cell population comprising said cell or the medium.
19	An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of embodiment 17.
20	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of embodiment 17 in a physiologically effective carrier.
21	A method of producing an isolated or recombinant polypeptide, the method comprising: culturing the cell of embodiment 10 in a suitable culture medium under conditions permitting expression of said polynucleotide; and, isolating the polypeptide from the cell or the medium.
22	A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, the virus of embodiment 17 in an amount effective to produce an immunogenic response against the viral infection.
23	The method of embodiment 22, wherein the subject is a human.
24	The immunogenic composition of embodiment 19, wherein the hemagglutinin comprises a modified polybasic cleavage site.
25	A live attenuated influenza vaccine comprising the composition of embodiment 19.
26	A split virus or killed virus vaccine comprising the composition of embodiment 19.
27	A live attenuated influenza vaccine comprising the composition of embodiment 24.
28	A split virus or killed virus vaccine comprising the composition of embodiment 24.
29	A method for producing an influenza virus in cell culture, the method comprising: i) introducing into a population of host cells, which population of host cells is capable of supporting replication of influenza virus, a plurality of vectors comprising nucleotide sequences corresponding to at least 6 internal genome segments of A/Ann Arbor/6/60; and, at least one genome segment comprising a polynucleotide encoding an HA and/or a NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44, ii) culturing the population of host cells at a temperature less than or equal to 35° C.; and, iii) recovering an influenza virus.
30	The method of embodiment 29, wherein the polynucleotide encoding the HA and/or NA polypeptide is selected from the group consisting of: a) a polynucleotide comprising the nucleotide sequence of any one of SEQ ID NOS: 21, 23-26 or 33-38, or 45, or a complementary nucleotide sequence thereof; b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44, or a complementary nucleotide sequence thereof; c) a polynucleotide which hybridizes under highly stringent conditions over substantially the entire length of the polynucleotide of (a); and d) a polynucleotide comprising a nucleotide sequence having at least 98% sequence identity to the polynucleotide of (a).
31	An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of embodiment 29.
32	An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of embodiment 30.
33	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus produced by the method of embodiment 29 in a physiologically effective carrier.
34	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus produced by the method of embodiment 30 in a physiologically effective carrier.
35	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of embodiment 31.
36	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of embodiment 32.
37	A live attenuated influenza vaccine comprising the immunogenic composition of embodiment 31.
38	A split virus or killed virus vaccine comprising the immunogenic composition of embodiment 32.

TABLE 4-continued

Specific embodiments.	
39	A 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from one or more donor viruses other than A/Ann Arbor/6/60 and 2 genome segments that encode an HA and/or a NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44.
40	The 6:2 reassortant influenza virus of embodiment 39, wherein said donor virus comprises one or more of the following phenotypes: temperature-sensitive, cold-adapted, or attenuated.
41	The 6:2 reassortant influenza virus of embodiment 39, wherein said donor virus is PR8.
42	The 6:2 reassortant influenza virus of embodiment 39, wherein said donor virus is A/Leningrad/17.
43	An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of embodiment 39.
44	An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of embodiment 40.
45	An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of embodiment 41.
46	An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of embodiment 42.
47	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of embodiment 39 in a physiologically effective carrier.
48	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of embodiment 40 in a physiologically effective carrier.
49	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of embodiment 41 in a physiologically effective carrier.
50	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of embodiment 42 in a physiologically effective carrier.
51	A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, the virus of embodiment 39 in an amount effective to produce an immunogenic response against the viral infection.
52	A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, the virus of embodiment 41 in an amount effective to produce an immunogenic response against the viral infection.
53	A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, the virus of embodiment 42 in an amount effective to produce an immunogenic response against the viral infection.
54	The method of embodiment 51, wherein said virus is killed or inactivated.
55	The method of embodiment 52, wherein said virus is killed or inactivated.
56	The method of embodiment 53, wherein said virus is killed or inactivated.
57	The immunogenic composition of embodiment 43, wherein the hemagglutinin comprises a modified polybasic cleavage site.
58	The immunogenic composition of embodiment 44, wherein the hemagglutinin comprises a modified polybasic cleavage site.
59	The immunogenic composition of embodiment 45, wherein the hemagglutinin comprises a modified polybasic cleavage site.
60	The immunogenic composition of embodiment 46, wherein the hemagglutinin comprises a modified polybasic cleavage site.
61	The method of embodiment 47, wherein the subject is a human.
62	The method of embodiment 48, wherein the subject is a human.
63	The method of embodiment 49, wherein the subject is a human.
64	A live attenuated influenza vaccine comprising the composition of embodiment 45.
65	A live attenuated influenza vaccine comprising the composition of embodiment 46.
66	A method for producing an influenza virus in cell culture, the method comprising: i) introducing into a population of host cells, which population of host cells is capable of supporting replication of influenza virus, a plurality of vectors comprising nucleotide sequences corresponding to: (a) at least 6 internal genome segments of a first influenza strain, wherein the first influenza strain is not A/Ann Arbor/6/60; and, at least one genome segment encoding an HA or an NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44; or (b) at least 6 internal genome segments of a first influenza strain, wherein the first influenza strain is not A/Ann Arbor/6/60 and which influenza strain comprises one or more phenotypic attributes selected from the group consisting of: attenuated, cold adapted and temperature sensitive; and, at least one genome segment encoding an HA or an NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS: 27-32, and 39-44,



TABLE 4-continued

Specific embodiments.	
	ii) culturing the population of host cells at a temperature less than or equal to 35° C.; and,
	iii) recovering an influenza virus.
67	An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of embodiment 66.
68	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus produced by the method of embodiment 66 in a physiologically effective carrier.
69	A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of embodiment 67.
70	A live attenuated influenza vaccine comprising the immunogenic composition of embodiment 67.
71	A split virus or killed virus vaccine comprising the immunogenic composition of embodiment 67.

## SEQUENCES

ca A/Vietnam/1203/04

[0251] Nucleotide Sequence of ca A/Vietnam/1203/04H5  
(SEQ ID NO:1)

[0252] Entire molecule length: 1767 nt

```

1  agcaaaagca ggggttcaat ctgtcaaaat ggagaaaata gtgcttcttt
51  ttgcaatagt cagtcctgtt aaaagtgatc agatttgcac tggttaccat
101 gcaaaacact cgacagagca ggttgacaca ataattggaaa agaactgtac
151 tggtacacat gccaagaca tactggaaaa gaaacacaac gggaagctct
201 gcgatctaga tggagtgaag cctctaattt tgagagattg tagcgtagct
251 ggatggctcc tcggaaaccc aatgtgtgac gaattcatca atgtgccgga
301 atggctttac atagtggaga aggccaatcc agtcaatgac ctctgttacc
351 caggggattt caatgactat gaagaattga aacacctatt gagcagaata
401 aaccattttg agaaaattca gatcatcccc aaaagttctt ggtccagtca
451 tgaagcctca ttaggggtga gctcagcatg tccataccag ggaaagtcct
501 cctttttcag aaatgtggtg tggcttatca aaaagaacag tacataccca
551 acaataaaga ggagctacaa taataccaac caagaagatc ttttgggtact
601 gtgggggatt caccatccta atgatgcggc agagcagaca agctctatc
651 aaaacccaac cacctatatt tccgttggga catcaacact aaaccagaga
701 ttggtaccaa gaatagctac tagatccaaa gtaaacgggc aaagtggaag
751 gatggagttc ttctggacaa ttttaaagcc gaatgatgca atcaacttcg
801 agagtaatgg aaatttcatt gctccagaat atgcatacaa aattgtcaag
851 aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa
901 caccaagtgt caaactccaa tgggggagat aaactctagc atgccattcc
951 acaatataca cctctcacc attggggaat gccccaaata tgtgaaatca
1001 aacagattag tccttgcgac tgggctcaga aatagccctc aaagagagac
1051 tcgaggatta ttggagcta tagcaggttt tatagaggga ggatggcagg

```

-continued

1101 gaatggtaga tggttggtat ggggtaccacc atagcaatga gcaggggagt  
1151 gggtagcgtg cagacaaaga atccactcaa aaggcaatag atggagtcac  
1201 caataaggtc aactcgatca ttgacaaaat gaacactcag tttgaggccg  
1251 ttggaaggga atttaacaac ttagaaagga gaatagagaa tttaacaag  
1301 aagatggaag acgggttcct agatgtctgg acttataatg ctgaacttct  
1351 ggttctcatg gaaaatgaga gaactctaga ctttcatgac tcaaattgtca  
1401 agaaccttta cgacaaggtc cgactacagc ttagggataa tgcaaaggag  
1451 ctgggtaacg gttgtttcga gttctatcat aaatgtgata atgaatgtat  
1501 ggaaagtgtg agaaatggaa cgtatgacta cccgcagtat tcagaagaag  
1551 cgagactaaa aagagaggaa ataagtggag taaaattgga atcaatagga  
1601 atttaccaaa tactgtcaat ttattctaca gtggcgagtt ccctagcact  
1651 ggcaatcatg gtagctggtc tatccttatg gatgtgctcc aatgggtcgt  
1701 tacaatgcag aatttgcatt taaatttggt agttcagatt gtagttaaaa  
1751 acacccttgt ttctact

[0253] Amino acid sequence of ca A/Vietnam/1203/04H5  
(SEQ ID NO:11)

[0254] Entire molecule length: 564 aa

1 mekivllfai vslvksdqic igyhannste qvdtimeknv tvthaqdile  
51 kkhngklcdl dgvkplilrd csvagwllgn pmcdefinvp ewsyivekan  
101 pvndlcypgd fndyeelkhl lsrinhfeki qiipksswss heaslgvssa  
151 cpyqgkssff rnvvwlikkn styptikrsy nntnqedllv lwgihhpnda  
201 aeqtikyqnp ttyisvgtst lnqrlvpria trskvngqsg rmeffwtilk  
251 pndainfesn gnfiapayay kivkkgdsti mkseleygnc ntkcqtpmga  
301 inssmpfhni hplrtigecpk yvksnrlvla tglrnsqpre trglfgaiag  
351 fieggwqgmv dgwygyhhsn eqgsgyaadk estqkaidgv tnkvnsiidk  
401 mntqfeavgr efnnerrie nlnkkmedgf ldwvtynael lvlmenertl  
451 dfhdsnvknl ydkvrlqlrd nakelngcgf efyhkcdnec mesvrngtyd  
501 ypqyseearl kreeisgvkl esigiyqils iystvassla laimvagls  
551 wmcsgslqc rici

[0255] Nucleotide Sequence of ca A/Vietnam/1203/04 N1  
(SEQ ID NO: 2)

[0256] Entire molecule length: 1398 nt

1 agcaaaagca ggagttcaaa atgaatccaa atcagaagat aataaccatc  
51 gggtagaatct gtatggtaac tggaatagtt agcttaatgt tacaaattgg  
101 gaacatgatc tcaatatggg tcagtcattc aattcacaca gggaatcaac  
151 accaatctga accaatcagc aataactaatt ttcttactga gaaagctgtg

-continued

201 gcttcagtaa aattagcggg caattcatct ctttgcccca ttaacggatg  
251 ggctgtatac agtaaggaca acagtataag gatcgggtcc aagggggatg  
301 tgtttgttat aagagagccg ttcactctcat gctcccactt ggaatgcaga  
351 actttctttt tgactcaggg agccttgctg aatgacaagc actccaatgg  
401 gactgtcaaa gacagaagcc ctacacagaac attaatgagt tgtcctgtgg  
451 gtgaggctcc ctccccatat aactcaaggt ttgagtctgt tgcttggtca  
501 gcaagtgcct gccatgatgg caccagttgg ttgacgattg gaatttctgg  
551 cccagacaat ggggctgtgg ctgtattgaa atacaatggc ataataacag  
601 acatatcaa gagttggagg aacaacatac tgagaactca agagtctgaa  
651 tgtgcatgtg taaatggctc ttgctttact gtaatgactg acggaccaag  
701 taatggtcag gcatcacata agatcttcaa aatggaaaaa gggaaagtgg  
751 ttaaatacgt cgaattggat gctcctaatt atcactatga ggaatgctcc  
801 tgttatccta atgccggaga aatcacatgt gtgtgcaggg ataattggca  
851 tggctcaaat cggccatggg tatctttcaa tcaaaatttg gagtatcaaa  
901 taggatatat atgcagtga gttttcggag acaatccacg ccccaatgat  
951 ggaacaggta gttgtgtgcc ggtgtcctct aacggggcat atggggtaaa  
1001 agggttttca tttaaatacg gcaatgggtg ctggatcggg agaaccaaaa  
1051 gcactaatc caggagcggc ttgaaatga ttgggatcc aaatgggtgg  
1101 actgaaacgg acagtagctt ttcagtga aaagatatcg tagcaataac  
1151 tgattggtea ggatatacg ggagtttgt ccagcatcca gaactgacag  
1201 gactagattg cataagacct tgtttctggg ttgagttgat cagagggcgg  
1251 cccaaagaga gcacaatttg gactagtggg agcagcatat ctttttggg  
1301 tgtaaatagt gacactgtgg gttggtcttg gccagacggt gctgagttgc  
1351 cattcaccat tgacaagtag tttgttcaa aaactccttg tttctact

[0257] Amino acid sequence of ca A/Vietnam/1203/04 N1  
(SEQ ID NO:12)

[0258] Entire molecule length: 449 aa

1 mnpnqkiiti gsicmvtgiv slmlqignmi siwvshsiht gnqhqssepis  
51 ntnfltekav asvklagnss lopingwavy skdnsirigs kgdvfvirep  
101 fiscshlecr tffltqgall ndkhsngtvk drsphrtlms cpvgeapsy  
151 nsrfesvaws asachdgtsw ltigisgpdn gavavlkyng iitdtikswr  
201 nniltqese cacvngscft vmtdgpengq ashkifkmeq gkvvksveld  
251 apnyhyeecs cypnageitc vcrdnwhgsn rpwvsfnqnl eyqigyicsg  
301 vfgdnprpnd gtgscgpvss ngaygvkgfs fkyngvwig rtkstnsrsq  
351 femiwdpangw tetdssfsvk qdivaitdws gysgsfvqhp eltglcdirp  
401 cfwvelirgr pkestiwtsg ssisfcgvns dtvgswpdg aelpftidk

ca A/Hong Kong/213/03

[0259] Nucleotide Sequence of ca A/Hong Kong/213/03H5 (SEQ ID NO: 3)

[0260] Entire molecule length: 1767 nt

```
1  agcaaaagca ggggttcaat ctgtcaaaat ggagaaaata gtgcttcttt
51  ttgcaatagt cagtcttggt aaaagtgatc agatttgcat tggttaccat
101 gcaaaacaact cgacagagca ggttgacaca ataatggaaa agaacgttac
151 tgttacacat gcccaagaca tactggaaaa gacacacaac gggaagctct
201 gcgatctaga tggagtgaag cctctaattt tgagagattg tagtgtagct
251 ggatggctcc tcggaacccc aatgtgtgac gaattcatca atgtgccgga
301 atggctttac atagtggaga aggccaatcc agccaatgac ctctgttacc
351 caggggattt caacgactat gaagaattga aacacctatt gagcagaata
401 aaccattttg agaaaattca gatcatcccc aaaaattctt ggtccagtca
451 tgaagcctca ttaggggtga gctcagcatg tccataccaa ggaaagtctt
501 cctttttcag gaatgtggtg tggcttatca aaaagaacaa tgcataccca
551 acaataaaga ggagctacaa taataccaac caagaagatc ttttgggtatt
601 gtgggggatt caccatccta atgatgcggc agagcagact aggctctatc
651 aaaacccaac cacctacatt tccgttgga catcaacact aaaccagaga
701 ttggtaccaa aaatagctac tagatccaaa gtaaacgggc aaaatggaag
751 gatggagttc ttctggacaa ttttaaaacc gaatgatgca atcaacttcg
801 agagcaatgg aaatttcatt gctccagaat atgcatacaa aattgtcaag
851 aaaggggact cagcaattat gaaaagtga ttggaatatg gtaactgcaa
901 caccaagtgt caaactccaa tgggggcgat aaactctagt atgccattcc
951 acaatataca cctctcacc atcggggaat gccccaaata tgtgaaatca
1001 aacagattag tccttgcgac tgggctcaga aatagccctc aaagagagac
1051 tcgaggatta tttggagcta tagcaggttt tatagagga ggatggcagg
1101 gaatggtaga tggttggtat gggtagcacc atagcaatga gcaggggagt
1151 gggtagcgtg cagacaaaga atccactcaa aaggcaatag atggagtcat
1201 caataaggtc aactcgatc ttagacaaat gaacactcag tttgaggccg
1251 ttggaaggga atttaataac ttagaaagga gaatagagaa tttaaacaag
1301 aagatggaag acggattcct agatgtctgg acttataatg ctgaacttct
1351 ggttctcatg gaaaatgaga gaactctaga ctttcatgac tcaaagtca
1401 agaaccttta cgacaaggtc cgactacagc ttagggataa tgcaaaggag
1451 ctgggtaacg gttgtttcga gttctatcac aaatgtgata atgaatgtat
1501 ggaaagtgtg agaaacggaa cgtatgacta cccgcagtat tcagaagaag
1551 caagactaaa aagagaggaa ataagtggag taaaattgga gtcaatagga
1601 acttaccaaa tactgtcaat ttattctaca gtggcgagtt ccctagcact
1651 ggcaatcatg gtagctggtc tatctttatg gatgtgctcc aatgggtcgt
```

-continued

1701 tacaatgcag aatttgcatt taaatttggt agttcagatt gtagttaaaa  
1751 acacccttqt ttctact

[0261] Amino acid sequence of ca A/Hong Kong/213/  
03H5 (SEQ ID NO:13)

[0262] Entire molecule length: 564 aa

1 mekivllfai vslvksdqic igyhannste qvdtimeknv tvthaqdile  
51 kthngklcdl dgvkplilrd csvagwllgn pmcdefinvp ewsyivekan  
101 pandlcypgd fndyeelkhl lsrinhfeki qiipknswws heaslgvssa  
151 cpyqgkssff rnvvwlkkn nayptikrsy nntnqedllv lwgihhpnda  
201 aeqtrlyqnp ttyisvgtst lnqrlvpkia trskvngqng rmeffwtilk  
251 pndainfesn gnfiapayay kivkkgdsai mkseleygnc ntckqtpmga  
301 inssmpfhni hpltigecpk yvksnrlvla tglrnspgre trglfgaiag  
351 fiegwwqgmw dgwygyhhsn eggsgyaadk estqkaidgv tnkvnsiidk  
401 mntqfeavgr efnnlerrrie nlnkkmedgf ldvwtynael lvlmenertl  
451 dfhdnsvknk ydkvrlqlrd nakelngngcf efyhkcdnec mesvrngtyd  
501 ypyyseearl kreeisgvkl esigtyqils iystvassla laimvagls1  
551 wmcsgslqc rici

[0263] Nucleotide Sequence of ca A/Hong Kong/213/03  
N1 (SEQ ID NO: 4)

[0264] Entire molecule length: 1458 nt

1 agcaaaagca ggagttcaaa atgaatccaa atcagaagat aacaaccatt  
51 ggatcaatct gtatggtaat tggaatagtt agcttgatgt tacaaattgg  
101 gaacataatc tcaatatggg ttagtcattc aattcaaaca gggaatcaac  
151 accaggctga accatgcaat caaagcatta ttacttatga aaacaacacc  
201 tgggtaaacc agacatatgt caacatcagc aataccaatt ttcttactga  
251 gaaagctgtg gcttcagtaa cattagcggg caattcatct ctttgcccca  
301 ttagtggatg ggctgtatac agtaaggaca acggtataag aatcggttcc  
351 aagggggatg tgtttgttat aagagagccg ttcattctcat gctcccactt  
401 ggaatgcaga actttctttt tgactcaggg agccttgetg aatgacaagc  
451 attctaattg gaccgtcaaa gacagaagcc ctacagaac attaatgagt  
501 tgtcccgtag gtgaggtcc ttcccatac aactcgaggt ttgagtctgt  
551 tgcttggtcg gcaagtgtt gtcattgatg cactagttag ttgacaattg  
601 gaatttctgg ccagacaat ggggctgtgg ctgtattgaa atacaatggc  
651 ataataacag aactatcaa gagttggagg aacaacataa tgagaactca  
701 agagtctgaa tgtgcatgtg taaatggctc ttgctttact gttatgactg  
751 atggaccaag taatgggcag gcttcataca aaatcttcag aatagaaaaa

-continued

801 gggaaagtag ttaaatcagc cgaattaaat gcccctaatt atcactatga  
 851 ggagtgtcc tggtatcctg atgctggaga aatcacatgt gtgtgcaggg  
 901 ataactggca tggctcaaat cggccatggg tatctttcaa tcaaaatttg  
 951 gagtatcgaa taggatatat atgcagtgga gttttcggag acaatccacg  
 1001 ccccaatgat gggacaggca gttgtgtcc ggtgtcccct aaaggggcat  
 1051 atggaataaa agggttctca tttaaatacg gcaatggtgt ttggatcggg  
 1101 agaaccacaaa gcactaattc caggagcggc ttgaaatga ttgggatcc  
 1151 aaatggatgg actggtacgg acagtaattt ttcagtaaag caagatattg  
 1201 tagctataac cgattggta ggatatagcg ggagttttgt ccagcatcca  
 1251 gaactgacag gattagattg cataagacct tgtttctggg ttgagctaat  
 1301 cagagggcgg cccaaagaga gcacaatttg gactagtggg agcagcatat  
 1351 ccttttgtgg tgtaaatagt gacactgtgg gttggtcttg gccagacggt  
 1401 gctgagttgc cattcaccat tgacaagtag tttgttcaaa aaactccttg  
 1451 tttctact

[0265] Amino acid sequence of ca A/Hong Kong/213/03  
 N1 (SEQ ID NO:14)

[0266] Entire molecule length: 469 aa

1 mnpnqkitti gsicmvigiv slmlqignii siwvshsiqt gnqhqaepcn  
 51 qsiityennt wvnqtyvnis ntnfltekav asvtlagncs lcpisgwavy  
 101 skdngirigs kgdvvfirep fiscshlecr tffltqgall ndkhsngtvk  
 151 drsphrtlms cpvgeapsy nsrfesvaws asachdgtsw ltigisgpdn  
 201 gavavlk yng iitdtikswr nnimrtqese cacvngscft vmtdgpsngq  
 251 asykifriek gkvvksaeln apnyhyeecs cypdageitc vcrdnwhgsn  
 301 rpwvsnfnql eyrigyicsg vfgdnprpnd gtgscgpvsp kgaygikgfs  
 351 fkyngngwig rtkstnsrsg femiwdpngw tgtdsnfsvk qdivaitdws  
 401 gysgsvfghp eltgldcirp cfwvelirgr pkestiwtsg ssisfcgvns  
 451 dtvgwswpdg aelpftidk

ca A/Hong Kong/491/97 (HA)+A/Hong Kong/486/97 (NA)

[0267] Nucleotide Sequence of ca A/Hong Kong/491/  
 97H5 (SEQ ID NO: 5)

[0268] Entire molecule length: 1767 nt

1 agcaaaagca ggggtataat ctgtcaaaat ggagaaaata gtgcttcttc  
 51 ttgcaacagt cagccttggt aaaagtgacc agatttgcatt tggttacat  
 101 gcaaacact cgacagagca agttgacaca ataattggaaa agaattgtac  
 151 tgttacacat gccaagaca tactggaaa gacacacaac gggaagctct  
 201 gcgatctaaa tggagtgaag cctctgattt tgagggattg tagttagct

-continued

251 ggatggctcc tcggaacccc tatgtgtgac gaattcatca atgtgccgga  
301 atgggtcttac atagtggaga aggccagtc agccaatgac ctctgttatc  
351 caggggaattt caacgactat gaagaactga aacacctatt gagcagaata  
401 aaccatthttg agaaaattca gataatcccc aaaagttctt ggtccaatca  
451 tgatgcctca tcaggggtga gctcagcatg tccatacctt gggaggtcct  
501 cctttttcag aaatgtggta tggcttatca aaaagaacag tagctaccca  
551 acaataaaga ggagctacaa taataccaac caagaagatc ttttgggtact  
601 gtgggggatt caccatccta atgatgcggc agagcagaca aggcctctatc  
651 aaaacccaac cacctacatt tccgttgga catcaacact gaaccagaga  
701 ttggttccag aaatagctac tagacccaaa gtaaacgggc aaagtggag  
751 aatggagtgc ttctggacaa ttttaaagcc gaatgatgcc atcaatttcg  
801 agagtaatgg aaatttcatt gctccagaat atgcatacaa aattgtcaag  
851 aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa  
901 caccaagtgt caaactccaa tgggggcaat aaactctagt atgccattcc  
951 acaacataca cccctcacc atcggggaat gccccaaata tgtgaaatca  
1001 aacagattag tccttgcaac tggactcaga aataccctc aacgagagac  
1051 gcgaggacta tttggagcta tagcaggttt tatagaggga ggatggcagg  
1101 gaatggtaga tggttggtat gggtagcacc atagcaatga gcaggggagt  
1151 ggatacgtg cagaccaaga atccacaaa aaggcaatag atggagtac  
1201 caataaggtc aactcgatca ttaacaaaat gaacactcag tttgaggccg  
1251 ttggaaggga atttaataac ttggaaagga ggatagagaa tttaaacaag  
1301 aaaatggaag acggattcct agatgtctgg acttacaatg ccgaacttct  
1351 ggttctcatg gaaaatgaga gaactctaga ctttcatgac tcaaattgca  
1401 agaaccttta cgacaaggtc cgactacagc ttagggataa tgcaaaggag  
1451 ctgggtaatg gttgtttcga attctatcac aaatgtgata acgaatgtat  
1501 ggaaagtgtg aaaaacggaa cgtatgacta cccgcagtat tcagaagaag  
1551 caagactaaa cagagaggaa ataagtggag taaaattgga atcaatggga  
1601 acttaccaaa tactgtcaat ttattcaaca gtggcgagtt ccctagcact  
1651 ggcaatcatg gtagctggtc tatctttatg gatgtgctcc aatggatcgt  
1701 tacaatgcag aatttgcatt taaatttggt agttcagatt gtagttaaaa  
1751 acacccttgt ttctact

[0269] Amino acid sequence of ca A/Hong Kong/491/  
97H5 (SEQ ID NO:15)

[0270] Entire molecule length: 564 aa

1 mekivllllat vslvksdqic igyhannste qvdtimeknv tvthaqdile  
51 rthngklcdl ngvklplilrd csvagwllgn pmcdefinvp ewsyivekas  
101 pandlccypgn fndyeelkhl lsrinhfeki qiipksswsn hdassgvssa

-continued

151 cpylgrssff rnvvlikkn ssyptikrsy nntnqedllv lwgihhpnda  
201 aeqtrlyqnp ttyisvgtst lnqrlvpeia trpkvngqsg rmeffwtilk  
251 pndainfesn gnfiapeyay kivkkgdsti mkseleygnc ntkcqtpmga  
301 inssmpfhni hpltigecpk yvksnrlvla tglrntpqre trglfgaiag  
351 fieggwqgmv dgwygyhhsn egsggyaadq estqkaidgv tnkvnsiink  
401 mntqfeavgr efnnerrie nlnkkmedgf ldvwtynael lvlmenertl  
451 dfhdenvknl ydkvrlqlrd nakelngngcf efyhkdneec mesvkngtyd  
501 ypqyseearl nreeisgvkl esmgtyqils iystvassla laimvagls1  
551 wmcsgslqc rici

[0271] Nucleotide Sequence of ca A/Hong Kong/486/97  
N1 (SEQ ID NO: 6)

[0272] Entire molecule length: 1401 nt

1 agcaaaagca ggagtttaaa atgaatccaa atcagaagat aataaccatt  
51 ggatcaatct gcatggtagt tgggataatc agcttgatgt tacaatttgg  
101 aaacacaata tcagtatggg tcagccacat aattaaaact tggcacccaa  
151 accagcctga accatgcaac caaagcatca atttttacac tgagcaggct  
201 gcagcttcag tgacattagc gggcaattcc tctctctgcc ctattagtgg  
251 atgggctata tacagcaagg acaatagtat aagaattggt tccaaagggg  
301 atgtgtttgt tataagagaa ccattcatct catgctccca ttggaatgc  
351 agaacctttt tcttgaccca aggagcccta ttgaatgaca agcattctaa  
401 tgggaccgtc aaagacagga gccctatag aactttaatg agctgtcctg  
451 ttggtgaggc cccctcccca tacaactcaa ggtttgagtc tgttgcttgg  
501 tcagcaagtg cttgccatga tggcattagt tggctaacaa ttggaatttc  
551 cggtcgggat aatggggtg tggctgtgtt gaaatacaat ggcataataa  
601 cagacaccat caagagttgg aggaacaaca cactgaggac gcaagagtct  
651 gaatgtgcat gtgtgaatgg ttcttgtttt actgtaatga cagatggacc  
701 gagtaatgaa caggcctcat acaagatttt caagatagaa aaggggaggg  
751 tagtcaaate agttgagttg aacgccccta attatcatta cgaggaaatgc  
801 tcctgttate ctgatgctgg cgaaatcaca tgtgtgtgca gggataattg  
851 gcatggctcg aaccgaccat ggggtgtctt caatcagaat ctggagtatc  
901 aaataggata tatatgcagt ggggttttcg gagacagtcc acgcccacat  
951 gatgggacag gcagttgtgg tccagtgtct cttaacggag cgtatggagt  
1001 aaaagggttt tcatttaaat acggcaatgg tgtttgatc gggagaacca  
1051 aaagcactag ttccaggagc ggttttgaaa tgatttggga tccaaatggg  
1101 tggaccgaaa cagacagtag cttctcgttg aagcaagaca tcatagcgat  
1151 aactgattgg tcaggataca gcgggagttt tattcaacat ccagaactga  
1201 caggattaaa ttgcatgaga ccttgcttct gggttgaact aatcagaggg



-continued

1251 aggcccaaaag agaaaaacaat ctggactagt gggagcagta tatctttctg  
1301 tgggtgtaaat agtgacactg tgggttggtc ttggccagac ggtgctgagt  
1351 tgccatacac cattgacaag tagtttggtc aaaaaactcc ttgtttctac  
1401 t

[0273] Amino acid sequence of ca A/Hong Kong/486/97  
N1 (SEQ ID NO:16)

[0274] Entire molecule length: 450 aa

1 mnpnqkiiti gsicmvvgii slmlqignti svwvshiikt whpnqpepcn  
51 qsinfyteqa aasvtlagns slcpisgwai yskdnsirig skgdvfvire  
101 pfiscshlec rtffltqgal lndkhsngtv kdrspyrtlm scpvgeapsp  
151 ynsrfesvaw sasachdgis wltigisgpd ngavavlkyn giitdtiksw  
201 rnntlrtqes ecacvngscf tvmtdgpsne qasykifkie kgrvvksvel  
251 napnyhyeec scypdageit cvcrdnwhgs nrpwvsfnqn leyqigyics  
301 gvfgdsprpn dgtgscgpvs lngaygvkgf sfkygngvwi grtkstssrs  
351 gfemiwdpng wtetdssfs1 kqdiiatdw sgysgsfihq peltglncmr  
401 pcfwvelirg rpkektiws gssisfcgvn sdtvgwswpd gaelpytidk

ca A/Hong Kong/491/97 (Ser211) (HA)+ca A/Hong Kong/  
486/97 (NA)

[0275] Nucleotide Sequence of ca A/Hong Kong/491/97  
(Ser211) H5 (SEQ ID NO: 7)

[0276] Entire molecule length: 1767 nt

1 agcaaaagca ggggtataat ctgtcaaaat ggagaaaata gtgcttcttc  
51 ttgcaacagt cagccttggt aaaagtgacc agatttgcat tggttaccat  
101 gcaaacact cgacagagca agttgacaca ataatggaaa agaattgtac  
151 tgttacacat gcccaagaca tactggaaag gacacacaac gggaagctct  
201 gcgatctaaa tggagtgaag cctctgattt tgagggattg tagtgtagct  
251 ggatggctcc tcggaaaccc tatgtgtgac gaattcatca atgtgccgga  
301 atggtcttac atagtggaga aggccagtcc agccaatgac ctctgttata  
351 caggggaattt caacgactat gaagaactga aacacctatt gacgagaata  
401 aaccattttg agaaaattca gataatcccc aaaagttcctt ggtccaatca  
451 tgatgctca tcaggggtga gctcagcatg tccatacctt gggaggtcct  
501 cctttttcag aaatgtggta tggcttatca aaaagaacag tagctaccca  
551 acaataaaga ggagctacaa taataccaac caagaagatc ttttggtagt  
601 gtgggggatt caccatccta atgatgcggc agagcagaca aggtctctatc  
651 aaaacccaac cacctacatt tccgttgga catcaacact gaaccagaga  
701 ttggtttcag aaatagctac tagacccaaa gtaaacgggc aaagtggaag

-continued

751 aatggagttc ttctggacaa ttttaaagcc gaatgatgcc atcaatttcg  
801 agagtaatgg aaatttcatt gctccagaat atgcatacaa aattgtcaag  
851 aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa  
901 caccaagtgt caaactccaa tgggggcaat aaactctagt atgccattcc  
951 acaacataca cccctcacc atcggggaat gcccacaata tgtgaaatca  
1001 aacagattag tccttgcaac tggactcaga aataccctc aacgagagac  
1051 gcgaggacta tttggagcta tagcaggttt tatagagga ggatggcagg  
1101 gaatggtaga tggttggtat gggtaaccacc atagcaatga gcaggggagt  
1151 ggatacgctg cagaccaaga atccacacaa aaggcaatag atggagtcac  
1201 caataaggtc aactcgatca ttaacaaaat gaacactcag tttgagggcg  
1251 ttggaaggga atttaataac ttggaagga ggatagagaa tttaacaag  
1301 aaaatggaag acggattcct agatgtctgg acttacaatg ccgaacttct  
1351 ggttctcatg gaaaatgaga gaactctaga ctttcatgac tcaaattgtca  
1401 agaaccttta cgacaaggtc cgactacagc ttagggataa tgcaaaggag  
1451 ctgggtaatg gttgtttcga attctatcac aaatgtgata acgaatgtat  
1501 ggaaagtgtg aaaaacggaa cgtatgacta cccgcagtat tcagaagaag  
1551 caagactaaa cagagaggaa ataagtgag taaaattgga atcaatggga  
1601 acttaccaaa tactgtcaat ttattcaaca gtggcgagtt ccctagcact  
1651 ggcaatcatg gtagctggtc tatctttatg gatgtgctcc aatggatcgt  
1701 tacaatgcag aatttgcatt taaatttggt agttcagatt gtagttaaaa  
1751 acacccttgt ttctact

[0277] Amino acid sequence of ca A/Hong Kong/491/97  
(Ser211) H5 (SEQ ID NO:17)

[0278] Entire molecule length: 564 aa

1 mekivlllat vslvksdqic igyhannste qvdtimeknv tvthaqdile  
51 rthngklcdl ngvklplilrd csvagwllgn pmcdefinvp ewsyivekas  
101 pandlcypgn fndyeelkhl lsrinhfeki qiipksswsn hdassgvssa  
151 cpylgrssff rnvvwlikkn ssyptikrsy nntnqedllv lwgihhpnda  
201 aeqtrlyqnp ttyisvgtst lnqrlvseia trpkvngqsg rmeffwtilk  
251 pndainfesn gnfiapayay kivkkgdsti mkseleygnc ntkcqtpmga  
301 insmpfhni hpltigecpk yvksnrlvla tglrntpqre trglfgaiag  
351 fiegwwgmw dgwygyhhsn eggsgyaadq estqkaidgv tnkvnsiink  
401 mntqfeavgr efnnerrie nlnkkmedgf ldvwtynael lvlmenertl  
451 dfhdsvknkl ydkvrlqlrd nakelngcgf efyhkcdnec mesvkngtyd  
501 ypqyseearl nreeisgvkl esmgtqyils iystvassla laimvagls1  
551 wmcsgslqc rici

[0279] Nucleotide Sequence of ca A/Hong Kong/486/97  
N1 (SEQ ID NO: 8)

[0280] Entire molecule length: 1401 nt

```
1  agcaaaagca ggagtttaaa atgaatccaa atcagaagat aataaccatt
51  ggatcaatct gcatggtagt tgggataatc agcttgatgt tacaatttgg
101  aaacacaata tcagtatggg tcagccacat aattaaaact tggcacccaa
151  accagcctga accatgcaac caaagcatca atttttacac tgagcaggct
201  gcagcttcag tgacattagc gggcaattcc tctctctgcc ctattagtgg
251  atgggctata tacagcaagg acaatagtat aagaattggg tccaaagggg
301  atgtgtttgt tataagagaa ccattcatct catgctccca ttggaatgc
351  agaacctttt tcttgaccca aggagcccta ttgaatgaca agcattctaa
401  tgggaccgtc aaagacagga gccctatag aactttaatg agctgtcctg
451  ttggtgaggc cccttcccca tacaactcaa ggtttgagtc tgttgcttgg
501  tcagcaagtg cttgccatga tggcattagt tggctaacaa ttggaatttc
551  cggtcctgat aatggggctg tggctgtgtt gaaatacaat ggcataataa
601  cagacaccat caagagttgg aggaacaaca cactgaggac gcaagagtct
651  gaatgtgcat gtgtgaatgg ttcttgtttt actgtaatga cagatggacc
701  gagtaatgaa caggcctcat acaagatttt caagatagaa aaggggaggg
751  tagtcaaadc agttgagttg aacgccccta attatcatta cgaggaatgc
801  tcctgttadc ctgatgctgg cgaaatcaca tgtgtgtgca gggataattg
851  gcatggctcg aaccgacat ggggtgtctt caatcagaat ctggagtatc
901  aaataggata tatatgcagt ggggttttcg gagacagtcc acgcccacat
951  gatgggacag gcagttgtgg tccagtgtct cttaacggag cgtatggagt
1001  aaaagggttt tcatttaaat acggcaatgg tgtttgatc gggagaacca
1051  aaagcactag ttccaggagc gggtttgaaa tgatttggga tccaaatggg
1101  tggaccgaaa cagacagtag cttctcgttg aagcaagaca tcatagcgat
1151  aactgattgg tcaggataca gcgggagttt tattcaacat ccagaactga
1201  caggattaaa ttgcatgaga ccttgcttct ggggtgaact aatcagaggg
1251  aggcccaaag agaaaacaat ctggactagt gggagcagta tatctttctg
1301  tgggtgaaat agtgacactg tgggttggtc ttggccagac ggtgctgagt
1351  tgccatacac cattgacaag tagtttgttc aaaaaactcc ttgtttctac
1401  t
```

[0281] Amino acid sequence of ca A/Hong Kong/486/97  
N1 (SEQ ID NO:18)

[0282] Entire molecule length: 450 aa

```
1  mnpnqkiiti gsicmvvgii slmlqignti svwvshiikt whpnqpepcn
51  qsinfytega aasvtaglgn slcpisgwai yskdnsirig skgdvfvire
101  pfiscshlec rtffltqgal lndkhsngtv kdrspyrtrlm scpvgeapsp
```

-continued

151 ynsrfesvaw sasachdgis wltigisgpd ngavavlkyn giitdtiksw  
201 rnntlirtqes ecacvngscf tvmtgdpsne qasykifkie kgrvvksvel  
251 napnyhyeec scypdageit cvcrdnwhgs nrpwvsfnqn leyqigyics  
301 gvfgdsprpn dgtgscgpvs lngaygvkgf sfkyngngvwi grtkstssrs  
351 gfemiwdpng wtetdssfs1 kqdiiatdw sgysgsfiqh peltglncmr  
401 pcfwvelirg rpkektiws gssisfcgvn sdtvgwswpd gaelpyidk

ca A/ck/Hong Kong/G9/97

[0283] Nucleotide sequence of ca A/ck/Hong Kong/G9/97  
(SEQ ID NO: 9)

[0284] Entire molecule length: 1690 bp

1 ttaaccactc aagatggaag caataccact aataactata ctactagtag  
51 taacagcaag caatgcagac aaaatctgca tcggctacca atcaacaaac  
101 tccacagaaa ccgtagacac gctaacagaa aacaatgttc ctgtgacaca  
151 tgccaaagaa ttgctccaca cagagcacia tgggatgctg tgtgcaacaa  
201 atctgggacg tcctcttatt ctagacactt gcaccattga aggactgatc  
251 tatggcaacc cttcttgtga tctactgttg ggaggaagag aatggtccta  
301 catcgctgaa agaccatcgg ctgttaatgg aatgtgttac cccgggaatg  
351 tagaaaacct agaggaacta aggtcatttt ttagttctgc tagttcctac  
401 caaagaatcc agatctttcc agacacaatc tggaatgtgt cttacagtgg  
451 aacaagcaaa gcatgttcag attcattcta caggagcatg agatggttga  
501 ctcaaaagaa caacgcttac cctattcaag acgccaata cacaataat  
551 agaggaaaga gcattctttt catgtggggc ataaatcacc cacctaccga  
601 tactgcacag acaaatctgt acacaaggac tgacacaaca acaagtgtgg  
651 caacagaaga tataaatagg accttcaaac cagtgatagg gccaggcccc  
701 cttgtcaatg gtctgcaggg aagaattgat tattattggt cggatttgaa  
751 accaggtcag acattgcgag taagatcaa tgggaatcta atcgctccat  
801 ggtatgggca cattctttca ggagagagcc acggaagaat cctgaagact  
851 gatttaaaaca gtggtagctg tgtagtgcaa tgtcaaacag aaagagggtg  
901 cttaataact actttgccat tccacaatgt cagtaaatat gcatttggaa  
951 actgccccaa atatgttggg gtaaagagtc tcaaactggc agttggtctg  
1001 aggaatgtgc ctgctagatc aagtagagga ctatttgggg ccatagettg  
1051 attcatagag ggaggttggg cagggctggt cgctggttgg tatgggttcc  
1101 agcattcaaa tgatcaaggg gttggtatag ctgcagatag agactcaact  
1151 caaagggcaa ttgacaaaat aacgtccaaa gtgaataata tagtcgataa  
1201 aatgaacaag caatatgaaa ttattgatca tgaattcagc gaggttgaaa  
1251 atagactcaa tatgatcaat aataagattg atgaccagat acaagacata

-continued

1301 tgggcatata acgctgaatt gctagtgtctg cttgaaaacc agaaaacact  
1351 cgatgagcat gatgcgaatg taaacaatct atataacaaa gtgaagaggg  
1401 cactgggttc caatgcaatg gaagatggga aaggatgttt cgagctatac  
1451 cataaatgtg atgatcagtg catggagaca attcggaaacg ggacctataa  
1501 caggaggaag tataaagagg aatcaagact agaaagacag aaaatagaag  
1551 gggccaagct ggaatctgaa ggaacttaca aaatcctcac cattttattcg  
1601 actgtcgct catctctgt gattgcaatg gggtttctg ccttcttgtt  
1651 ctgggccatg tccaatggat cttgcagatg caacatttga

[0285] Amino acid sequence of ca A/ck/Hong Kong/G9/  
97H9 (SEQ ID NO:19)

[0286] Entire molecule length: 558 aa

1 meaiplitil lvvtasnadk icigyqstns tetvdtlten nvpvthakel  
51 lhtehngmlc atnlgrplil dtctieglyy gnpscdlllg grewsyiver  
101 psavngmcp gnvenleelr sffssassyq riqifpdtiw nvsygskska  
151 csdsfyrsmr wltqknnayp iqdaqytnnr gksilfmwgi nhpptdtaqt  
201 nlytrtdttt svatedinrt fkpvigprpl vnglqgridy ywsvlkpgqt  
251 lrvrsgnli apwyghilsg eshgrilktl lnsqscvvc qtergglntt  
301 lpfhnskyia fgncpkyvgv kslklavglr nvparssrgl fgaiaqfieg  
351 gwsglvagwy gfqhsndqgv giaadrstq raidkitskv nnivdkmnkq  
401 yeiidhefse venrlminn kiddqiqdiw aynaellvll enqktldehd  
451 anvnnlynkv kralgsname dgkgcfelyh kcdqcmeti rngtynrrky  
501 keesrlrqk iegvkleseg tykiltiyst vasslviamg faaflfwams  
551 ngscrcni

[0287] Nucleotide sequence of ca A/ck/Hong Kong/G9/97  
N2 (SEQ ID NO:10)

[0288] Entire molecule length: 1428 bp

1 aaatgaatcc aaatcagaag ataatagcaa ttggctctgt ttctctaact  
51 attgcgacaa tatgcctcct catgcagatt gctatcttag caacgactat  
101 gacactacat ttcaagcaga atgaatgcat caactcctcg aataatcaag  
151 tagtgccatg tgaaccaatc ataatagaaa ggaacataac agagatagtg  
201 catttgaata gtactacctt agagaaggaa atttgccta aagtagcaga  
251 ctacaggaat tgggtcaaac cacaatgtca aatcacaggg ttcgctcctt  
301 tctccaagga caattcaatt aggtctctcg cagggtggaga tatttgggtg  
351 acaagagaac cttatgtatc gtgcgggtctt ggtaaatgtt atcaatttgc  
401 acttgggcag ggaaccactt tggagaacaa aactcaaac ggcacagcac

-continued

451 atgatagaac tcctcataga acccttttaa tgaatgagtt ggggtgttccg  
501 ttctatttgg caaccaaaca agtgtgcata gcatgggccca gctcaagctg  
551 ccatgatggg aaagcatggt tacatgtttg tgtcactggg gatgatagaa  
601 atgcaacggc tagcatcatt tatgatggga tacttgttga cagtattggt  
651 tcatggctca aaaacatcct cagaactcag gagtcagaat gcgtttgcat  
701 caatggaacc tgtgcagtag taatgactga tggaagtgca tcaggaaggg  
751 ctgacactag aatactatctt attagagagg ggaaaattgc acacattagc  
801 ccattgtcag gaagtgtcga gcatgtggag gaatgtcctt gttacccccg  
851 atatccagaa gttagatgtg ttgacagaga caattggaag ggatccaata  
901 ggcccgttct atatatataat atggcaaatt atagtattga ttccagttat  
951 gtgtgctcag gacttggttg cgacacacca agaaatgatg ataggtctag  
1001 cagcagcaac tgcagagatc ctaataacga gagagggggc ccaggagtaa  
1051 aaggggtggc ctttgacaat ggaaatgaca ttgggatggg aagaacaatc  
1101 aaaaaggatt cgcgctcagg ttatgagact ttcagggtca ttggtggttg  
1151 gaccactgct aattccaagt cacagataaa tagacaagtc atagttgaca  
1201 gtgataactc gtctgggtat tctggtatct tctctgttga aggcaaaagc  
1251 tgcatacaaca ggtgttttta cgtggagttg ataagaggaa gaccaaagga  
1301 gactaggggtg tgggtggactt caaatagcat cattgtatct tgtggaactt  
1351 caggtaccta tggaacaggc tcatggcctg atggggcgaa tatcaatttc  
1401 atgcctatat aagcttttcgc aattttag

[0289] Amino acid sequence of ca A/c/Hong Kong/G9/  
97 N2 (SEQ ID NO: 20)

[0290] Entire molecule length: 469 aa

1 mnpnqkii ai gsvsltiati cllmqiaila tmttlhfkqn ecinsennqv  
51 vpcepiiier niteivhl ns ttlekeicpk vadyrnwskp qcqitgfpaf  
101 skdnsirlsa ggdiwvtrep yvscglgkcy qfalgqgttl enkhngtah  
151 drtphrllm nelgvpfhla tkqvciawss sschdgkawl hvcvtgddrn  
201 atasilydgi lvdsgswsk nhlrtqesec vcingtcavv mtdgsasgra  
251 dtrilfireg kiahisplsg saqhveecsc ypripevrcv crdnwkgsnr  
301 pvlyinmany sidssyvcsg lvgdtprndd rsssnncrdp nnergapgvk  
351 gwafdnngndi wmgtrtkkds rsgyetfrvi ggwtansks qinrqivids  
401 dnssgygif svegkscinr cfyvelirgr pketrvwts nsiivfcgts  
451 gtygtgswpd ganinfmpi

A/Netherlands/219/03

[0291] Nucleotide sequence of A/Netherlands/219/03H7  
(SEQ ID NO: 21)

[0292] Entire molecule length: 1737 bp

```
1  agcaaaagca ggggatacaa aatgaacact caaatcctgg tattcgctct
51  ggtggcgagc attccgacaa atgcagacaa gatctgcctt gggcatcatg
101 ccggtgtcaaa cgggactaaa gtaaacacat taactgagag aggagtggaa
151 gtcgttaatg caactgaaac ggtggaacga acaaacgttc ccaggatctg
201 ctcaaaaggg aaaaggacag ttgacctcgg tcaatgtgga cttctgggaa
251 caatcactgg gccaccccaa tgtgaccaat tcctagaatt ttcgcccgac
301 ttaattattg agaggcgaga aggaagtgat gtctgttata ctgggaaatt
351 cgtgaatgaa gaagctctga ggcaaattct cagagagtca ggcggaattg
401 acaaggagac aatgggattc acctacagcg gaataagaac taatggaaca
451 accagtgcac gtaggagatc aggatcttca ttctatgcag agatgaaatg
501 gtcctgttca aacacagaca atgctgcttt cccgcaaatg actaagtcac
551 acaagaacac aaggaaagac ccagctctga taatatgggg gatccaccat
601 tccggatcaa ctacagaaca gaccaagcta tatgggagtg gaaacaaact
651 gataacagtt gggagttcta attaccaaca gtcctttgta ccgagtcacg
701 gagcgagacc acaagtgaat ggccaatctg gaagaattga ctttcattgg
751 ctgatactaa accctaatac cacggtcact ttcagtttca atggggcctt
801 catagctcca gaccgtgcaa gctttctgag agggaagtcc atgggaattc
851 agagtgaagt acaggttgat gccaatgtg agggagattg ctatcatagt
901 ggagggacaa taataagtaa ttgcccctt cagaacataa atagcagggc
951 agtaggaaaa tgtccgagat atgttaagca agagagtctg ctgttggtgaa
1001 caggaatgaa gaatgttccc gaaatcccaa agaggaggag gagaggccta
1051 tttggtgcta tagcgggttt cattgaaaat ggatgggaag gtttgattga
1101 tgggtggtat ggcttcaggg atcaaaatgc acaaggggag ggaactgctg
1151 cagattacaa aagcacccaa tcagcaattg atcaaataac agggaaatta
1201 aatcggctta tagaaaaaac taaccaacag tttgagtaa tagacaacga
1251 attcactgag gttgaaaggg aaattggcaa tgtgataaac tggaccagag
1301 attccatgac agaagtgtgg tcctataacg ctgaactctt agtagcaatg
1351 gagaatcagc acacaattga tctggccgac tcagaaatga acaaactgta
1401 cgaacgagtg aagagacaac tgagagagaa tgccgaagaa gatggcactg
1451 gttgcttcca aatatttcac aagtgtgatg acgactgcac ggccagtatt
1501 agaaacaaca cctatgatca cagcaagtac agggaagaag caatacaaaa
1551 tagaatacag attgaccacg tcaaaactaag cagcggctac aaagatgtga
1601 tactttgggt tagcttcggg gcatcatgtt tcatacttct ggccattgca
1651 atgggccttg tcttcatatg tgtgaagaat ggaacatgc ggtgcactat
1701 ttgtatataa gtttgaaaaa acacccttgt ttctact
```

[0293] Amino acid sequence of A/Netherlands/219/03H7  
(SEQ ID NO: 27)

[0294] Entire molecule length: 562 aa

```
1  mntqilvfal vasiptnadk iclghhavs n gtkvntlter gvevvnatet
51  vertnvpric skgkrtvdlg qcglgtitg ppqcdqflef sadliierre
101 gsdvcypgkf vneearqil resggidket mgftysgirt ngtsacrrs
151 gssfyaemkw llstndnaaf pqmtksyknt rkdpaliwg ihhsgstteq
201 tklygsnkl itvgssnyqq sfvpspgarp qvngqsgrid fhwlilnpnd
251 tvtfsfngaf iapdrasflr gksmgisqev qvdancegdc yhsggtiisn
301 lpfqninsra vgkcpvryvkq eslllatgmk nvpeipkrrr rglfgaiagf
351 iengweglid gwygfrhna qgegtaadyk stqsaidqit gklnrliect
401 nqqfelidne fteverqign vinwtrdsmt evwsynaell vamenqhtid
451 ladsemnkly ervkrqlren aeedgtgcfe ifhkcdcdcm asirnttydh
501 skyreeaign riqidpvkls sgykdvilwf sfgascfill aiamglvfic
551 vkngnmrcti ci
```

[0295] Nucleotide Sequence of A/Netherlands/219/03 N7  
(SEQ ID NO: 22)

[0296] Entire molecule length: 1465 nt

```
1  agcaaaagca gggatgatcg gaataaatcc aaatcagaaa ctatttgcac
51  tatctggagt ggcaatagca cttagtgtac tgaacttatt gataggaatc
101 tcaaacgtcg gattgaacgt atctctacat ctaaaggaaa aaggacccaa
151 acaggaggag aatttaacat gcacgacccat taatcaaaac aacactactg
201 tagtagaaaa cacatatgta aataatacaa caataattac caagggaact
251 gatttgaaaa caccaagcta tctgctgttg aacaagagcc tgtgcaatgt
301 tgaaggggtg gtcgtgatag caaaagacaa tgcagtaaga tttggggaaa
351 gtgaacaaat cattgttacc agggagccat atgtatcatg cgaccaaca
401 ggatgcaaaa tgtatgcctt gcaccaaggg actaccatta ggaacaaaca
451 ttcaaatgga acgattcatg acagaacagc tttcagaggt ctcatctcca
501 ctccattggg cactccacca accgtaagta acagtgaact tatgtgtgtt
551 ggatgggtcaa gcacaacttg ccatgatggg attgctagga tgactatctg
601 tatacaagga aataatgaca atgctacagc aacggtttat tacaacagaa
651 ggctgaccac taccattaag acctgggcca gaaacattct gaggactcaa
701 gaatcagaat gtgtgtgcca caatggcaca tgtgcagttg taatgaccga
751 cggatcggtc agtagtcaag cctatacaaa agtaaatgtat ttccacaagg
801 gattagtagt taaggaggag gagttaaggg gatcagccag acatattgag
851 gaatgctcct gttatggaca caatcaaaag gtgacctgtg tgtgcagaga
901 taactggcag ggagcaaaac gccctattat agaaattgat atgagcacat
951 tggagcacac aagtagatag gtgtgcactg gaattctcac agacaccagc
```



-continued

1001 agacctgggg acaaatctag tggatgtgt tccaatccaa taactgggag  
1051 tcccggcggt cccggagtga agggattcgg gtttctaaat ggggataaca  
1101 catggcttgg taggaccatc agcccagat caagaagtgg attcgaaatg  
1151 ttgaaaatac ctaatgcagg tactgatccc aattctagaa tagcagaacg  
1201 acaggaaatt gtcgacaata acaattgggc aggcatttcc ggaagcttta  
1251 ttgactattg gaatgataac agtgaatgct acaatccatg cttttacgta  
1301 gagttaatta gaggaagacc cgaagaggct aaatacgtat ggtgggcaag  
1351 taacagtcta attgccctat gtggaagccc attccagtt gggctcgtgt  
1401 ccttccccga tggggcacia atccaatact tttcgtaaaa tgcaaaaaaa  
1451 ctcttggtt ctact

[0297] Nucleotide Sequence of A/Netherlands/219/03 N7  
(SEQ ID NO: 45)

[0298] Entire molecule length: 1464 nt

1 agcaaaagca gggatgatga gaatgaatcc aaatcagaaa ctatttgcac tatctggagt  
61 ggcaatagca cttagtgtac tgaacttatt gataggaatc tcaaacgtcg gattgaacgt  
121 atctctacat ctaaggaaa aaggacccaa acaggaggag aatttaacat gcacgaccat  
181 taatcaaaac aacactactg tagtagaaaa cacatatgta aataatacaa caataattac  
241 caagggaact gatttgaaaa caccaagcta tctgctgttg aacaagagcc tgtgcaatgt  
301 tgaagggtgg gtcgtgatag caaaagacaa tgcagtaaga tttggggaaa gtgaacaaat  
361 cattgttacc agggagccat atgtatcatg cgaccaaca ggatgcaaaa tgtatgcctt  
421 gcaccaaggg actaccatta ggaacaaaca ttcaaatgga acgattcatg acagaacagc  
481 tttcagaggt ctcatctcca ctccattggg cactccacca accgtaagta acagtgactt  
541 tatgtgtgtt ggatgggtcaa gcacaacttg ccatgatggg attgctagga tgactatctg  
601 tatacaagga aataatgaca atgctacagc aacggtttat tacaacagaa ggctgaccac  
661 taccattaag acctgggcca gaaacattct gaggactcaa gaatcagaat gtgtgtgcca  
721 caatggcaca tgtgcagttg taatgaccga cggatcggct agtagtcaag cctatacaaa  
781 agtaatgtat ttccacaagg gattagtagt taaggaggag gagttaaggg gatcagccag  
841 acatattgag gaatgctcct gttatggaca caatcaaaag tgacactgtg tgtgcagaga  
901 taactggcag ggagcaaaaca ggcctattat agaaattgat atgagcacat tggagcacac  
961 aagtagatac gtgtgcactg gaattctcac agacaccagc agacctgggg acaaatctag  
1021 tggatgtgtt tccaatccaa taactgggag tcccggcggt cccggagtga agggattcgg  
1081 gtttctaaat ggggataaca catggcttgg taggaccatc agcccagat caagaagtgg  
1141 attcgaaatg ttgaaaatac ctaatgcagg tactgatccc aattctagaa tagcagaacg  
1201 acaggaaatt gtcgacaata acaattgggc aggcatttcc ggaagcttta ttgactattg  
1261 gaatgataac agtgaatgct acaatccatg cttttacgta gagttaatta gaggaagacc  
1321 cgaagaggct aaatacgtat ggtgggcaag taacagtcta attgccctat gtggaagccc

-continued

1381 attcccagtt gggctctggt cctccccga tggggcacaa atccaatact ttctgtaaaa  
1441 tgcaaaaaca cccttgtttc tact

[0299] Amino acid sequence of A/Netherlands/219/03 N7  
(SEQ ID NO: 28)

[0300] Entire molecule length: 471 aa

1 mnpnqklfal sgvaialsvl nlligisnvg lnvslhlkek gpkqeenltc  
51 ttingnnttv ventyvntt iitkgtdlkt psylllnksl cnvegwwvia  
101 kdnavrfges eqiivtrepy vscdptgckm yalhggttir nkhsngtihd  
151 rtafrglist plgtppptvsn sdfmcvgwss ttchdgiarm ticiqgnndn  
201 atatvyyrrr ltttikttwar nilrtqesec vchngtcavv mtdgsassqa  
251 ytkvmyfhkg lvvkeelrg sarhieecsc yghnqkvctv crdnwqganr  
301 piieidmstl ehtsryvctg ilttdtsrpgd kssgdcnpi tgspgvpvgvk  
351 gfgfngdnt wlgrtisprs rsgfemlkip nagtdpnsri aergeivdnn  
401 nwsygsygf dywndnsecy npcfyvelir grpeeakyvw wasnslialc  
451 gspfpvsgs fpdgaqiyf s

A/ck/BC/CN-7/04

[0301] Nucleotide sequence of A/ck/BC/CN-7/04H7  
(SEQ ID NO: 23)

[0302] Entire molecule length: 1754 bp

1 agcaaaagca ggggatacaa aatgaatact caaatTTTtg cattcattgc  
51 ttgtatgctg attggaacta aggagacaa aatatgtctt gggcaccatg  
101 ctgtggcaaa tgggacaaaa gtgaacacac taacagagag ggaattgaa  
151 gtagtcaatg ccacggagac ggtggaaact gtaaataatta aaaaaaatg  
201 cactcaagga aaaaggccaa cagatctggg acaatgtgga cttctaggaa  
251 ccctaatagg acctcccaa tgcgatcaat ttctggagtt tgacgctaata  
301 ttgataattg aacgaagaga aggaaccgat gtgtgctatc ccgggaagtt  
351 cacaaatgaa gaatcactga ggcagatcct tcgagggta ggaaggaattg  
401 ataaagatc aatgggtttc acctatagtg gaataagaac caatggggcg  
451 acgagtgcct gcagaagatc aggttcttct ttctatgcgg agatgaagtg  
501 gttactgtcg aattcagaca atgcggcatt tccccaatg actaagtcgt  
551 ataggaatcc caggaacaaa ccagctctga taatctgggg agtgcacac  
601 tctggatcag ctactgagca gaccaaactc tatggaagtg gaaacaagtt  
651 gataacagta ggaagctcga aataccagca atcattcact ccaagtccgg  
701 gagcacggcc acaagtgaat ggacaatcag gaaggattga ttttcattgg

-continued

751 ctactccttg accccaatga cacagtgacc ttcactttca atggggcatt  
801 catagccctt gacagggcaa gtttctttag aggagaatcg ctaggagtcc  
851 agagtgatgt tcctttggat tctgggtgtg aaggggattg ctccacagt  
901 gggggtacga tagtcagttc cctgccattc caaaacatca accctagaac  
951 agtggggaaa tgcctcgcgt atgtcaaaca gacaagcctc cttttggcta  
1001 caggaatgag aaacgtccca gagaacccca agcaggccta ccggaacgg  
1051 atgaccagag gccttttttg agcgattgct ggattcatag agaattgatg  
1101 ggaaggtctc atcgatggat ggtatggttt cagacatcaa aatgcacaag  
1151 gagaaggaac tgcagctgac taaaaagca ccaatctgc aatagatcag  
1201 atcacaggca aattgaatcg tctgattgac aaaacaaacc agcagtttga  
1251 actgatatag aatgaattca gtgagataga acaacaaatc gggaaatgtca  
1301 ttaactggac acgagactca atgactgagg tatggtcgtg taatgctgag  
1351 ctggttggtg caatggagaa tcagcataca atagatcttg cagactcaga  
1401 aatgaacaaa ctttacgaac gcgtcagaaa acaactaagg gaaatgctg  
1451 aagaagatgg aactggatgc ttgagatat tccataagtg tgatgatcag  
1501 tgtatggaga gcataaggaa caacacttat gaccataccc aatacaggac  
1551 agagtcatcg cagaatagaa tacagataga cccagtgaag ttgagtagtg  
1601 gatacaaaga cataatctta tggtttagct tcggggcatt atgttttctt  
1651 cttctagcca ttgcaatggg attggttttc atttgcataa agaattgaaa  
1701 catgcggtgc actatgtgta tatagtttga gaaaaaaca cccttgtttc  
1751 tact

[0303] Amino acid sequence of A/ck/BC/CN-7/04H7  
(SEQ ID NO: 29)

[0304] Entire molecule length: 567 aa

1 mntqilafia cmligtkgdk iclghhavan gtkvntltter gievvnatet  
51 vetvnikkic tqgkrptdlg qcglgtlig ppqcdqlef danlillerre  
101 gtdvcypgkf tneeslrqil rgsggidkes mgftygirt ngatsacrrs  
151 gssfyamkw llsnsdnaaf pqmtksyrnp rnkpaliiwg vhhsgsateq  
201 tklysgnkl itvgeskyqq sftpspgarp qvngqsgrid fhwlldpnd  
251 tvtftfngaf iapdrasffr geslgvqsdv pldsgcegdh fhsggtivss  
301 lpfqninprt vgkcpvykq tslllatgmr nvpenpkqay rkrmtgrlfg  
351 aiagfiengw eglidgwygf rhqnaqgegt aadykstqsa idqitgklnr  
401 lidktnqqfe lidnefseie qqignvinwt rdsmtewvsw naellvamen  
451 qhtidladse mnklyervrk qlreneeeg tgcfefihkc ddqcmesirn  
501 ntydhtqyrt eslnriqid pvklssgykd iilwfsfgas cillaiamg  
551 lvficikngn mrtctici

[0305] Nucleotide Sequence of A/ck/BC/CN-7/04 N3  
(SEQ ID NO: 24)

[0306] Entire molecule length: 1453 nt

```
1 agcaaaagca ggtgcgagat gaatccgaat cagaagataa taacaatcgg
51 ggtagtgaat accactctgt caacaatagc ccttctcatt ggagtgggaa
101 acttagtttt caacacagtc atacatgaga aaataggaga ccatcaaata
151 gtgacccatc caacaataat gacccctgaa gtaccgaact gcagtgcac
201 tataataaca tacaataaca ctgttataaa caacataaca acaacaataa
251 taactgaagc agaaaggcct ttcaagtctc cactaccgct gtgccccttc
301 agaggattct tcccttttca caaggacaat gcaatacgac tgggtgaaaa
351 caaagacgtc atagtcacaa gggagcctta tgttagctgc gataatgaca
401 actgctggtc ctttgctctc gcacaaggag cattgctagg gactaaacat
451 agcaatggga ccattaaaga cagaacacca tataggcttc taattcggtt
501 cccaatagga acagctccag tactaggaaa ttacaaagag atatgcattg
551 cttggtcgag cagcagttgc ttgacggga aagagtggat gcatgtgtgc
601 atgacaggga atgataatga tgcaagtgcc cagataatat atggaggaag
651 aatgacagac tccattaaat catggaggaa ggacatacta agaaccagg
701 agtctgaatg tcaatgcatt gacgggactt gtgttggtgc tgtcacagat
751 ggccctgctg ctaatagtgc agatcacagg gtttactgga tacgggaggg
801 aagaataata agtatgaaa atgttcccaa aacaaagata caacacttag
851 aagaatgttc ctgctatgtg gacattgatg tttactgtat atgtagggac
901 aattggaagg gctctaacag accttgatg agaatcaaca acgagactat
951 actggaaaca ggatatgtat gtagtaaatt tctactcagac acccccagge
1001 cagctgaccc ttcaataatg tcatgtgact cccaagcaa tgtcaatgga
1051 ggaccggag tgaaggggtt tggtttcaaa gctggcaatg atgtatggtt
1101 aggtagaaca gtgtcaacta gtggtagatc gggtttgaa attatcaaag
1151 ttacagaagg gtggatcaac tctcctaacc atgtcaaac aattacacaa
1201 aactagtgt ccaacaatga ctggtcaggc tattcaggta gcttcattgt
1251 caaagccaag gactgttttc agccctgttt ttatgttgag cttatacgag
1301 ggaggcccaa caagaatgat gacgtctctt ggacaagtaa tagtatagtt
1351 actttctgtg gactagacaa tgaacctgga tcgggaaatt ggccagatgg
1401 ttctaacatt gggtttatgc ccaagtaata gaaaaagca cttgtttct
1451 act
```

[0307] Amino acid sequence of A/ck/BC/CN-7/04 N3  
(SEQ ID NO: 30)

[0308] Entire molecule length: 469 aa

```
1 mnpnqkiiti gvnnttlsti alligvgnlv fntvihekig dhqivthpti
51 mtpevpncsd tiitynntvi nnitttiite aerpfkspfp lcpfrgffpf
```

-continued

101 hkdnairlge nkdvivtrep yvscdndncw sfalaaggall gtkhsngtik  
 151 drtpyrslir fpigtapvlg nykeiciaws ssscfdgkew mhvcmtgndn  
 201 dasaqiiygg rmtdsikswr kdilrtqese cqcidgtcvv avtdgpaans  
 251 adhrvywire griikyenvp ktkiqhleec scyvdidvyc icrdnwkgsn  
 301 rpwmrinnet iletgyvcsk fhsdtpypad psimscdspd nvnggpgvkg  
 351 fgfkagndvw lgrtvstsgs sgfeiikvte gwinspnvhk sitqtlvsnn  
 401 dwsgygsfsi vkakdcfqpz fyvelirgrp nknddvsmts nsivtfcgld  
 451 nepgsgnwps gsnigfmpk

ca A/ck/BC/CN-6/04

[0309] Nucleotide sequence of ca A/ck/BC/CN-6/04H7  
 (SEQ ID NO: 25)

[0310] Entire molecule length: 1734 bp

1 agcgaaagca ggggatacaa aatgaatact caaatTTTtg cattcattgc  
 51 ttgtatgtcg attggaacta aaggagacaa aatatgtctt gggcaccatg  
 101 ctgtggcaaa tgggacaaaa gtgaacacac taacagagag gggaattgaa  
 151 gtagtcaatg ccacggagac ggtggaaact gtaaataatta agaaaaatg  
 201 cactcaagga aaaaggccaa cagatctggg acaatgtgga cttctaggaa  
 251 ccctaataag acctcccaa tgcgatcaat ttctggagtt tgacgcta  
 301 ttgataattg aacgaagaga aggaaccgat gtgtgctatc ccgggaagt  
 351 cacaatgaa gaatcactga ggcagatcct tcgagggta ggaggaattg  
 401 ataaagatg aatgggttcc acctatagtg gaataagaac caatggggcg  
 451 acgagtgcct gcagaagatc aggttcttct ttctatgcgg agatgaagt  
 501 gttactgtcg aattcagaca atgcggcatt tccccaaatg actaagtcgt  
 551 ataggaatcc caggaacaaa ccagctctga taatctgggg agtgcacac  
 601 tctggatcag ctactgagca gaccaaactc tatggaagt gaaacaagt  
 651 gataacagta ggaagctcga aataccagca atcattcact ccaagtccgg  
 701 gagcagggc acaagtgaat ggacaatcag gaaggattga ttttcattgg  
 751 ctactccttg accccaatga cacagtgacc ttcactttca atggggcatt  
 801 catagcccct gacagggcaa gtttctttag aggagaatcg ctaggagtcc  
 851 agagtgatgt tcctttggat tctggtgtg aaggggattg cttccacagt  
 901 ggggtacga tagtcagttc cctgccattc caaaacatca accctagaac  
 951 agtggggaaa tgcctcgat atgtcaaaca gacaagcctc cttttggcta  
 1001 caggaatgag aaacgtccca gagaaccca agaccagagg ctttttggga  
 1051 gcgattgtcg gattcataga gaatggatgg gaaggtctca tcgattgatg  
 1101 gtatggttcc agacatcaaa atgcacaagg agaaggaact gcagctgact  
 1151 acaaaagcac ccaatctgca atagatcaga tcacaggcaa attgaatcgt  
 1201 ctgattgaca aaacaaacca gcagtttgaa ctgatagaca atgaattcag

-continued

1251 tgagatagaa caacaaatcg ggaatgtcat taactggaca cgagactcaa  
 1301 tgactgaggt atggctgtat aatgctgagc tgttggtggc aatggagaat  
 1351 cagcatacaa tagatcttgc agactcagaa atgaacaaac ttacgaacg  
 1401 cgtcagaaaa caactaaggg aaaatgctga agaagatgga actggatgct  
 1451 ttgagatatt ccataagtgt gatgatcagt gtatggagag cataaggaac  
 1501 aacacttatg accataccca atacaggaca gagtcattgc agaatagaat  
 1551 acagatagac ccagtgaat tgagtagtgg atacaaagac ataatttat  
 1601 ggtttagctt cggggcatca tgttttcttc ttctagccat tgcaatggga  
 1651 ttggttttca ttgcataaa gaatggaaac atgcggtgca ctatttgtat  
 1701 atagtttgag aaaaaaacac ccttgtttct act

[0311] Amino acid sequence of ca A/ck/BC/CN-6/04H7  
 (SEQ ID NO: 31)

[0312] Entire molecule length: 560 aa

1mntqilafia cmligtkgdk iclghhavan gtkvntlter gievvnatet  
 51vetvnikkic tggkrptdlg qcgllgtlig ppqcdqflef danliierre  
 101gtdvcypgkf tneeslrqil rgsggidkes mgftysgirt ngatsacrrs  
 151gssfyamkw llnsdnaaf pqmtksyrnp rnkpaliiwg vhhsgsateq  
 201tklygsgnkl itvgsskyqq sftspgarp qvngqsgrid fhwlildpnd  
 251tvtftfngaf iapdrasffr geslgvqsdv pldsqcegdh fhsggtivss  
 301lpfqninprt vgkcpryvkq tslllatgmr nvpenpktrg lfgaiagfie  
 351ngweglidgw ygfrhqnagg egtaadykst qsaidqitgk lnrlidktnq  
 401qfelidnefs eieqqignvi nwtrdsmtew wsynaellva menqhtidla  
 451dsemnklyer vrkqlrenae edgtgcfeif hkoddqcmes irrnttydhtq  
 501yrteslqnri qidpvklssg ykdiilwfsf gascfllai amglvficik  
 551ngnmrctici

[0313] Nucleotide sequence of ca A/ck/BC/CN-6/04 N3  
 (SEQ ID NO: 26)

[0314] Entire molecule length: 1453 nt

1 agcaaaagca ggtgcgagat gaatccgaat cagaagataa taacaatcgg  
 51 ggtagtgaat accactctgt caacaatagc ccttctcatt ggagtgggaa  
 101 acttagtttt caacacagtc atacatgaga aaataggaga ccatcaaata  
 151 gtgacccatc caacaataat gaccctgaa gtaccgaact gcagtgcac  
 201 tataataaca tacaataaca ctgttataaa caacataaca acaacaataa  
 251 taactgaagc agaaaggcct ttcaagtctc cactaccgct gtgccccttc  
 301 agaggattct tcccttttca caaggacaat gcaatacgac tgggtgaaaa

-continued

351 caaagacgtc atagtcacaa gggagcetta tgttagctgc gataatgaca  
401 actgctggtc ctttgccttc gcacaaggag cattgctagg gactaaacat  
451 agcaatggga ccattaaaga cagaacacca tataggcttc taattcgttt  
501 cccaatagga acagctccag tactaggaaa ttacaaagag atatgcattg  
551 cttggtcgag cagcagttgc ttgacggga aagagtggat gcatgtgtgc  
601 atgacaggga atgataatga tgcaagtgcc cagataatat atggaggaag  
651 aatgacagac tccattaaat catggaggaa ggacatacta agaaccagg  
701 agtctgaatg tcaatgcatt gacgggactt gtgttggtgc tgtcacagat  
751 ggccctgctg ctaatagtgc agatcacagg gtttactgga tacgggaggg  
801 aagaataata aagtatgaaa atgttcccaa aacaaagata caacacttag  
851 aagaatgttc ctgctatgtg gacattgatg tttactgtat atgtagggac  
901 aattggaagg gctctaacag accttgatg agaatcaaca acgagactat  
951 actggaaaca ggatatgtat gtagtaaatt tctactcagac acccccaggc  
1001 cagctgaccc ttcaataatg tcatgtgact cccaagcaa tgtcaatgga  
1051 ggacccggag tgaaggggtt tggtttcaa gctggcaatg atgtatggtt  
1101 aggtagaaca gtgtcaacta gtggtagatc gggctttgaa attatcaaag  
1151 ttacagaagg gtggatcaac tctcctaacc atgtcaaac aattacacaa  
1201 aactagtgt ccaacaatga ctggtcaggc tattcaggta gcttcattgt  
1251 caaagccaag gactgttttc agccctgttt ttatgttgag cttatacgag  
1301 ggaggcccaa caagaatgat gacgtctctt ggacaagtaa tagtatagtt  
1351 actttctgtg gactagacaa tgaacctgga tcgggaaatt ggccagatgg  
1401 ttctaacatt gggtttatgc ccaagtaata gaaaaaagca ccttgtttct  
1451 act

[0315] Amino acid sequence of ca A/ck/BC/CN-6/04 N3  
(SEQ ID NO: 32)

[0316] Entire molecule length: 469 aa

1mnpnqkiiti gvnnttltsti alligvgnlv fntvihekig dhqivthpti  
51mtpevpncsd tiitynnvti nnitttiite aerpfkspfp lcpfrgffpf  
101hkdnairlge nkdvivtrep yvscdndncw sfalaqgall gtkhngtik  
151drtpyrslir fpigtapvlg nykeiciaws ssscfdgkew mhvcmtgndn  
201dasaqiiygg rmtdsikswr kdilrtqese cqcidgtcvv avtdgpaans  
251adhrvywire griikyenvp ktkiqhlee scyvdivdyc icrdnwksn  
301rpwmrinnet iletgyvcsk fhstdprpad psimscdps nvngpgpvkg  
351fgfkagndvw lgrtvstsgsr sgfeiikvte gwinspnvhk sitqtlvsnn  
401dwsygsfsi vkakdcfqp fyvelirgrp nknddsvwts nsivtfcgl  
451nepgsgnwpd gsnigfmpk

ca A/Duck

[0317] Nucleotide sequence of ca A/Duck H6 (SEQ ID NO: 33)

[0318] Entire molecule length: 1743 bp

```
1  agcaaaagca ggggaaaatg attgcagtca ttataatagc ggtactggca acggccggaa
61  aatcagacaa gatctgcatt gggatatcatg ccaacaattc aacaacacaa gtggatacga
121 tacttgagaa gaatgtaacc gtcacacact cagttgaatt gctggagaac caaaaagaag
181 aaagattctg caagatcttg aacaaggccc ctctcgattt aagaggatgt accatagagg
241 gttggatctt ggggaatccc caatgcgacc tattgcttgg tgatcaaagc tggatcatata
301 tagtggaag acctacagct caaaatggga tctgctaccc aggaattttg aatgaagtag
361 aagaactgaa ggcacttatt ggatcaggag aaagagtgga gagatttgaa atgtttccca
421 aaagtacatg ggcaggagta gacaccagca gtggggtaac aaaggcttgc ctttatacta
481 gtggttcgtc tttctacaga aacctcctat ggataataaa aaccaagtcc gcagcatatc
541 cagtaattaa gggaacctac aataaactg gaagccagcc aatcctctat ttctggggtg
601 tgcaccatcc tcctgacacc aatgagcaaa acactttgta tggctctggt gatcgatatg
661 tcaggatggg aactgaaagc atgaattttg ccaagagccc agaaattgag gcaaggcctg
721 ctgtgaatgg tcaaaagggc agaattgatt attactggtc tgttttaaag ccgggggaaa
781 ccttgaatgt ggaatctaata ggaaatctaa tcgcccttg gtatgcatac aaatttgta
841 gcaccaatag taaaggagcc gtcttcaagt caaatctacc aatcgagaac tgtgatgcca
901 catgccagac tattgcagga gtcttaagaa ccaataaaac atttcagaat gtaagccctc
961 tgtggatagg agaatgcccc aaatatgtga aaagtgaag tttgaggctt gcaactggac
1021 taagaaatat tccacagatt gagactagag gacttttcgg agctatcgca gggtttattg
1081 aaggaggatg gactggaatg atagatgggt ggtatggcta tcaccatgaa aattctcaag
1141 gctcagggta tgcggcagac agagaaagca ctcaaagggc tatagacgga attacaaata
1201 aggtcaattc cattatagac aaaatgaaca cacaattcga agctatagac cacgaattct
1261 caaatgtgga gagaagaatt gacagtctga acaaaagaat ggaagatgga tttctggacg
1321 tttggacata caatgctgaa ctgttggttc ttcttgaaaa cgaaaggaca ctagacctac
1381 atgacgcgaa tgtgaagaac ctgtatgaaa aggtcaaacc acaactacgg gacaatgcta
1441 atgatctagg aaatggatgc tttgaatttt ggcataagtg tgacaatgaa tgcataagat
1501 ctgtcaaaaa tggtaacctat gactatccca aatatcagga tgaaagcaaa ttgaacaggc
1561 aggaaataga atcggtgaag ctggagaacc ttgggtgtgta tcaaatcctc gccatttata
1621 gtacgggatc gagcagtcta gtcttggtag ggctgattat agcaatgggt ctttgatgt
1681 gttcaaatgg ttcaatgcaa tgcaggatat gtatataatt aagaaaaaca cccttggtct
1741 act
```



[0319] Amino acid sequence of ca A/Duck H6 (SEQ ID NO: 39)

[0320] Entire molecule length: 566 aa

```
1 miaviiiavl atagksdkic igyhannstt qvdtileknv tvthsvelle nqkeerfcki
61 lnkapldlrg ctiegwilgn pqcdlllgdq swsyiverpt aqngicypgi lneveelkal
121 igsgerverf emfpkstwag vdtssgvtkc cpytsgssfy rnllwiiktk saaypvikgt
181 ynntgspil yfwgvhphpd tneqntlygs gdryvmgte smnfakspei aarpavngqr
241 gridyywsvl kpgetlnves ngnlapwya ykfvstnsgk avfknlpie ncdatcqtia
301 gvlrtnktf nvsplwige pkyvkseslr latglnipq ietrglfgai agfieggtg
361 midgwygyhh ensqgsyaa drestqraid gitnkvnsl dkmntqfeai dhefsnlerr
421 idslnkrmed gldvwtyna ellvllener tldlhdanvk nlyekvksql rdnandlgn
481 cfefwhkcdn eciesvkngt ydypkyqdes klnrgeiesv klenlgvyqi laiystvsss
541 lylvgliiam glwmcsngsm qcrici
```

[0321] Nucleotide sequence of ca A/Duck N9 (SEQ ID NO: 34)

[0322] Entire molecule length: 1460 nt

```
1 agcaaaagca gggtaaat gaattccaaat cagaagattc tatgcacatc tgctactgcc
61 attgcaatag gcacaattgc tgtattaata ggaatagcaa acctgggttt gaacatagga
121 ctacacctga aaccgagctg caactgctcc aacctctc ctgaaacaac aatgtaagc
181 caaacaataa taaacaatta ctacaatgaa acaaatgtta cccaaataag taacacaaac
241 attcaacata tggggggaac cgaaaaggac ttcaacaatc tgactaaagg gctctgcaca
301 ataaattcat ggcatatatt cggaaaggac aatgctataa gaataggga gaactctgat
361 gtttttagtca caagagagcc atatgtttct tgtgatccag atgaatgcag attctatgct
421 ctacagccaag gaacaacaat acgggggaaag cactcaaatg gaacaataca cgatagatcc
481 caataccgtg ctttagtgag ctggccttta tcatcaccac ccactgtgta caataccaga
541 gtagaatgca ttggatggtc cagtacaagc tgccatgatg ggaaagcacg aatgtctata
601 tgtgtctcag gtcccaacaa caatgcatca gcagtgtttt ggtacaaagg gcggcctatc
661 acggaaatca atacgtgggc ccgaaacata ttgagaacct aagaatctga gtgtgtatgc
721 cacaatggaa tatgtccagt agtggtcact gacggttctg ccaccgttc agcagaaact
781 aggatatact atttcaaaga ggggaaaatc ctcaaatggg agccactaac tggaaccgcc
841 aagcacattg aagaatgctc ttgctatggg aaagactcag aaataacgtg cacatgtaga
901 gacaattggc aaggctcgaa tagaccagta atacaaataa accccacaat gatgactcac
961 actagtcaat acatatgcag cctgtctc acagacaatc cagcccccac tgacccacg
1021 gtaggcaagt gtaatgatcc ttatccagga aacaacaata atggagtcaa aggattctca
1081 tatttagatg gtgacaatac atggctagga agaacgataa gcacagctc taggtctggg
1141 tatgaaatgc tgaaagtgcc taatgcattg acagatgata gatcaaaacc tactcaagg
1201 cagacaattg tattaacac agactggagt gggtacagt ggtcttcat tgattactgg
1261 gcaaaagggg agtctatag agcatgcttc tacgttgagc tgatccgtgg aaggccaaaa
```

-continued

1321 gaggacaaaag tgtggtggac cagtaatagt atagtgtcga tgtgttccag cacagagttc  
1381 cttggacaat ggaactggcc agatggggct aaaatagagt acttcctcta agatgtagaa  
1441 aaaagaccct tgtttctact

[0323] Amino acid sequence of ca A/Duck N9 (SEQ ID  
NO: 40)

[0324] Entire molecule length: 470 aa

1mnpnqkilt sataiaigti avilgianlg lniglhlkps cncsnpppet tnvsqtiinn  
61yynetnvtqi sntniqhmvg tekdfnnltk glctinswhi fgkdnairig ensdvlvtre  
121pyvscdpdec rfyalsqgtt irgkhsngti hdsrqyralv swplsspptv yntrvecigw  
181sstschdgka rmsicvsgpn nnasaviwyk grpiteitw arnilrtqes ecvchngicp  
241vvftdgsatg paetriyyfk egkilkwepl tgtakhieec scygdseite ctcrdnwqgs  
301nrpviginp mmthtsqyic spvltndnprp ndptvgkcnd pypgnnnngv kgfsyldgdn  
361twlgrtista srsqyemlkv pnaltdrsk ptqggtivln tdwsgysgsf idywakgecy  
421racfyvelir grpkedkvww tsnsivsmcs steflgqwnw pdgakieyfl

ca A/Teal

[0325] Nucleotide sequence of ca A/Teal H6 (SEQ ID NO:  
35)

[0326] Entire molecule length: 1747 bp

1 agcaaaagca ggggaaaatg attgcaatca ttgtaatagc aatactggca gcagccggaa  
61 aatcagacaa gatctgcatt gggatatcatg ccaacaattc aacaacacag gtagatacga  
121 tacttgagaa gaattgtgact gtcacacact caattgaatt gctggaaaat cagaaggaag  
181 aaagattctg caagatattg aacaaggccc ctctcgactt aagggaatgt accatagagg  
241 gttggatctt ggggaatccc caatgcgacc tattgcttgg tgatcaaagc tggtcataca  
301 ttgtggaaag acctactgct caaaacggga tctgctaccc aggaacctta aatgaggtag  
361 aagaactgag ggcacttatt ggatcaggag aaagggtaga gagatttgag atgtttcccc  
421 aaagcacctg gcaaggagtt gacaccaaca gtggaacaac aagatcctgc ccttattcta  
481 ctggtgatcc gtctttctac agaaacctcc tatggataat aaaaaccaag acagcagaat  
541 atccagtaat taagggaatt tacaacaaca ctggaacca gccaatctc tatttctggg  
601 gtgtgcatca tcctcctaac accgacgagc aagatactct gtatggctct ggtgatcgat  
661 acgttagaat gggaaactgaa agcatgaatt ttgccaagag tccggaaatt gcggcaaggc  
721 ctgctgtgaa tggacaaaga ggcagaattg attattattg gtcggtttta aaaccagggg  
781 aaaccttgaa tgtggaatct aatggaaatc taatcgcccc ttggtatgca taaaatttg  
841 tcaacacaaa tagtaaagga gccgtcttca ggtcagattt accaatcgag aactgcatg  
901 ccacatgcc aactattgca ggggttctaa ggaccaataa aacatttcag aatgtgagtc  
961 ccctgtggat aggagaatgt cccaaatagc tgaagagtga aagtctgagg cttgcaactg  
1021 gactaagaaa tgttccacag attgaaacta gaggactctt cggagctatt gcagggttta

-continued

1081 ttgaaggagg atggactggg atgatatagtg ggtggtatgg ctatcaccat gaaaattctc  
 1141 aagggtcagg atatgcagcg gacagagaaa gcactcaaaa ggctgtaaac agaattacaa  
 1201 ataagggtcaa ttccatcatc aacaaaatga acacacaatt tgaagctgtc gatcacgaat  
 1261 tttcaaatct ggagaggaga atcgacaatc tgaacaaaag aatgcaagat ggatttctgg  
 1321 atgtttggac atacaatgct gaactgttgg ttcttcttga aaacgaaaga acactagaca  
 1381 tgcattgacgc aaatgtgaag aacctacatg aaaagggtcaa atcacaaacta agggacaatg  
 1441 ctaacgatct aggggaatgg tgccttgaat ttgggcataa gtgtgacaat gaatgcatag  
 1501 agtctgtcaa aaatggtaca tatgactatc ccaaatacca gactgaaagc aaattaaaca  
 1561 ggctaaaaat agaatcagta aagctagaga accttgggtg gtatcaaat cttagcattt  
 1621 atagtacggt atcgagcagc ctagtgttgg tagggctgat catggcaatg ggtctttgga  
 1681 tgtgttcaaa tgggtcaatg cagtgcaatg tgtgtatatg attaagaaaa acacccttgt  
 1741 ttctact

[0327] Amino acid sequence of ca A/Teal H6 (SEQ ID NO: 41)

[0328] Entire molecule length: 567 aa

1 miaiivaiail aaagksdkic igyhannstt qvdtileknv tvthsielle nqkeerfcki  
 61 lnkapldlre ctiegwilgn pqcdlllgdq swsyiverpt aqngicypgt lneveelral  
 121 lgsgerverf emfpqstwgq vdtngttrs cpystgdpsf yrnllwiikt ktaeypvikg  
 181 iynntgtqpi lyfwgvhphp ntdeqdtlyg sgdryvrmgt esmnfakspe iaarpavngq  
 241 rgridyywsv lkpgetlnve sngnliapwy aykfvntnsk gavfrsdlpi encdatcqi  
 301 agvlrtnktf qnvsplwige cpkyvksesl rlatglrnvp qietrglfga iagfieggwt  
 361 gmidgwygyh hensqgsqya adrestqkav nritnkvnsl inkmntqfea vdhefsnler  
 421 ridnlnkrmq dgfldvwtyn aellvllene rtdmhdanv knlhekvsq lrdnandlgn  
 481 gcfefwhkcd neciesvkng tydypkyqte sklnrlkies vklenlgvyq ilaiystvss  
 541 slvlvglima mglwmcngs mqcnvci

[0329] Nucleotide sequence of ca A/Teal N1 (SEQ ID NO: 36)

[0330] Entire molecule length: 1401 nt

1 agcaaaagca ggagttaaac atgaatccaa atcagaagat aataaccatt gggtaactct  
 61 gtatggtagt tggaataatc agcttgatgt tacaaattgg aaacataata tcaatatggg  
 121 ttagccatcat aattcagact gggcatccaa accagcctgg gccatgcaat caaagcatca  
 181 atttttacac tgagcaggct gcagcttcag tgacattagc gggtaattcc tctctctgcc  
 241 ctattagtgg atgggctata tacagtaaag acaatagtat aagaattggt tccaagggg  
 301 atgtgtttgt tatgagagaa ccattcggtt catgctocca ttggaatgc agaacccttt  
 361 tcttgactca aggagcccta ttgaatgaca agcattctaa tgggaccgtt aaagacagaa  
 421 gccctatag aactttaatg agctgtcctg ttggtgaggc tccttcccca tacaactcaa

-continued

481 ggtttgagtc tgttgcttgg tcagcaagtg cttgccatga tggcattagt tggctaacaa  
 541 ttggaatttc cggtcggat aatggggctg tggtgtgtt gaaatacaat ggcataataa  
 601 cagacaccat caagagtgg aggaacaaca tactgaggac acaagagtct gaatgtgcat  
 661 gtgtgaatgg ttcttgtttt actgtaatga cagatggacc gagtaatgaa caggccctcat  
 721 acaagatttt caagatagag aaggggaaag tagtcaaatac agttgagttg aacgccccta  
 781 attatcatta cgaggaatgc tctgtttatc ctgatgctgg cgaaatcaca tgtgtgtgca  
 841 gggataattg gcattggctg aaccgaccgt ggggtgtctt caatcagaat ctggagtatc  
 901 aaataggata tatatgcagt ggggttttctg gagacagtcc acgcccgaat gatggaacag  
 961 gcagttgcgg tccagtgtct cttaacggag agtatggagt aaaagggttt tcatttaagt  
 1021 acggtgatgg tgtttggatc gggagaacca aaagcactag ttccaggagc gggtttgaaa  
 1081 tgatttggga tccaaatggg tggaccgaaa cagatagtaa cttctcattg aagcaagaca  
 1141 tcatagcaat aactgattgg tcaggataga gggggagttt tgtccaacat ccagaactga  
 1201 caggattaaa ttgcatgagg ccttgcctct gggttgaact aatcagaggg aggcccgaag  
 1261 agaaaacaat ctggactagt gggagcagta tatctttctg tgggtgaaat agtgacactg  
 1321 tgggttggtc ttggccagac ggtgctgagg tgccattcac cattgacaag tagtttggtc  
 1381 aaaaaactcc ttgtttctac t

[0331] Amino acid sequence of ca A/Teal N1 (SEQ ID NO: 42)

[0332] Entire molecule length: 450 aa

1mnpnqkiiti gsicmvvgii slmlqignii siwvshiiqt ghpnqpgpcn qsinfyeteqa  
 61aasvtlagns slcpisgwai yskdnsirig skgdvfvrmre pfvscshlec rtffltgga  
 121lndkhsngtv kdrspyrthm scpvgeapsp ynsrfesvaw sasachdgis wltigisgpd  
 181ngavavlkyn giitdtiksw rnnilrtqes ecacvngscf tvmtdgpsne qasykifkie  
 241kgkvksvel napnyhyec scypdageit cvcrdnwhgs nrpwvsfnqn leyqigyics  
 301gvfgdsprpn dgtgscgpvs lngeygvkgf sfkygdgvi grtkstssrs gfemiwdpng  
 361wtetdsnfs lkdiaitdw sgysgsfvqh peltglnmr pcfwvelirg rpkektiws  
 421gssisfcgyn sdtvgwswpd gaevpftidk

ca A/Mallard

[0333] Nucleotide sequence of ca A/Mallard H6 (SEQ ID NO: 37)

[0334] Entire molecule length: 1745 bp

1 agcaaaagca ggggaaaatg attgcaatca taatacttgc aatagtggtc tctaccagca  
 61 agtcagacag gatctgcatt ggttaccatg caaacaactc gacaacacaa gtggacacaa  
 121 tattagagaa gaatgtgaca gtgacacact cagtggagct cctagaaaac cagaaggaga  
 181 atagattctg cagagtcttg aataaagcgc cactggatct aatggactgc accactgagg  
 241 gttggatcct tggaaacccc cgatgtgata acttactcgg tgatcaaagt tggtcataca

-continued

301 tagtagagag gacctgatgcc caaaatggga tatgttacct aggggtattg aaggagacgg  
 361 aagagctgaa agcactcatt gggctctatag atagcataca aagatttgaa atgtttccca  
 421 agagcacgtg gaccggggta gatactaata gcggagttag gagcgcttgc ccctacaatg  
 481 gtgaatcttc cttttacagg aatctgttgt ggataataaa aataagatct gatccgtact  
 541 cattgatcaa ggggacatat accaatacag gctctcagcc aatcttatat ttctggggtg  
 601 tgcaccatcc tccagatgaa gttgagcaag ctaacttgta tgggaattggt acccggtatg  
 661 ttaggatggg aactgaaagt atgaattttg ccaaagggtcc tgaaatagca ggcagaccac  
 721 ctgcgaatgg gcaacgagga agaattgatt attattggtc tgtgttgaag ccaggagaaa  
 781 ccttgaatgt ggaatccaat ggaaatttaa tagctccttg gtatgcttac aagttcacta  
 841 gttccagaaa caaggagct attttcaaat cagaccttcc aattgagaat tgtgatgctg  
 901 tctgtcaaac tttagctgga gcaataaata caaacaaaac ctcccaaat attagtccag  
 961 tctggattgg agaatgcccc aaatatgtta aaagtaagag cctaaaacta gcaactggtc  
 1021 tgagaaatgt tccacaggca gaaacaagag gattgtttgg agcaatagct gggtttatag  
 1081 aaggaggatg gacaggtagt gtagacggat ggtacggata ccaccatgaa aattcacagg  
 1141 ggtctggtta tgcagcagat aaagaaagca ctacagaaagc aatagacggg atcaccaata  
 1201 aagtcaatc aatcattgac aaaatgaaca cacaatttga ggcagtagag catgagtctt  
 1261 caagtctcga aaggagaata ggcaatctga acaaaagaat ggaagatgga ttttttagacg  
 1321 tgtggacata caatgctgaa cttctggttc tactggaaaa tgagaggact ttggacatgc  
 1381 atgatgctaa tgtaaagaat ctacatgaaa aggtgaaatc acaattaagg gataatgcaa  
 1441 aggatttggg taatgggtgt tttgaatttt ggcacaaatg cgacaatgaa tgcacaaact  
 1501 cagttaaaaa tggcacatat gactacccaa agtaccagga agagagcaga cttaataggc  
 1561 aggaaataaa atcagttagt ctggaaaatc tgggagtata ccaaatcctt gctatttata  
 1621 gtacggatc gagcagctctg gttttggtgg gactgatcat tgccatgggt ctttggatgt  
 1681 gctcaaatgg ctcaatgcaa tgcaagatat gtatataatt agaaaaaac acccttggtt  
 1741 ctact

[0335] Amino acid sequence of ca A/Mallard H6 (SEQ ID NO: 43)

[0336] Entire molecule length: 566 aa

1miaiilaiv vstksdric igyhannstt qvdtileknv tvthsvelle nqkenrfcrv  
 61lnkapldlmd cttegwilgn prcdnllgdq swsyiverpd aqngicypgv lketeelkal  
 121lgsidsiqrf emfpkstwtg vdtngvtsa cpyngessfy rnllwiikir sdpslikgt  
 181yntnsgspil yfwgvhhppd eveqanlygi gtryvmgte smnfakgpei agrppangqr  
 241gridyywsvl kpgetlnves ngliapwya ykftssrnkg aifksdlpie ncdavcqtla  
 301gaintnktfq nispvwigec pkyvkskslk latglrnvpq aetrglfgai agfieggwtg  
 361mvdgwygyhh ensqsgyaa dkestqkaid gitnkvnsl dkmntqfeav ehefsslerr  
 421ignlnkrmed gfldvwtyna ellvllener tldmhdanvk nlhekvsqsl rdnakdlng

-continued

481cfefwhkcdn ecinsvkngt ydyppyqees rlnrqiexv mlenlgvyqi laiystvsss  
541lvlgliiam glwmcengsm qckici

[0337] Nucleotide sequence of ca A/Mallard N2 (SEQ ID  
NO: 38)

[0338] Entire molecule length: 1467 nt

1 agcaaaagca ggagtgaata tgaatccaaa tcagaggata ataacaattg gatccgtctc  
61 tctaactatt gcaacagtgt gtttcctcat gcagattgcc atcctagcaa cgactgtgac  
121 actgcatttc aaacaaaatg aatgcagcat tcccgcaaac aaccaagtaa cgccatgtga  
181 accaatagta atagagagga acataacaga gatagtgtat ttgaataata ctaccataga  
241 aaaagagatt tgtcctgaag tagtagaata caggaattgg tcaaaaccgc aatgtcaaat  
301 tacagggttt gctcctttct ccaaggacaa ctcaattcgg ctttctgctg gtggggacat  
361 ttggataaca agagaacctt atgtgtcatg cgaccccggt aatgtttatc aatttgactc  
421 cgggcagggg accacgctgg acaacaaaca ctcaaatggc acaatacatg atagaatccc  
481 tcatcgagac cttttgatga atgaattggg tgttcggtt catttgggaa ccaacaagt  
541 gtgcatagca tggtcagct caagctgtca tgatgggaaa gcatggttgc acgtttgtgt  
601 cactggggat gatagaaatg caactgctag tttcatttat gatgggatgc ttattgacag  
661 tattggttcc tgggtcctaa atatcctcag gactcaggag tcagaatgcg tttgtatcag  
721 tggaaactgt acagtagtaa tgactgatgg aagtgcata ggaaggcgag acactagaat  
781 actattcatt agagagggga aaattgtcca cattagtcca ttgtcaggaa gtgctcagca  
841 tgtagaggaa tgttctgtt atccccgta ccaaacgctc agatgtgtct gcagagacaa  
901 ctggaagggc tctaataaggc cgttataga tataaatatg gcagattata gcattgactc  
961 aagttatgtg tgctcaggac ttgttgaga cacaccaagg aacgatgata gctctagcag  
1021 cagcaactgc agggatccta ataagagag agggaacca ggagtgaag ggtgggcctt  
1081 tgataatgga aatgatgtgt ggatgggaag aacaatcagt aaagattcgc gctcaggcta  
1141 tgagaccttc aaggtcattg gtggttgggc cattgcta atccaagtcac agaccaatag  
1201 acaagtcata gttgataata acaactggc tggttattct ggtattttct ctgttgaaag  
1261 caaaggctgc atcaataggt gtttttatgt ggagttgata agaggaaggc cacaggagac  
1321 tagagtatgg tggacctcaa acagtattgt cgtattttgt ggcacttcag ggacatatgg  
1381 aacaggctca tggcctgatg gggcgaatat cgatttcatt cctatataag ctttcgcaat  
1441 tttagaaaaa aactccttgt ttctact

[0339] Amino acid sequence of ca A/Mallard N2 (SEQ ID  
NO: 44)

[0340] Entire molecule length: 469 aa

1mnpnqriiti gsvltiatv cfmqiaila ttvtlhfkn ecsipannqv tpcepivier  
6lniteivylm ttiekeicpe vveyrnwskp qcqitgfapf skdnsirlsa ggdiwitrep  
121yvscdpskcy qfalgggttl dnkhsngtih driphrtllm nelgvpfhlq tkqvciawss

-continued

181sschdskawl hvctvgddrn atasfydgm lidsigswsq nilrtqesec vcisgtctvv  
 241mtdgsasgra dtrilfireg kivhisplsg saqhveecsc yprypnvrcv crdnwkgsnr  
 301pvidinmady sidssyvcsg lvgdtprndd sssssncrdp nnergngpvk gwafdnngdv  
 361wmgrtiskds rsgyetfkvi ggwaiaansks qtnrqvivdn nnwsgysgif sveskgcinc  
 421cfyvelirgr pqetrvwts nsivvfegts gtygtgswpd ganidfmpi

## [0341] Summary of SEQ ID NO Designations

SEQ ID NO	HA or NA	STRAIN NAME	Amino Acid or Nucleotide
SEQ ID NO: 1	HA (H5)	ca A/Vietnam/1203/04	Nucleotide
SEQ ID NO: 2	NA (N1)	ca A/Vietnam/1203/04	Nucleotide
SEQ ID NO: 3	HA (H5)	ca A/Hong Kong/213/03	Nucleotide
SEQ ID NO: 4	NA (N1)	ca A/Hong Kong/213/03	Nucleotide
SEQ ID NO: 5	HA (H5)	ca A/Hong Kong/491/97	Nucleotide
SEQ ID NO: 6	NA (N1)	ca A/Hong Kong/486/97	Nucleotide
SEQ ID NO: 7	HA (H5)	ca A/Hong Kong/491/97 (Ser211)	Nucleotide
SEQ ID NO: 8	NA (N1)	ca A/Hong Kong/486/97	Nucleotide
SEQ ID NO: 9	HA (H9)	ca A/ck/Hong Kong/G9/97	Nucleotide
SEQ ID NO: 10	NA (N2)	ca A/ck/Hong Kong/G9/97	Nucleotide
SEQ ID NO: 21	HA (H7)	A/Netherlands/219/03	Nucleotide
SEQ ID NOS: 22 & 45	NA (N7)	A/Netherlands/219/03	Nucleotide
SEQ ID NO: 23	HA (H7)	A/ck/BC/CN-7/04	Nucleotide
SEQ ID NO: 24	NA (N3)	A/ck/BC/CN-7/04	Nucleotide
SEQ ID NO: 25	HA (H7)	ca A/ck/BC/CN-6/04	Nucleotide
SEQ ID NO: 26	NA (N3)	ca A/ck/BC/CN-6/04	Nucleotide
SEQ ID NO: 33	HA (H6)	ca A/Duck	Nucleotide
SEQ ID NO: 34	NA (N9)	ca A/Duck	Nucleotide
SEQ ID NO: 35	HA (H6)	ca A/Teal	Nucleotide
SEQ ID NO: 36	NA (N1)	ca A/Teal	Nucleotide
SEQ ID NO: 37	HA (H6)	ca A/Mallard	Nucleotide
SEQ ID NO: 38	NA (N2)	ca A/Mallard	Nucleotide
SEQ ID NO: 11	HA (H5)	ca A/Vietnam/1203/04	Amino Acid
SEQ ID NO: 12	NA (N1)	ca A/Vietnam/1203/04	Amino Acid

-continued

SEQ ID NO	HA or NA	STRAIN NAME	Amino Acid or Nucleotide
SEQ ID NO: 13	HA (H5)	ca A/Hong Kong/213/03	Amino Acid
SEQ ID NO: 14	NA (N1)	ca A/Hong Kong/213/03	Amino Acid
SEQ ID NO: 15	HA (H5)	ca A/Hong Kong/491/97	Amino Acid
SEQ ID NO: 16	NA (N1)	ca A/Hong Kong/486/97	Amino Acid
SEQ ID NO: 17	HA (H5)	ca A/Hong Kong/491/97 (Ser211)	Amino Acid
SEQ ID NO: 18	NA (N1)	ca A/Hong Kong/486/97	Amino Acid
SEQ ID NO: 19	HA (H9)	ca A/ck/Hong Kong/G9/97	Amino Acid
SEQ ID NO: 20	NA (N2)	ca A/ck/Hong Kong/G9/97	Amino Acid
SEQ ID NO: 27	HA (H7)	A/Netherlands/219/03	Amino Acid
SEQ ID NO: 28	NA (N7)	A/Netherlands/219/03	Amino Acid
SEQ ID NO: 29	HA (H7)	A/ck/BC/CN-7/04	Amino Acid
SEQ ID NO: 30	NA (N3)	A/ck/BC/CN-7/04	Amino Acid
SEQ ID NO: 31	HA (H7)	ca A/ck/BC/CN-6/04	Amino Acid
SEQ ID NO: 32	NA (N3)	ca A/ck/BC/CN-6/04	Amino Acid
SEQ ID NO: 39	HA (H6)	ca A/Duck	Amino Acid
SEQ ID NO: 40	NA (N9)	ca A/Duck	Amino Acid
SEQ ID NO: 41	HA (H6)	ca A/Teal	Amino Acid
SEQ ID NO: 42	NA (N1)	ca A/Teal	Amino Acid
SEQ ID NO: 43	HA (H6)	ca A/Mallard	Amino Acid
SEQ ID NO: 44	NA (N2)	ca A/Mallard	Amino Acid

[0342]

## SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 45

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 1767

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 1

agcaaaagca ggggttcaat ctgtcaaaat ggagaaaata gtgcttcttt ttgcaatagt 60

cagtcttggtt aaaagtgatc agatttgcac tggttaccat gcaaacact cgacagagca 120

ggttgacaca ataattgaaa agaactgtac tgttacacat gcccaagaca tactggaaaa 180

-continued

gaaacacaac gggaagctct gcgatctaga tggagtgaag cctctaattt tgagagattg	240
tagcgtagct ggatggctcc tcggaacccc aatgtgtgac gaattcatca atgtgccgga	300
atggctttac atagtggaga aggccaatcc agtcaatgac ctctgttacc caggggattt	360
caatgactat gaagaattga aacacctatt gagcagaata aaccattttg agaaaattca	420
gatcatcccc aaaagttctt ggtccagtca tgaagcctca ttaggggtga gctcagcatg	480
tcataaccag ggaaagtctt cctttttcag aaatgtggta tggcttatca aaaagaacag	540
tacataccca acaataaaga ggagctacaa taataccaac caagaagatc ttttggtagt	600
gtgggggatt caccatccta atgatgcggc agagcagaca aagctctatc aaaacccaac	660
cacctatatt tccgttggga catcaacact aaaccagaga ttggtaccaa gaatagctac	720
tagatccaaa gtaaacgggc aaagtgaag gatggagttc ttctggacaa ttttaagcc	780
gaatgatgca atcaacttcg agagtaatgg aaatttcatt gctccagaat atgcatacaa	840
aattgtcaag aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa	900
caccaagtgt caaactccaa tggggggcat aaactctagc atgccattcc acaatatata	960
ccctctcacc attggggaat gcccacaata tgtgaaatca aacagattag tccttgcgac	1020
tgggctcaga aatagccctc aaagagagac tcgaggatta tttggagcta tagcaggttt	1080
tatagagga ggatggcagg gaatggtaga tggttggtat ggggtaccacc atagcaatga	1140
gcaggggagt gggtagctg cagacaaaga atccactcaa aaggcaatag atggagtcac	1200
caataaggtc aactcgatca ttgacaaaat gaacactcag tttgaggccg ttggaaggga	1260
atttaacaac ttagaaagga gaatagagaa tttaacaag aagatggaag acgggttcct	1320
agatgtctgg acttataatg ctgaacttct ggttctcatg gaaaatgaga gaactctaga	1380
ctttcatgac tcaaatgtca agaacttta cgacaaggtc cgactacagc ttagggataa	1440
tgcaaggag ctgggtaacg gttgtttcga gttctatcat aaatgtgata atgaatgtat	1500
ggaaagtgt agaaatggaa cgtatgacta cccgcagtat tcagaagaag cgagactaaa	1560
aagagaggaa ataagtggag taaaattgga atcaatagga atttaccaaa tactgtcaat	1620
ttattctaca gtggcgagt ccctagcact ggcaatcatg gtactgtgac tacccttatg	1680
gatgtgctcc aatgggtcgt tacaatgcag aatttgcatt taaatttgat agttcagatt	1740
gtagttaaaa acacccttgt ttctact	1767

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 1398

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 2

agcaaaagca ggagtccaat atgaatccaa atcagaagat aataaccatc ggggtcaatct	60
gtatggtaac tggaaatggt agcttaatgt tacaatttgg gaacatgac tcaatatggg	120
tcagtcattc aattcacaca gggaatcaac accaatctga accaatcagc aataactaatt	180
ttcttactga gaaagctgtg gcttcagtaa aattagcggg caattcatct ctttgcccca	240
ttaacggatg ggctgtatag agtaaggaca acagtataag gatcggttcc aagggggatg	300
tgtttgttat aagagagccg ttcattcatg gctccactt ggaatgcaga actttctttt	360
tgactcaggg agccttgctg aatgacaagc actccaatgg gactgtcaaa gacagaagcc	420



-continued

---

ctcacagaac attaatgagt tgtcctgtgg gtgaggctcc ctcccatat aactcaaggt	480
ttgagtctgt tgcttggtca gcaagtgtt gccatgatgg caccagtgg ttgacgattg	540
gaatttctgg ccagacaat ggggctgtgg ctgtattgaa atacaatggc ataataacag	600
acactatcaa gaggtaggag aacaacatac tgagaactca agagtctgaa tgtgcatgtg	660
taaatggctc ttgctttact gtaatgactg acggaccaag taatggtcag gcatcacata	720
agatcttcaa aatggaaaaa gggaaagtgg ttaaatcagt cgaattggat gtcctaatt	780
atcactatga ggaatgctcc tgttatccta atgccggaga aatcacatgt gtgtgcaggg	840
ataattggca tggctcaaat cggccatggg tatctttcaa tcaaaatttg gagtatcaaa	900
taggatatat atgcagtgga gttttcggag acaatccacg cccaatgat ggaacaggta	960
gttgtgttcc ggtgtcctct aacggggcat atggggtaaa agggttttca tttaaatacg	1020
gcaatgggtg ctggatcggg agaaccaaaa gcaactaattc caggagcggc tttgaaatga	1080
tttgggatcc aaatgggtgg actgaaacgg acagtagctt ttcagtgaag caagatatcg	1140
tagcaataac tgattggtca ggatatagcg ggagttttgt ccagcatcca gaactgacag	1200
gactagattg cataagacct tgtttctggg ttgagttgat cagagggcgg cccaaagaga	1260
gcacaatttg gactagtggg agcagcatat ctttttgtgg tgtaaatagt gacactgtgg	1320
gttggctctg gccagacggt gctgagttgc cattcaccat tgacaagtag tttgttcaaa	1380
aaactccttg tttctact	1398

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 1767

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 3

agcaaaagca ggggttcaat ctgtcaaaat ggagaaaata gtgcttcttt ttgcaatagt	60
cagtcttgtt aaaagtgate agatttgcat tggttaccat gcaacaact cgacagagca	120
ggttgacaca ataatggaaa agaacgttac tggtacacat gcccaagaca tactggaaaa	180
gacacacaac gggaagctct gcgactaga tggagtgaag cctctaattt tgagagattg	240
tagtgtagct ggatggctcc tcggaaaccc aatgtgtgac gaattcatca atgtgccgga	300
atggtcttac atagtggaga aggccaatcc agccaatgac ctctgttacc caggggattt	360
caacgactat gaagaattga aacacctatt gagcagaata aaccattttg agaaaattca	420
gatcatcccc aaaaattctt ggtccagtca tgaagcctca ttaggggtga gctcagcatg	480
tcataccaa ggaaagtctt cttttttcag gaatgtggtg ttgcttatca aaaagaacaa	540
tgcataccca acaataaaga ggagctacaa taataccaac caagaagatc ttttggattt	600
gtgggggatt caccatccta atgatgcggc agagcagact aggtctctatc aaaacccaac	660
cacctacatt tccgttggga catcaacact aaaccagaga ttggtaccaa aaatagctac	720
tagatccaaa gtaaacgggc aaaatggaag gatggagttc ttctggacaa ttttaaaacc	780
gaatgatgca atcaacttcg agagcaatgg aaatttcatt gctccagaat atgcatacaa	840
aattgtcaag aaaggggact cagcaattat gaaaagtga ttggaatatg gtaactgcaa	900
caccaagtgt caaactccaa tggggggcat aaactctagt atgccattcc acaatatata	960
ccctctcacc atcgggggaat gcccacaaata tgtgaaatca aacagattag tccttgcgac	1020

-continued

tgggctcaga aatagccctc aaagagagac tggaggatta tttggagcta tagcaggttt	1080
tatagagggg ggatggcagg gaatggtaga tgggttggtat ggggtaccacc atagcaatga	1140
gcaggggagt gggtagcgtg cagacaaaga atccactcaa aaggcaatag atggagtcac	1200
caataaggtc aactcgatca ttgacaaaat gaacactcag tttgaggccg ttggaaggga	1260
atttaataac ttagaaagga gaatatagaa tttaacaag aagatggaag acggattcct	1320
agatgtctgg acttataatg ctgaacttct ggttctcatg gaaaatgaga gaactctaga	1380
ctttcatgac tcaaagtca agaaccttta cgacaaggtc cgactacagc ttagggataa	1440
tgcaaaggag ctgggtaacg gttgtttcga gttctatcac aaatgtgata atgaatgtat	1500
ggaaagtgtg agaaacggaa cgtatgacta cccgcagtat tcagaagaag caagactaaa	1560
aagagaggaa ataagtggag taaaattgga gtcaatagga acttaccaaa tactgtcaat	1620
ttattctaca gtggcgagtt ccctagcact ggcaatcatg gtagctggtc tatctttatg	1680
gatgtgctcc aatgggtcgt tacaatgcag aatttgcatt taaatttggtg agttcagatt	1740
gtagttaaaa acacccttgt ttctact	1767

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 1458

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 4

agcaaaaagca ggagtcaaaa atgaatccaa atcagaagat aacaaccatt ggatcaatct	60
gtatggtaat tggaatgatt agcttgatgt tacaattgg gaacataatc tcaatatggg	120
ttagtcattc aattcaaaca gggaatcaac accaggctga accatgcaat caaagcatta	180
ttacttatga aaacaacacc tgggtaaacc agacatatgt caacatcagc aataccaatt	240
ttcttactga gaaagctgtg gcttcagtaa cattagcggg caattcatct ctttgcccca	300
ttagtggatg ggctgtatac agtaaggaca acggtataag aatcggttcc aagggggatg	360
tgtttgttat aagagagcgg ttcattcatg gctccactt ggaatgcaga actttctttt	420
tgactcaggg agccttctgt aatgacaagc attctaattg gaccgtcaaa gacagaagcc	480
ctcacagaac attaatgagt tgtcccgtag gtgaggctcc tccccatac aactcgaggt	540
ttgagtctgt tgcttggtcg gcaagtgtt gtcattgatg cactagttag ttgacaattg	600
gaatttcttg cccagacaat ggggctgtgg ctgtattgaa atacaatggc ataataacag	660
acactatcaa gagttggagg aacaacataa tgagaactca agagtctgaa tgtgcatgtg	720
taaatggctc ttgctttact gttatgactg atggaccaag taatgggcag gcttcataca	780
aaatcttcag aatagaaaaa gggaaagtag ttaaatcagc cgaattaaat gccctaatt	840
atcactatga ggagtgtccc tgttatcctg atgctggaga aatcacatgt gtgtgcaggg	900
ataactggca tggctcaaat cggccatggg tatctttcaa tcaaaatttg gagtatcgaa	960
taggatatat atgcagtgga gttttcggag acaatccacg cccaatgat gggacaggca	1020
gttggtggtcc ggtgtccct aaaggggcat atggaataaa agggttctca tttaaatacg	1080
gcaatgggtg ttggatcggg agaaccataa gcaactaatc caggagcggc tttgaaatga	1140
tttgggatcc aaatggatgg actggtacgg acagtaattt ttcagtaaag caagatattg	1200
tagctataac cgattgtgca ggatatagcg ggagttttgt ccagcatcca gaactgacag	1260

-continued

gattagattg cataagacct tgtttctggg ttgagctaata cagagggcgg cccaaagaga	1320
gcacaatttg gactagtggg agcagcatat ccttttgttg tgtaaatagt gacactgttg	1380
gttggtcttg gccagacggt gctgagttgc cattcaccat tgacaagtag tttgttcaaa	1440
aaactccttg tttctact	1458

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 1767

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 5

agcaaaagca ggggtataat ctgtcaaaat ggagaaaata gtgcttcttc ttgcaacagt	60
cagccttggtt aaaagtgacc agatttgcatt tggttaccat gcaacaact cgacagagca	120
agttgacaca ataattgaaa agaattgtac tgttacacat gccaagaca tactggaaag	180
gacacacaac gggaagctct gcgatctaaa tggagtgaag cctctgattt tgagggattg	240
tagtgtagct ggatggctcc tcggaaaccc tatgtgtgac gaattcatca atgtgccgga	300
atggtcttac atagtggaga aggccagtc agccaatgac ctctgttatc cagggaattt	360
caacgactat gaagaactga aacacctatt gagcagaata aaccattttg agaaaattca	420
gataatcccc aaaagttctt ggtccaatca tgatgcctca tcaggggtga gctcagcatg	480
tccatacctt gggaggtcct cctttttcag aaatgtggta tggcttatca aaaagaacag	540
tagctaccca acaataaaga ggagctacaa taataccaac caagaagatc ttttggtagt	600
gtgggggatt caccatccta atgatgcggc agagcagaca aggtctctac aaaacccaac	660
cacctacatt tccgttgga catcaacact gaaccagaga ttggttccag aaatagctac	720
tagaccctaa gtaaacgggc aaagtggaag aatggagttc ttctggacaa ttttaagcc	780
gaatgatgcc atcaatttcg agagttaagg aaatttcatt gctccagaat atgcatacaa	840
aattgtcaag aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa	900
caccaagtgt caaactccaa tgggggcaat aaactctagt atgccattcc acaacataca	960
ccccctcacc atcggggaat gccccaaata tgtgaaatca aacagattag tccttgcaac	1020
tggactcaga aatacccctc aacgagagac gcgaggacta tttggagcta tagcaggttt	1080
tatagagggg gatggcagg gaattgtaga tgggttggtat gggtagccac atagcaatga	1140
gcaggggagt ggatacgtg cagaccaaga atccacacaa aggcgaatag atggagtcac	1200
caataaggtc aactcgatca ttaacaaaat gaacactcag tttgaggccg ttggaaggga	1260
atttaataac ttggaaggga ggatagagaa ttttaacaag aaaatggaag acggattcct	1320
agatgtctgg acttacaatg ccgaacttct ggttctcatg gaaaatgaga gaactctaga	1380
ctttcatgac tcaaatgtca agaacttta cgacaaggtc cgactacagc ttagggataa	1440
tgcaaaggag ctgggtaatg gttgtttcga attctatcac aaatgtgata acgaatgtat	1500
ggaaagtgtg aaaaacggaa cgtatgacta cccgcagtat tcagaagaag caagactaaa	1560
cagagaggaa ataagtggag taaaattgga atcaatggga acttaccaaa tactgtcaat	1620
ttattcaaca gtggcgagtt ccctagcact ggcaatcatg gtagctggtc tatctttatg	1680
gatgtgctcc aatggatcgt tacaatgcag aatttgcatt taaattgtg agttcagatt	1740
gtagttaaaa acacccttgt ttctact	1767

-continued

---

<210> SEQ ID NO 6  
 <211> LENGTH: 1401  
 <212> TYPE: DNA  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 6

```

agcaaaagca ggagtttaaa atgaatccaa atcagaagat aataaccatt ggatcaatct    60
gcatggtagt tgggataatc agcttgatgt taaaaattgg aaacacaata tcagtatggg    120
tcagccacat aattaaaact tggcacccaa accagcctga accatgcaac caaagcatca    180
atttttacac tgagcaggct gcagcttcag tgacattagc gggcaattcc tctctctgcc    240
ctattagtgg atgggcctata tacagcaagg acaatagtat aagaattggg tccaaagggg    300
atgtgtttgt tataagagaa ccattcatct catgctccca tttggaatgc agaacccttt    360
tcttgacca aggagcccta ttgaatgaca agcattctaa tgggaccgtc aaagacagga    420
gcccctatag aactttaatg agctgtcctg ttggtgaggc cccttcccca tacaactcaa    480
ggtttgagtc tgttgcttgg tcagcaagtg cttgccatga tggcattagt tggctaacaa    540
ttggaatttc cggtcggat aatggggctg tggctgtgtt gaaatacaat ggcataataa    600
cagacaccat caagagtgg aggaacaaca cactgaggac gcaagagtct gaatgtgcat    660
gtgtgaatgg ttctgtttt actgtaatga cagatggacc gagtaatgaa caggcctcat    720
acaagatttt caagatagaa aaggggaggg tagtcaaac agttgagttg aacgccccta    780
attatcatta cgaggaatgc tcctgttacc ctgatgctgg cgaaatcaca tgtgtgtgca    840
gggataattg gcatggctcg aaccgaccat ggggtgtctt caatcagaat ctggagtatc    900
aaataggata tatatgcagt ggggttttcg gagacagtcc acgcccctaat gatgggacag    960
gcagttgtgg tccagtgtct cttaacggag cgtatggagt aaaaggggtt tcattttaat    1020
acggcaatgg tgtttggatc gggagaacca aaagcactag ttccaggagc ggttttgaaa    1080
tgatttggga tccaaatggg tggaccgaaa cagacagtag cttctcgttg aagcaagaca    1140
tcatagcgat aactgattgg tcaggataca gcgggagttt tattcaacat ccagaactga    1200
caggattaaa ttgcatgaga ccttgcttct ggggtgaact aatcagaggg aggcccaaag    1260
agaaaacaat ctggactagt gggagcagta tatctttctg tgggtgaaat agtgacactg    1320
tgggttggtc ttggccagac ggtgctgagt tgccatacac cattgacaag tagtttggtc    1380
aaaaaactcc ttgtttctac t                                     1401
  
```

<210> SEQ ID NO 7  
 <211> LENGTH: 1767  
 <212> TYPE: DNA  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 7

```

agcaaaagca ggggtataat ctgtcaaaat ggagaaaata gtgcttcttc ttgcaacagt    60
cagccttgtt aaaagtgacc agatttgcac tgggtaccat gcaacaact cgacagagca    120
agttgacaca ataatggaaa agaattgtac tggtacacat gccaagaca tactggaaag    180
gacacacaac gggaagctct cggatctaaa tggagtgaag cctctgattt tgagggattg    240
tagtgtagct ggtggtctcc tcggaaaccc tatgtgtgac gaattcatca atgtgccgga    300
atggtcttac atagtggaga aggccagtc agccaatgac ctctgttacc caggggaattt    360
  
```

-continued

caacgactat gaagaactga aacacctatt gagcagaata aaccattttg agaaaattca	420
gataatcccc aaaagttctt ggtccaatca tgatgcctca tcaggggtga gctcagcatg	480
tccatacctt gggaggtcct cctttttcag aaatgtggta tggcttatca aaaagaacag	540
tagctaccca acaataaaga ggagctacaa taataccaac caagaagatc ttttggtagt	600
gtgggggatt caccatccta atgatgcggc agagcagaca aggctctatc aaaacccaac	660
cacctacatt tccgttgga catcaacact gaaccagaga ttggtttcag aaatagctac	720
tagacccaaa gtaaacgggc aaagtggaag aatggagtgc tcttgacaa ttttaaagcc	780
gaatgatgcc atcaatttcg agagtaatgg aaatttcatt gctccagaat atgcatacaa	840
aattgtcaag aaaggggact caacaattat gaaaagtga ttggaatatg gtaactgcaa	900
caccaagtgt caaacccaa tgggggcaat aaactctagt atgccattcc acaacataca	960
ccccctacc atcggggaat gcccacaata tgtgaaatca aacagattag tccttgcaac	1020
tggactcaga aatacccctc aacgagagac gcgaggacta tttggagcta tagcaggttt	1080
tatagagga ggatggcagg gaatggtaga tggttggtat gggtagccac atagcaatga	1140
gcaggggagt ggatacgctg cagaccaaga atccacacaa aaggcaatag atggagtcac	1200
caataaggtc aactcgatca ttaacaaat gaacactcag tttgaggccg ttggaaggga	1260
atttaataac ttggaaggga ggatagagaa tttaaacaag aaaatggaag acggattcct	1320
agatgtctgg acttacaatg ccgaacttct ggttctcatg gaaaatgaga gaactctaga	1380
ctttcatgac tcaaatgtca agaacttta cgacaaggtc cgactacagc ttagggataa	1440
tgcaaggag ctgggtaatg gttgtttcga attctatcac aaatgtgata acgaatgtat	1500
ggaaagtgt aaaaacggaa cgtatgacta cccgcagtat tcagaagaag caagactaaa	1560
cagagaggaa ataagtggag taaaattgga atcaatggga acttaccaaa tactgtcaat	1620
ttattcaaca gtggcgagt ccctagcact ggcaatcatg gtactgtggtc tatctttatg	1680
gatgtgctcc aatggatcgt tacaatgcag aatttgcatt taaatttggt agttcagatt	1740
gtagttaaaa acacccttgt ttctact	1767

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 1401

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 8

agcaaaagca ggagtttaaa atgaatccaa atcagaagat aataaccatt ggatcaatct	60
gcatggtagt tgggataatc agcttgatgt tacaattgg aaacacaata tcagtatggg	120
tcagccacat aattaaaact tggcacccaa accagcctga accatgcaac caaagcatca	180
atttttacac tgagcaggct gcagcttcag tgacattagc gggcaattcc tctctctgcc	240
ctattagtgg atgggtata tacagcaagg acaatagtat aagaattggg tccaaagggg	300
atgtgtttgt tataagagaa ccattcatct catgctccca tttggaatgc agaactttt	360
tcttgacca aggagcccta ttgaatgaca agcattctaa tgggaccgtc aaagacagga	420
gcccctatag aactttaatg agctgtcctg ttggtgaggc ccttcccca tacaactcaa	480
ggtttgagtc tgttgcttgg tcagcaagtg cttgccatga tggcattagt tggctaacaa	540
ttggaatttc cggtcggat aatggggctg tggctgtgtt gaaatacaat ggcataataa	600

-continued

---

cagacacccat caagagttgg aggaacaaca cactgaggac gcaagagtct gaatgtgcat	660
gtgtgaatgg ttcttgtttt actgtaatga cagatggacc gagtaatgaa caggcctcat	720
acaagatttt caagatagaa aaggggaggg tagtcaaac agttgagttg aacgccccta	780
attatcatta cgaggaaatgc tcctgttatc ctgatgctgg cgaatacaca tgtgtgtgca	840
gggataattg gcatggctcg aaccgacccat ggggtgtctt caatcagaat ctggagtatc	900
aaataggata tatatgcagt ggggttttcg gagacagtcc acgcccgaat gatgggacag	960
gcagttgtgg tccagtgtct cttaacggag cgtatggagt aaaagggttt tcatttaaat	1020
acggcaatgg tgtttggatc gggagaacca aaagcactag tccaggagc ggttttgaaa	1080
tgatttggga tccaaatggg tggaccgaaa cagacagtag cttctcgttg aagcaagaca	1140
tcatagcgat aactgattgg tcaggataca gcgggagttt tattcaacat ccagaactga	1200
caggattaaa ttgcatgaga ccttgcttct ggggtgaact aatcagaggg aggcccaaag	1260
agaaaaaat ctggactagt gggagcagta tatctttctg tgggtgaaat agtgacactg	1320
tgggttggtc ttggccagac ggtgctgagt tgccatacac cattgacaag tagtttgttc	1380
aaaaaactcc ttgtttctac t	1401

&lt;210&gt; SEQ ID NO 9

&lt;211&gt; LENGTH: 1690

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 9

ttaaccactc aagatggaag caataccact aataactata ctactagtag taacagcaag	60
caatgcagac aaaatctgca tcggctacca atcaacaaac tccacagaaa ccgtagacac	120
gctaacagaa aacaatgttc ctgtgacaca tgccaaagaa ttgctccaca cagagcacia	180
tgggatgctg tgtgcaacaa atctgggacg tcctcttatt ctgacactt gcaccattga	240
aggactgac tatggcaacc cttcttgtga tctactgttg ggaggaagag aatggtecta	300
catcgctgaa agaccatcgg ctgttaatgg aatgtgttac cccgggaatg tagaaaacct	360
agaggaaact aggtcatttt ttagttctgc tagttcctac caaagaatcc agatctttcc	420
agacacaatc tggaatgtgt cttacagtgg aacaagcaaa gcatgttcag attcattcta	480
caggagcatg agatggttga ctcaaaagaa caacgcctac cctattcaag acgcccata	540
cacaaataat agaggaaaga gcattctttt catgtggggc ataaatcacc cacctaccga	600
tactgcacag acaaatctgt acacaaggac tgacacaaca acaagtgttg caacagaaga	660
tataaatagg accttcaaac cagtgatagg gccaaaggccc cttgtcaatg gtctgcaggg	720
aagaattgat tattattggt cggatttgaa accaggctcag acattgagag taagatccaa	780
tgggaatcta atcgctccat ggtatgggca cattctttca ggagagagcc acggaagaat	840
cctgaagact gatttaaaca gtggtagctg ttagtgcaa tgtcaaacag aaagaggtgg	900
cttaataact actttgccat tccacaatgt cagtaaatat gcatttgga actgcccata	960
atatgttga gtaaagagtc tcaaactggc agttgggtctg aggaatgtgc ctgctagatc	1020
aagtagagga ctatttgggg ccatagctgg attcatagag ggaggttggg cagggtctgg	1080
cgctggttgg tatgggttcc agcattcaaa tgatcaaggg gttggtatag ctgcagatag	1140
agactcaact caaagggcaa ttgacaaaat aacgtccaaa gtgaataata tagtcgataa	1200

-continued

---

aatgaacaag caatatgaaa ttattgatca tgaattcagc gaggttgaaa atagactcaa	1260
tatgatcaat aataagattg atgaccagat acaagacata tgggcatata acgctgaatt	1320
gctagtgctg cttgaaaacc agaaaacact cgatgagcat gatgcgaatg taaacaatct	1380
atataacaaa gtgaagaggg cactgggttc caatgcaatg gaagatggga aaggatgttt	1440
cgagctatac cataaatgtg atgatcagtg catggagaca attcggaacg ggacctataa	1500
caggaggaag tataaagagg aatcaagact agaaagacag aaaatagaag gggccaagct	1560
ggaatctgaa ggaacttaca aaatcctcac catttattcg actgtcgcct catctcttgt	1620
gattgcaatg gggtttgctg ccttcttgtt ctgggccatg tccaatggat cttgcagatg	1680
caacatttga	1690

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 1428

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 10

aaatgaatcc aaatcagaag ataatagcaa ttggctctgt ttctctaact attgcgacaa	60
tatgcctcct catgcagatt gctatcttag caacgactat gacactacat ttcaagcaga	120
atgaatgcat caactcctcg aataatcaag tagtgccatg tgaaccaatc ataatagaaa	180
ggaacataac agagatagtg catttgaata gtactacctt agagaaggaa atttgtccta	240
aagtagcaga ctacaggaat tgggtcaaac cacaatgtca aatcacaggg ttgcctcctt	300
tctccaagga caattcaatt aggcctctcg cagggtggaga tatttgggtg acaagagAAC	360
cttatgtatc gtgcggtcct ggtaaatgtt atcaatttgc acttgggcag ggaaccactt	420
tggagaacaa acactcaaac ggcacagcac atgatagaac tcctcataga acccttttaa	480
tgaatgagtt ggggtgttccg tttcatttgg caaccaaaca agtgtgcata gcatggacca	540
gctcaagctg ccatgatggg aaagcatggg tacatgtttg tgtcaactggg gatgatagaa	600
atgcaacggc tagcatcatt tatgatggga tacttggtga cagtattggg tcatgggtcta	660
aaaaatcctc cagaactcag gagtcagaat gcgtttgcat caatggaacc tgtgcagtag	720
taatgactga tggaagtgca tcaggaaggg ctgacactag aatactatct attagagagg	780
ggaaaattgc acacattagc ccattgtcag gaagtgtcga gcatgtggag gaatgtcct	840
gttaccctcc ataccagaa gttagatgtg tttgcagaga caattggaag ggatccaata	900
ggcccggtct atataaaat atggcaaatt atagtattga ttccagttat gtgtgctcag	960
gacttggttg cgacacacca agaaatgatg ataggtctag cagcagcaac tgcagagatc	1020
ctaataacga gagagggggc ccaggagtaa aagggtgggc ctttgacaat ggaaatgaca	1080
tttggtatgg aagaacaatc aaaaaggatt cgcgctcagg ttatgagact ttcagggtca	1140
ttggtgggtg gaccactgct aattccaagt cacagataaa tagacaagtc atagttgaca	1200
gtgataactc gtctgggtat tctggtatct tctctgttga aggcaaaagc tgcacaaaca	1260
gggtgtttta cgtggagttg ataagaggaa gaccaaagga gactaggggtg tgggtgactt	1320
caaatagcat cattgtatct tgtggaactt caggtaccta tggaacaggc tcatggcctg	1380
atggggcgaa tatcaatttc atgcctatat aagctttcgc aatttttag	1428

&lt;210&gt; SEQ ID NO 11

-continued

---

```

<211> LENGTH: 564
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 11
Met Glu Lys Ile Val Leu Leu Phe Ala Ile Val Ser Leu Val Lys Ser
 1             5             10             15
Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val
          20             25             30
Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile
          35             40             45
Leu Glu Lys Lys His Asn Gly Lys Leu Cys Asp Leu Asp Gly Val Lys
 50             55             60
Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn
 65             70             75             80
Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val
          85             90             95
Glu Lys Ala Asn Pro Val Asn Asp Leu Cys Tyr Pro Gly Asp Phe Asn
          100            105            110
Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu
          115            120            125
Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Ser Ser His Glu Ala Ser
          130            135            140
Leu Gly Val Ser Ser Ala Cys Pro Tyr Gln Gly Lys Ser Ser Phe Phe
          145            150            155            160
Arg Asn Val Val Trp Leu Ile Lys Lys Asn Ser Thr Tyr Pro Thr Ile
          165            170            175
Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp
          180            185            190
Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Lys Leu Tyr Gln
          195            200            205
Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg
          210            215            220
Leu Val Pro Arg Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Ser Gly
          225            230            235            240
Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn
          245            250            255
Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile
          260            265            270
Val Lys Lys Gly Asp Ser Thr Ile Met Lys Ser Glu Leu Glu Tyr Gly
          275            280            285
Asn Cys Asn Thr Lys Cys Gln Thr Pro Met Gly Ala Ile Asn Ser Ser
          290            295            300
Met Pro Phe His Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys
          305            310            315            320
Tyr Val Lys Ser Asn Arg Leu Val Leu Ala Thr Gly Leu Arg Asn Ser
          325            330            335
Pro Gln Arg Glu Thr Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe Ile
          340            345            350
Glu Gly Gly Trp Gln Gly Met Val Asp Gly Trp Tyr Gly Tyr His His
          355            360            365
Ser Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Lys Glu Ser Thr Gln

```



-continued

370	375	380
Lys Ala Ile Asp Gly Val Thr Asn Lys Val Asn Ser Ile Ile Asp Lys		
385	390	395 400
Met Asn Thr Gln Phe Glu Ala Val Gly Arg Glu Phe Asn Asn Leu Glu		
	405	410 415
Arg Arg Ile Glu Asn Leu Asn Lys Lys Met Glu Asp Gly Phe Leu Asp		
	420	425 430
Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Met Glu Asn Glu Arg		
	435	440 445
Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Asp Lys Val		
	450	455 460
Arg Leu Gln Leu Arg Asp Asn Ala Lys Glu Leu Gly Asn Gly Cys Phe		
	465	470 475 480
Glu Phe Tyr His Lys Cys Asp Asn Glu Cys Met Glu Ser Val Arg Asn		
	485	490 495
Gly Thr Tyr Asp Tyr Pro Gln Tyr Ser Glu Glu Ala Arg Leu Lys Arg		
	500	505 510
Glu Glu Ile Ser Gly Val Lys Leu Glu Ser Ile Gly Ile Tyr Gln Ile		
	515	520 525
Leu Ser Ile Tyr Ser Thr Val Ala Ser Ser Leu Ala Leu Ala Ile Met		
	530	535 540
Val Ala Gly Leu Ser Leu Trp Met Cys Ser Asn Gly Ser Leu Gln Cys		
	545	550 555 560
Arg Ile Cys Ile		

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 449

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 12

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Cys Met Val		
1	5	10 15
Thr Gly Ile Val Ser Leu Met Leu Gln Ile Gly Asn Met Ile Ser Ile		
	20	25 30
Trp Val Ser His Ser Ile His Thr Gly Asn Gln His Gln Ser Glu Pro		
	35	40 45
Ile Ser Asn Thr Asn Phe Leu Thr Glu Lys Ala Val Ala Ser Val Lys		
	50	55 60
Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Asn Gly Trp Ala Val Tyr		
	65	70 75 80
Ser Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly Asp Val Phe Val		
	85	90 95
Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu Cys Arg Thr Phe		
	100	105 110
Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly Thr		
	115	120 125
Val Lys Asp Arg Ser Pro His Arg Thr Leu Met Ser Cys Pro Val Gly		
	130	135 140
Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp Ser		
	145	150 155 160
Ala Ser Ala Cys His Asp Gly Thr Ser Trp Leu Thr Ile Gly Ile Ser		

-continued

165					170					175					
Gly	Pro	Asp	Asn	Gly	Ala	Val	Ala	Val	Leu	Lys	Tyr	Asn	Gly	Ile	Ile
			180					185					190		
Thr	Asp	Thr	Ile	Lys	Ser	Trp	Arg	Asn	Asn	Ile	Leu	Arg	Thr	Gln	Glu
			195				200					205			
Ser	Glu	Cys	Ala	Cys	Val	Asn	Gly	Ser	Cys	Phe	Thr	Val	Met	Thr	Asp
			210				215					220			
Gly	Pro	Ser	Asn	Gly	Gln	Ala	Ser	His	Lys	Ile	Phe	Lys	Met	Glu	Lys
					230					235					240
Gly	Lys	Val	Val	Lys	Ser	Val	Glu	Leu	Asp	Ala	Pro	Asn	Tyr	His	Tyr
					245				250					255	
Glu	Glu	Cys	Ser	Cys	Tyr	Pro	Asn	Ala	Gly	Glu	Ile	Thr	Cys	Val	Cys
			260					265					270		
Arg	Asp	Asn	Trp	His	Gly	Ser	Asn	Arg	Pro	Trp	Val	Ser	Phe	Asn	Gln
			275				280					285			
Asn	Leu	Glu	Tyr	Gln	Ile	Gly	Tyr	Ile	Cys	Ser	Gly	Val	Phe	Gly	Asp
			290				295					300			
Asn	Pro	Arg	Pro	Asn	Asp	Gly	Thr	Gly	Ser	Cys	Gly	Pro	Val	Ser	Ser
					310					315					320
Asn	Gly	Ala	Tyr	Gly	Val	Lys	Gly	Phe	Ser	Phe	Lys	Tyr	Gly	Asn	Gly
					325				330					335	
Val	Trp	Ile	Gly	Arg	Thr	Lys	Ser	Thr	Asn	Ser	Arg	Ser	Gly	Phe	Glu
			340					345					350		
Met	Ile	Trp	Asp	Pro	Asn	Gly	Trp	Thr	Glu	Thr	Asp	Ser	Ser	Phe	Ser
			355				360					365			
Val	Lys	Gln	Asp	Ile	Val	Ala	Ile	Thr	Asp	Trp	Ser	Gly	Tyr	Ser	Gly
			370				375					380			
Ser	Phe	Val	Gln	His	Pro	Glu	Leu	Thr	Gly	Leu	Asp	Cys	Ile	Arg	Pro
					390					395					400
Cys	Phe	Trp	Val	Glu	Leu	Ile	Arg	Gly	Arg	Pro	Lys	Glu	Ser	Thr	Ile
					405				410					415	
Trp	Thr	Ser	Gly	Ser	Ser	Ile	Ser	Phe	Cys	Gly	Val	Asn	Ser	Asp	Thr
			420					425					430		
Val	Gly	Trp	Ser	Trp	Pro	Asp	Gly	Ala	Glu	Leu	Pro	Phe	Thr	Ile	Asp
			435				440					445			

Lys

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 564

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 13

Met	Glu	Lys	Ile	Val	Leu	Leu	Phe	Ala	Ile	Val	Ser	Leu	Val	Lys	Ser
1				5					10					15	

Asp	Gln	Ile	Cys	Ile	Gly	Tyr	His	Ala	Asn	Asn	Ser	Thr	Glu	Gln	Val
			20					25					30		

Asp	Thr	Ile	Met	Glu	Lys	Asn	Val	Thr	Val	Thr	His	Ala	Gln	Asp	Ile
			35				40					45			

Leu	Glu	Lys	Thr	His	Asn	Gly	Lys	Leu	Cys	Asp	Leu	Asp	Gly	Val	Lys
			50			55				60					

Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn

-continued

65	70	75	80
Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val	85	90	95
Glu Lys Ala Asn Pro Ala Asn Asp Leu Cys Tyr Pro Gly Asp Phe Asn	100	105	110
Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu	115	120	125
Lys Ile Gln Ile Ile Pro Lys Asn Ser Trp Ser Ser His Glu Ala Ser	130	135	140
Leu Gly Val Ser Ser Ala Cys Pro Tyr Gln Gly Lys Ser Ser Phe Phe	145	150	155
Arg Asn Val Val Trp Leu Ile Lys Lys Asn Asn Ala Tyr Pro Thr Ile	165	170	175
Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp	180	185	190
Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Arg Leu Tyr Gln	195	200	205
Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg	210	215	220
Leu Val Pro Lys Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Asn Gly	225	230	235
Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn	245	250	255
Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile	260	265	270
Val Lys Lys Gly Asp Ser Ala Ile Met Lys Ser Glu Leu Glu Tyr Gly	275	280	285
Asn Cys Asn Thr Lys Cys Gln Thr Pro Met Gly Ala Ile Asn Ser Ser	290	295	300
Met Pro Phe His Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys	305	310	315
Tyr Val Lys Ser Asn Arg Leu Val Leu Ala Thr Gly Leu Arg Asn Ser	325	330	335
Pro Gln Arg Glu Thr Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe Ile	340	345	350
Glu Gly Gly Trp Gln Gly Met Val Asp Gly Trp Tyr Gly Tyr His His	355	360	365
Ser Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Lys Glu Ser Thr Gln	370	375	380
Lys Ala Ile Asp Gly Val Thr Asn Lys Val Asn Ser Ile Ile Asp Lys	385	390	395
Met Asn Thr Gln Phe Glu Ala Val Gly Arg Glu Phe Asn Asn Leu Glu	405	410	415
Arg Arg Ile Glu Asn Leu Asn Lys Lys Met Glu Asp Gly Phe Leu Asp	420	425	430
Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Met Glu Asn Glu Arg	435	440	445
Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Asp Lys Val	450	455	460
Arg Leu Gln Leu Arg Asp Asn Ala Lys Glu Leu Gly Asn Gly Cys Phe	465	470	475
			480

-continued

---

Glu Phe Tyr His Lys Cys Asp Asn Glu Cys Met Glu Ser Val Arg Asn  
 485 490 495

Gly Thr Tyr Asp Tyr Pro Gln Tyr Ser Glu Glu Ala Arg Leu Lys Arg  
 500 505 510

Glu Glu Ile Ser Gly Val Lys Leu Glu Ser Ile Gly Thr Tyr Gln Ile  
 515 520 525

Leu Ser Ile Tyr Ser Thr Val Ala Ser Ser Leu Ala Leu Ala Ile Met  
 530 535 540

Val Ala Gly Leu Ser Leu Trp Met Cys Ser Asn Gly Ser Leu Gln Cys  
 545 550 555 560

Arg Ile Cys Ile

<210> SEQ ID NO 14  
 <211> LENGTH: 469  
 <212> TYPE: PRT  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 14

Met Asn Pro Asn Gln Lys Ile Thr Thr Ile Gly Ser Ile Cys Met Val  
 1 5 10 15

Ile Gly Ile Val Ser Leu Met Leu Gln Ile Gly Asn Ile Ile Ser Ile  
 20 25 30

Trp Val Ser His Ser Ile Gln Thr Gly Asn Gln His Gln Ala Glu Pro  
 35 40 45

Cys Asn Gln Ser Ile Ile Thr Tyr Glu Asn Asn Thr Trp Val Asn Gln  
 50 55 60

Thr Tyr Val Asn Ile Ser Asn Thr Asn Phe Leu Thr Glu Lys Ala Val  
 65 70 75 80

Ala Ser Val Thr Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Ser Gly  
 85 90 95

Trp Ala Val Tyr Ser Lys Asp Asn Gly Ile Arg Ile Gly Ser Lys Gly  
 100 105 110

Asp Val Phe Val Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu  
 115 120 125

Cys Arg Thr Phe Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His  
 130 135 140

Ser Asn Gly Thr Val Lys Asp Arg Ser Pro His Arg Thr Leu Met Ser  
 145 150 155 160

Cys Pro Val Gly Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser  
 165 170 175

Val Ala Trp Ser Ala Ser Ala Cys His Asp Gly Thr Ser Trp Leu Thr  
 180 185 190

Ile Gly Ile Ser Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr  
 195 200 205

Asn Gly Ile Ile Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Ile Met  
 210 215 220

Arg Thr Gln Glu Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr  
 225 230 235 240

Val Met Thr Asp Gly Pro Ser Asn Gly Gln Ala Ser Tyr Lys Ile Phe  
 245 250 255

Arg Ile Glu Lys Gly Lys Val Val Lys Ser Ala Glu Leu Asn Ala Pro  
 260 265 270

-continued

---

Asn Tyr His Tyr Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile  
 275 280 285  
 Thr Cys Val Cys Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val  
 290 295 300  
 Ser Phe Asn Gln Asn Leu Glu Tyr Arg Ile Gly Tyr Ile Cys Ser Gly  
 305 310 315 320  
 Val Phe Gly Asp Asn Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly  
 325 330 335  
 Pro Val Ser Pro Lys Gly Ala Tyr Gly Ile Lys Gly Phe Ser Phe Lys  
 340 345 350  
 Tyr Gly Asn Gly Val Trp Ile Gly Arg Thr Lys Ser Thr Asn Ser Arg  
 355 360 365  
 Ser Gly Phe Glu Met Ile Trp Asp Pro Asn Gly Trp Thr Gly Thr Asp  
 370 375 380  
 Ser Asn Phe Ser Val Lys Gln Asp Ile Val Ala Ile Thr Asp Trp Ser  
 385 390 395 400  
 Gly Tyr Ser Gly Ser Phe Val Gln His Pro Glu Leu Thr Gly Leu Asp  
 405 410 415  
 Cys Ile Arg Pro Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys  
 420 425 430  
 Glu Ser Thr Ile Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val  
 435 440 445  
 Asn Ser Asp Thr Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Leu Pro  
 450 455 460  
 Phe Thr Ile Asp Lys  
 465

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 564

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 15

Met Glu Lys Ile Val Leu Leu Leu Ala Thr Val Ser Leu Val Lys Ser  
 1 5 10 15  
 Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val  
 20 25 30  
 Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile  
 35 40 45  
 Leu Glu Arg Thr His Asn Gly Lys Leu Cys Asp Leu Asn Gly Val Lys  
 50 55 60  
 Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn  
 65 70 75 80  
 Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val  
 85 90 95  
 Glu Lys Ala Ser Pro Ala Asn Asp Leu Cys Tyr Pro Gly Asn Phe Asn  
 100 105 110  
 Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu  
 115 120 125  
 Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Ser Asn His Asp Ala Ser  
 130 135 140  
 Ser Gly Val Ser Ser Ala Cys Pro Tyr Leu Gly Arg Ser Ser Phe Phe

-continued

145	150	155	160
Arg Asn Val Val Trp 165	Leu Ile Lys Lys Asn Ser Ser Tyr Pro Thr Ile 170		
Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp 180	185		190
Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Arg Leu Tyr Gln 195	200		205
Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg 210	215		220
Leu Val Pro Glu Ile Ala Thr Arg Pro Lys Val Asn Gly Gln Ser Gly 225	230		235
Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn 245	250		255
Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile 260	265		270
Val Lys Lys Gly Asp Ser Thr Ile Met Lys Ser Glu Leu Glu Tyr Gly 275	280		285
Asn Cys Asn Thr Lys Cys Gln Thr Pro Met Gly Ala Ile Asn Ser Ser 290	295		300
Met Pro Phe His Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys 305	310		315
Tyr Val Lys Ser Asn Arg Leu Val Leu Ala Thr Gly Leu Arg Asn Thr 325	330		335
Pro Gln Arg Glu Thr Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe Ile 340	345		350
Glu Gly Gly Trp Gln Gly Met Val Asp Gly Trp Tyr Gly Tyr His His 355	360		365
Ser Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Glu Ser Thr Gln 370	375		380
Lys Ala Ile Asp Gly Val Thr Asn Lys Val Asn Ser Ile Ile Asn Lys 385	390		395
Met Asn Thr Gln Phe Glu Ala Val Gly Arg Glu Phe Asn Asn Leu Glu 405	410		415
Arg Arg Ile Glu Asn Leu Asn Lys Lys Met Glu Asp Gly Phe Leu Asp 420	425		430
Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Met Glu Asn Glu Arg 435	440		445
Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Asp Lys Val 450	455		460
Arg Leu Gln Leu Arg Asp Asn Ala Lys Glu Leu Gly Asn Gly Cys Phe 465	470		475
Glu Phe Tyr His Lys Cys Asp Asn Glu Cys Met Glu Ser Val Lys Asn 485	490		495
Gly Thr Tyr Asp Tyr Pro Gln Tyr Ser Glu Glu Ala Arg Leu Asn Arg 500	505		510
Glu Glu Ile Ser Gly Val Lys Leu Glu Ser Met Gly Thr Tyr Gln Ile 515	520		525
Leu Ser Ile Tyr Ser Thr Val Ala Ser Ser Leu Ala Leu Ala Ile Met 530	535		540
Val Ala Gly Leu Ser Leu Trp Met Cys Ser Asn Gly Ser Leu Gln Cys 545	550		555
			560

-continued

Arg Ile Cys Ile

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 450

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 16

```

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Cys Met Val
1           5           10           15

Val Gly Ile Ile Ser Leu Met Leu Gln Ile Gly Asn Thr Ile Ser Val
20          25          30

Trp Val Ser His Ile Ile Lys Thr Trp His Pro Asn Gln Pro Glu Pro
35          40          45

Cys Asn Gln Ser Ile Asn Phe Tyr Thr Glu Gln Ala Ala Ala Ser Val
50          55          60

Thr Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Ser Gly Trp Ala Ile
65          70          75          80

Tyr Ser Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly Asp Val Phe
85          90          95

Val Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu Cys Arg Thr
100         105         110

Phe Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly
115         120         125

Thr Val Lys Asp Arg Ser Pro Tyr Arg Thr Leu Met Ser Cys Pro Val
130         135         140

Gly Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp
145         150         155         160

Ser Ala Ser Ala Cys His Asp Gly Ile Ser Trp Leu Thr Ile Gly Ile
165         170         175

Ser Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr Asn Gly Ile
180         185         190

Ile Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Thr Leu Arg Thr Gln
195         200         205

Glu Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr Val Met Thr
210         215         220

Asp Gly Pro Ser Asn Glu Gln Ala Ser Tyr Lys Ile Phe Lys Ile Glu
225         230         235         240

Lys Gly Arg Val Val Lys Ser Val Glu Leu Asn Ala Pro Asn Tyr His
245         250         255

Tyr Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile Thr Cys Val
260         265         270

Cys Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val Ser Phe Asn
275         280         285

Gln Asn Leu Glu Tyr Gln Ile Gly Tyr Ile Cys Ser Gly Val Phe Gly
290         295         300

Asp Ser Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly Pro Val Ser
305         310         315         320

Leu Asn Gly Ala Tyr Gly Val Lys Gly Phe Ser Phe Lys Tyr Gly Asn
325         330         335

Gly Val Trp Ile Gly Arg Thr Lys Ser Thr Ser Ser Arg Ser Gly Phe
340         345         350

```

-continued

---

Glu Met Ile Trp Asp Pro Asn Gly Trp Thr Glu Thr Asp Ser Ser Phe  
 355 360 365  
 Ser Leu Lys Gln Asp Ile Ile Ala Ile Thr Asp Trp Ser Gly Tyr Ser  
 370 375 380  
 Gly Ser Phe Ile Gln His Pro Glu Leu Thr Gly Leu Asn Cys Met Arg  
 385 390 395 400  
 Pro Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys Glu Lys Thr  
 405 410 415  
 Ile Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val Asn Ser Asp  
 420 425 430  
 Thr Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Leu Pro Tyr Thr Ile  
 435 440 445  
 Asp Lys  
 450

<210> SEQ ID NO 17  
 <211> LENGTH: 564  
 <212> TYPE: PRT  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 17

Met Glu Lys Ile Val Leu Leu Leu Ala Thr Val Ser Leu Val Lys Ser  
 1 5 10 15  
 Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val  
 20 25 30  
 Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile  
 35 40 45  
 Leu Glu Arg Thr His Asn Gly Lys Leu Cys Asp Leu Asn Gly Val Lys  
 50 55 60  
 Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn  
 65 70 75 80  
 Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val  
 85 90 95  
 Glu Lys Ala Ser Pro Ala Asn Asp Leu Cys Tyr Pro Gly Asn Phe Asn  
 100 105 110  
 Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu  
 115 120 125  
 Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Ser Asn His Asp Ala Ser  
 130 135 140  
 Ser Gly Val Ser Ser Ala Cys Pro Tyr Leu Gly Arg Ser Ser Phe Phe  
 145 150 155 160  
 Arg Asn Val Val Trp Leu Ile Lys Lys Asn Ser Ser Tyr Pro Thr Ile  
 165 170 175  
 Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp  
 180 185 190  
 Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Arg Leu Tyr Gln  
 195 200 205  
 Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg  
 210 215 220  
 Leu Val Ser Glu Ile Ala Thr Arg Pro Lys Val Asn Gly Gln Ser Gly  
 225 230 235 240  
 Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn



-continued

245					250					255					
Phe	Glu	Ser	Asn	Gly	Asn	Phe	Ile	Ala	Pro	Glu	Tyr	Ala	Tyr	Lys	Ile
			260					265					270		
Val	Lys	Lys	Gly	Asp	Ser	Thr	Ile	Met	Lys	Ser	Glu	Leu	Glu	Tyr	Gly
			275				280						285		
Asn	Cys	Asn	Thr	Lys	Cys	Gln	Thr	Pro	Met	Gly	Ala	Ile	Asn	Ser	Ser
			290				295						300		
Met	Pro	Phe	His	Asn	Ile	His	Pro	Leu	Thr	Ile	Gly	Glu	Cys	Pro	Lys
				310										320	
Tyr	Val	Lys	Ser	Asn	Arg	Leu	Val	Leu	Ala	Thr	Gly	Leu	Arg	Asn	Thr
				325					330					335	
Pro	Gln	Arg	Glu	Thr	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	Phe	Ile
				340				345						350	
Glu	Gly	Gly	Trp	Gln	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	Tyr	His	His
			355				360						365		
Ser	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Glu	Ser	Thr	Gln
			370				375						380		
Lys	Ala	Ile	Asp	Gly	Val	Thr	Asn	Lys	Val	Asn	Ser	Ile	Ile	Asn	Lys
				390										400	
Met	Asn	Thr	Gln	Phe	Glu	Ala	Val	Gly	Arg	Glu	Phe	Asn	Asn	Leu	Glu
				405					410					415	
Arg	Arg	Ile	Glu	Asn	Leu	Asn	Lys	Lys	Met	Glu	Asp	Gly	Phe	Leu	Asp
				420					425					430	
Val	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Met	Glu	Asn	Glu	Arg
				435				440						445	
Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Asp	Lys	Val
							455							460	
Arg	Leu	Gln	Leu	Arg	Asp	Asn	Ala	Lys	Glu	Leu	Gly	Asn	Gly	Cys	Phe
							470							480	
Glu	Phe	Tyr	His	Lys	Cys	Asp	Asn	Glu	Cys	Met	Glu	Ser	Val	Lys	Asn
				485					490					495	
Gly	Thr	Tyr	Asp	Tyr	Pro	Gln	Tyr	Ser	Glu	Glu	Ala	Arg	Leu	Asn	Arg
				500					505					510	
Glu	Glu	Ile	Ser	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Thr	Tyr	Gln	Ile
				515					520					525	
Leu	Ser	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Ala	Leu	Ala	Ile	Met
								535						540	
Val	Ala	Gly	Leu	Ser	Leu	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu	Gln	Cys
								550						555	
														560	

Arg Ile Cys Ile

&lt;210&gt; SEQ ID NO 18

&lt;211&gt; LENGTH: 450

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 18

Met	Asn	Pro	Asn	Gln	Lys	Ile	Ile	Thr	Ile	Gly	Ser	Ile	Cys	Met	Val
1				5					10					15	

Val	Gly	Ile	Ile	Ser	Leu	Met	Leu	Gln	Ile	Gly	Asn	Thr	Ile	Ser	Val
				20				25					30		

Trp	Val	Ser	His	Ile	Ile	Lys	Thr	Trp	His	Pro	Asn	Gln	Pro	Glu	Pro
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-continued

35					40					45					
Cys	Asn	Gln	Ser	Ile	Asn	Phe	Tyr	Thr	Glu	Gln	Ala	Ala	Ser	Val	
50					55					60					
Thr	Leu	Ala	Gly	Asn	Ser	Ser	Leu	Cys	Pro	Ile	Ser	Gly	Trp	Ala	Ile
65				70						75				80	
Tyr	Ser	Lys	Asp	Asn	Ser	Ile	Arg	Ile	Gly	Ser	Lys	Gly	Asp	Val	Phe
				85					90					95	
Val	Ile	Arg	Glu	Pro	Phe	Ile	Ser	Cys	Ser	His	Leu	Glu	Cys	Arg	Thr
			100					105					110		
Phe	Phe	Leu	Thr	Gln	Gly	Ala	Leu	Leu	Asn	Asp	Lys	His	Ser	Asn	Gly
		115					120					125			
Thr	Val	Lys	Asp	Arg	Ser	Pro	Tyr	Arg	Thr	Leu	Met	Ser	Cys	Pro	Val
	130					135					140				
Gly	Glu	Ala	Pro	Ser	Pro	Tyr	Asn	Ser	Arg	Phe	Glu	Ser	Val	Ala	Trp
145					150					155					160
Ser	Ala	Ser	Ala	Cys	His	Asp	Gly	Ile	Ser	Trp	Leu	Thr	Ile	Gly	Ile
				165					170					175	
Ser	Gly	Pro	Asp	Asn	Gly	Ala	Val	Ala	Val	Leu	Lys	Tyr	Asn	Gly	Ile
			180					185					190		
Ile	Thr	Asp	Thr	Ile	Lys	Ser	Trp	Arg	Asn	Asn	Thr	Leu	Arg	Thr	Gln
		195					200					205			
Glu	Ser	Glu	Cys	Ala	Cys	Val	Asn	Gly	Ser	Cys	Phe	Thr	Val	Met	Thr
	210					215					220				
Asp	Gly	Pro	Ser	Asn	Glu	Gln	Ala	Ser	Tyr	Lys	Ile	Phe	Lys	Ile	Glu
225					230					235					240
Lys	Gly	Arg	Val	Val	Lys	Ser	Val	Glu	Leu	Asn	Ala	Pro	Asn	Tyr	His
				245					250					255	
Tyr	Glu	Glu	Cys	Ser	Cys	Tyr	Pro	Asp	Ala	Gly	Glu	Ile	Thr	Cys	Val
			260					265					270		
Cys	Arg	Asp	Asn	Trp	His	Gly	Ser	Asn	Arg	Pro	Trp	Val	Ser	Phe	Asn
		275					280					285			
Gln	Asn	Leu	Glu	Tyr	Gln	Ile	Gly	Tyr	Ile	Cys	Ser	Gly	Val	Phe	Gly
	290					295					300				
Asp	Ser	Pro	Arg	Pro	Asn	Asp	Gly	Thr	Gly	Ser	Cys	Gly	Pro	Val	Ser
305					310					315					320
Leu	Asn	Gly	Ala	Tyr	Gly	Val	Lys	Gly	Phe	Ser	Phe	Lys	Tyr	Gly	Asn
				325					330					335	
Gly	Val	Trp	Ile	Gly	Arg	Thr	Lys	Ser	Thr	Ser	Ser	Arg	Ser	Gly	Phe
			340					345					350		
Glu	Met	Ile	Trp	Asp	Pro	Asn	Gly	Trp	Thr	Glu	Thr	Asp	Ser	Ser	Phe
		355					360					365			
Ser	Leu	Lys	Gln	Asp	Ile	Ile	Ala	Ile	Thr	Asp	Trp	Ser	Gly	Tyr	Ser
	370					375					380				
Gly	Ser	Phe	Ile	Gln	His	Pro	Glu	Leu	Thr	Gly	Leu	Asn	Cys	Met	Arg
385					390					395					400
Pro	Cys	Phe	Trp	Val	Glu	Leu	Ile	Arg	Gly	Arg	Pro	Lys	Glu	Lys	Thr
				405					410					415	
Ile	Trp	Thr	Ser	Gly	Ser	Ser	Ile	Ser	Phe	Cys	Gly	Val	Asn	Ser	Asp
			420					425					430		
Thr	Val	Gly	Trp	Ser	Trp	Pro	Asp	Gly	Ala	Glu	Leu	Pro	Tyr	Thr	Ile
		435					440					445			

---

-continued

---

Asp Lys  
450

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 558

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 19

Met Glu Ala Ile Pro Leu Ile Thr Ile Leu Leu Val Val Thr Ala Ser  
1 5 10 15

Asn Ala Asp Lys Ile Cys Ile Gly Tyr Gln Ser Thr Asn Ser Thr Glu  
20 25 30

Thr Val Asp Thr Leu Thr Glu Asn Asn Val Pro Val Thr His Ala Lys  
35 40 45

Glu Leu Leu His Thr Glu His Asn Gly Met Leu Cys Ala Thr Asn Leu  
50 55 60

Gly Arg Pro Leu Ile Leu Asp Thr Cys Thr Ile Glu Gly Leu Ile Tyr  
65 70 75 80

Gly Asn Pro Ser Cys Asp Leu Leu Leu Gly Gly Arg Glu Trp Ser Tyr  
85 90 95

Ile Val Glu Arg Pro Ser Ala Val Asn Gly Met Cys Tyr Pro Gly Asn  
100 105 110

Val Glu Asn Leu Glu Glu Leu Arg Ser Phe Phe Ser Ser Ala Ser Ser  
115 120 125

Tyr Gln Arg Ile Gln Ile Phe Pro Asp Thr Ile Trp Asn Val Ser Tyr  
130 135 140

Ser Gly Thr Ser Lys Ala Cys Ser Asp Ser Phe Tyr Arg Ser Met Arg  
145 150 155 160

Trp Leu Thr Gln Lys Asn Asn Ala Tyr Pro Ile Gln Asp Ala Gln Tyr  
165 170 175

Thr Asn Asn Arg Gly Lys Ser Ile Leu Phe Met Trp Gly Ile Asn His  
180 185 190

Pro Pro Thr Asp Thr Ala Gln Thr Asn Leu Tyr Thr Arg Thr Asp Thr  
195 200 205

Thr Thr Ser Val Ala Thr Glu Asp Ile Asn Arg Thr Phe Lys Pro Val  
210 215 220

Ile Gly Pro Arg Pro Leu Val Asn Gly Leu Gln Gly Arg Ile Asp Tyr  
225 230 235 240

Tyr Trp Ser Val Leu Lys Pro Gly Gln Thr Leu Arg Val Arg Ser Asn  
245 250 255

Gly Asn Leu Ile Ala Pro Trp Tyr Gly His Ile Leu Ser Gly Glu Ser  
260 265 270

His Gly Arg Ile Leu Lys Thr Asp Leu Asn Ser Gly Ser Cys Val Val  
275 280 285

Gln Cys Gln Thr Glu Arg Gly Gly Leu Asn Thr Thr Leu Pro Phe His  
290 295 300

Asn Val Ser Lys Tyr Ala Phe Gly Asn Cys Pro Lys Tyr Val Gly Val  
305 310 315 320

Lys Ser Leu Lys Leu Ala Val Gly Leu Arg Asn Val Pro Ala Arg Ser  
325 330 335

Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe Ile Glu Gly Gly Trp

-continued

340					345					350					
Ser	Gly	Leu	Val	Ala	Gly	Trp	Tyr	Gly	Phe	Gln	His	Ser	Asn	Asp	Gln
		355					360					365			
Gly	Val	Gly	Ile	Ala	Ala	Asp	Arg	Asp	Ser	Thr	Gln	Arg	Ala	Ile	Asp
		370					375					380			
Lys	Ile	Thr	Ser	Lys	Val	Asn	Asn	Ile	Val	Asp	Lys	Met	Asn	Lys	Gln
		385					390					395			400
Tyr	Glu	Ile	Ile	Asp	His	Glu	Phe	Ser	Glu	Val	Glu	Asn	Arg	Leu	Asn
				405					410					415	
Met	Ile	Asn	Asn	Lys	Ile	Asp	Asp	Gln	Ile	Gln	Asp	Ile	Trp	Ala	Tyr
				420					425					430	
Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn	Gln	Lys	Thr	Leu	Asp	Glu
				435					440					445	
His	Asp	Ala	Asn	Val	Asn	Asn	Leu	Tyr	Asn	Lys	Val	Lys	Arg	Ala	Leu
				450					455					460	
Gly	Ser	Asn	Ala	Met	Glu	Asp	Gly	Lys	Gly	Cys	Phe	Glu	Leu	Tyr	His
				465					470					475	480
Lys	Cys	Asp	Asp	Gln	Cys	Met	Glu	Thr	Ile	Arg	Asn	Gly	Thr	Tyr	Asn
				485					490					495	
Arg	Arg	Lys	Tyr	Lys	Glu	Glu	Ser	Arg	Leu	Glu	Arg	Gln	Lys	Ile	Glu
				500					505					510	
Gly	Val	Lys	Leu	Glu	Ser	Glu	Gly	Thr	Tyr	Lys	Ile	Leu	Thr	Ile	Tyr
				515					520					525	
Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Ile	Ala	Met	Gly	Phe	Ala	Ala	Phe
				530					535					540	
Leu	Phe	Trp	Ala	Met	Ser	Asn	Gly	Ser	Cys	Arg	Cys	Asn	Ile		
				545					550					555	

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 469

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 20

Met	Asn	Pro	Asn	Gln	Lys	Ile	Ile	Ala	Ile	Gly	Ser	Val	Ser	Leu	Thr
1				5					10					15	
Ile	Ala	Thr	Ile	Cys	Leu	Leu	Met	Gln	Ile	Ala	Ile	Leu	Ala	Thr	Thr
			20					25					30		
Met	Thr	Leu	His	Phe	Lys	Gln	Asn	Glu	Cys	Ile	Asn	Ser	Ser	Asn	Asn
			35				40						45		
Gln	Val	Val	Pro	Cys	Glu	Pro	Ile	Ile	Ile	Glu	Arg	Asn	Ile	Thr	Glu
			50				55						60		
Ile	Val	His	Leu	Asn	Ser	Thr	Thr	Leu	Glu	Lys	Glu	Ile	Cys	Pro	Lys
				70					75					80	
Val	Ala	Asp	Tyr	Arg	Asn	Trp	Ser	Lys	Pro	Gln	Cys	Gln	Ile	Thr	Gly
				85					90					95	
Phe	Ala	Pro	Phe	Ser	Lys	Asp	Asn	Ser	Ile	Arg	Leu	Ser	Ala	Gly	Gly
				100				105						110	
Asp	Ile	Trp	Val	Thr	Arg	Glu	Pro	Tyr	Val	Ser	Cys	Gly	Leu	Gly	Lys
				115				120					125		
Cys	Tyr	Gln	Phe	Ala	Leu	Gly	Gln	Gly	Thr	Thr	Leu	Glu	Asn	Lys	His
				130				135						140	

```
<210> SEQ ID NO 21
<211> LENGTH: 1737
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 21
```

```

agcaaaagca ggggatacaa aatgaacact caaatcctgg tattcgctct ggtggcgagc      60
attccgacaa atgcagacaa gatctgcctt gggcatcatg cgtgtcaaa cgggactaaa    120
gtaaacacat taactgagag aggagtggaa gtcgttaatg caactgaaac ggtggaacga    180

```

-continued

---

acaaacgttc ccaggatctg ctcaaaaggg aaaaggacag ttgacctcgg tcaatgtgga	240
cttctgggaa caatcactgg gccaccccaa tgtgaccaat tctagaatt ttcggccgac	300
ttaattattg agaggcgaga aggaagtgat gtctgttata ctgggaaatt cgtgaatgaa	360
gaagctctga ggcaattct cagagagtca ggcggaattg acaaggagac aatgggattc	420
acctacagcg gaataagaac taatggaaca accagtgcac gtaggagatc aggatcttca	480
ttctatgcag agatgaaatg gctcctgtca aacacagaca atgctgcttt cccgcaaatg	540
actaagtcac acaagaacac aaggaaagac ccagctctga taatatgggg gatccaccat	600
tccggatcaa ctacagaaca gaccaagcta tatgggagtg gaaacaaact gataacagtt	660
gggagttcta attaccaaca gtcctttgta ccgagtcacg gagcgagacc acaagtgaat	720
ggccaatctg gaagaattga ctttcattgg ctgatactaa accctaatga cagggtcact	780
ttcagtttca atggggcctt catagctcca gaccgtgcaa gctttctgag aggggaagtcc	840
atgggaattc agagtgaagt acagggtgat gccaatgtg aaggagattg ctatcatagt	900
ggagggacaa taataagtaa tttgcccttt cagaacataa atagcagggc agtaggaaaa	960
tgctcgagat atgttaagca agagagtctg ctgttggaac caggaatgaa gaatgttccc	1020
gaaatcccaa agaggaggag gagaggccta tttggtgcta tagcgggttt cattgaaaat	1080
ggatgggaag gtttgattga tgggtggtat ggcttcaggc atcaaatgc acaaggggag	1140
ggaactgctg cagattacaa aagcacccaa tcagcaattg atcaataaac agggaaatta	1200
aatcggctta tagaaaaaac taaccaacag tttgagttaa tagacaacga attcactgag	1260
gttgaaaggc aaattggcaa tgtgataaac tggaccagag attccatgac agaagtgtgg	1320
tcctataacg ctgaactctt agtagcaatg gagaatcagc acacaattga tctggccgac	1380
tcagaaatga acaaactgta cgaacgagtg aagagacaac tgagagagaa tgccgaagaa	1440
gatggcactg gttgcttcga aatatttcac aagtgtgatg acgactgcac ggccagtatt	1500
agaaacaaca cctatgatca cagcaagtac agggagaag caatacaaaa tagaatacag	1560
attgacccag tcaaactaag cagcggctac aaagatgtga tactttggtt tagcttcggg	1620
gcacatgtt tcatacttct ggccattgca atgggccttg tcttcatatg tgtgaagaat	1680
ggaaacatgc ggtgcactat ttgtatataa gtttgaaaa acacccttgt ttctact	1737

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 1465

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 22

agcaaaagca gggatgatga gaatgaatcc aaatcagaaa ctatttgcac tatctggagt	60
ggcaatagca cttagtgtac tgaacttatt gataggaatc tcaaacgtcg gattgaacgt	120
atctctacat ctaaaggaaa aaggacccaa acaggaggag aatttaacat gcacgaccat	180
taatcaaaac aacactactg tagtagaaaa cacatatgta aataatacaa caataattac	240
caagggaact gatttgaaaa caccaagcta tctgctgttg aacaagagcc tgtgcaatgt	300
tgaagggtgg gtcgtgatag caaaagacaa tgcagtaaga tttggggaaa gtgaacaaat	360
cattgttacc agggagccat atgtatcatg cgacccaaca ggatgcaaaa tgtatgcctt	420
gcaccaaggg actaccatta ggaacaaaca ttcaaatgga acgattcatg acagaacagc	480

-continued

---

```

tttcagaggt ctcatctcca ctccattggg cactccacca accgtaagta acagtgactt 540
tatgtgtgtt ggatgggtcaa gcacaacttg ccatgatggg attgctagga tgactatctg 600
tatacaagga aataatgaca atgctacagc aacggtttat tacaacagaa ggctgaccac 660
taccattaag acctgggcca gaaacattct gaggactcaa gaatcagaat gtgtgtgcca 720
caatggcaca tgtgcagttg taatgaccga cggatcggct agtagtcaag cctatacaaa 780
agtaatgtat ttccacaagg gattagtagt taaggaggag gagttaaggg gatcagccag 840
acatattgag gaatgctcct gttatggaca caatcaaaag gtgacctgtg tgtgcagaga 900
taactggcag ggagcaaaca ggcctattat agaaattgat atgagcacat tggagcacac 960
aagtagatac gtgtgcactg gaattctcac agacaccagc agacctgggg acaaatctag 1020
tggtgattgt tccaatccaa taactgggag tcccggcggt cggggagtga agggattcgg 1080
gtttctaaat ggggataaca catggcttgg taggaccatc agccccagat caagaagtgg 1140
attcgaaatg ttgaaaatac ctaatgcagg tactgatccc aattctagaa tagcagaacg 1200
acaggaaatt gtcgacaata acaattggtc aggtattccc ggaagcttta ttgactattg 1260
gaatgataac agtgaatgct acaatccatg cttttacgta gagttaatta gaggaagacc 1320
cgaagaggct aaatacgtat ggtgggcaag taacagtcta attgccctat gtggaagccc 1380
attcccagtt gggctctggt ccttccccga tggggcacia atccaatact tttcgtaaaa 1440
tgcaaaaaaa ctctctgttt ctact 1465

```

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 1754

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 23

```

agcaaaagca ggggatacaa aatgaatact caaattttgg cattcattgc ttgtatgctg 60
attggaacta aaggagacaa aatatgtctt gggcaccatg ctgtggcaaa tgggacaaaa 120
gtgaacacac taacagagag gggaattgaa gtagtcaatg ccacggagac ggtggaaact 180
gtaaatatta aaaaaatag cactcaagga aaaaggccaa cagatctggg acaatgtgga 240
cttctaggaa ccctaatagg acctcccaa tgcgatcaat ttctggagtt tgacgcta 300
ttgataattg aacgaagaga aggaaccgat gtgtgctatc ccgggaagtt cacaaatgaa 360
gaatcactga ggcagatcct tcgaggggca ggaggaattg ataaagagtc aatgggtttc 420
acctatagtg gaataagaac caatggggcg acgagtgcct gcagaagatc aggttcttct 480
ttctatgcgg agatgaagtg gttactgtcg aattcagaca atgcggcatt tccccaaatg 540
actaagtcgt ataggaatcc caggaacaaa ccagctctga taatctgggg agtgcacac 600
tctggatcag ctactgagca gaccaaactc tatggaagtg gaaacaagtt gataacagta 660
ggaagctcga aataccagca atcattcact ccaagtccgg gagcacggcc acaagtgaat 720
ggacaatcag gaaggattga ttttcatttg ctactccttg accccaatga cacagtgacc 780
ttcactttca atggggcatt catagccctt gacagggcaa gtttcttttag aggagaatcg 840
ctaggagtcc agagtgatgt tcctttggat tctggttgtg aaggggattg ctccacagt 900
gggggtacga tagtcagttc cctgccatcc caaaacatca accctagaac agtggggaaa 960
tgccctcgat atgtcaaaca gacaagcctc cttttggcta caggaatgag aaacgtccca 1020

```

-continued

---

```

gagaacccca agcaggccta ccggaacgag atgaccagag gccttttttg agcgattgct 1080
ggattcatag agaatggatg ggaagggtctc atcgatggat ggtatggttt cagacatcaa 1140
aatgcacaag gagaaggaac tgcagctgac tacaaaagca cccaatctgc aatagatcag 1200
atcacaggca aattgaatcg tctgattgac aaaacaaacc agcagtttga actgatagac 1260
aatgaattca gtgagataga acaacaaatc gggaaatgtca ttaactggac acgagactca 1320
atgactgagg tatggtcgta taatgctgag ctggtggtgg caatggagaa tcagcataca 1380
atagatcttg cagactcaga aatgaacaaa ctttacgaac gcgtcagaaa acaactaagg 1440
gaaaatgctg aagaatagg aactggatgc tttgagatat tccataagtg tgatgatcag 1500
tgtatggaga gcataaggaa caacacttat gaccataccc aatacaggac agagtcattg 1560
cagaatagaa tacagataga ccagtgaaa ttgagtagtg gatacaaaga cataatctta 1620
tggttttagct tcggggcctc atgtttttctt cttctagcca ttgcaatggg attggttttc 1680
atttgcataa agaatggaaa catgcggtgc actatttgta tatagtttga gaaaaaaca 1740
cccttgtttc tact 1754

```

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 1453

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 24

```

agcaaaagca ggtgcgagat gaatccgaat cagaagataa taacaatcgg ggtagtgaat 60
accactctgt caacaatagc cttctctatt ggagtgggaa acttagtttt caacacagtc 120
atacatgaga aaataggaga ccatcaaata gtgacccatc caacaataat gaccctgaa 180
gtaccgaact gcagtgcacac tataataaca tacaataaca ctgttataaa caacataaca 240
acaacaataa taactgaagc agaaaggcct ttcaagtctc cactaccgct gtgcccttc 300
agaggattct tcccttttca caaggacaat gcaatacgac tgggtgaaaa caaagacgtc 360
atagtcacaa gggagcctta tgtagctgc gataatgaca actgctggtc ctttgctctc 420
gcacaaggag cattgctagg gactaaacat agcaatggga ccattaaaga cagaacacca 480
tataggtctc taattcgttt cccaatagga acagctccag tactaggaaa ttacaaagag 540
atatgcattg cttggtcgag cagcagttgc tttgacggga aagagtggat gcatgtgtgc 600
atgacaggga atgataatga tgcaagtgcc cagataatat atggaggaag aatgacagac 660
tccattaaat catggaggaa ggacatacta agaaccagg agtctgaatg tcaatgcatt 720
gacgggactt gtgttgttgc tgtcacagat ggcctgctg ctaatagtgc agatcacagg 780
gtttactgga tacggggagg aagaataata aagtatgaaa atgttccaa aacaaagata 840
caacacttag aagaatgttc ctgctatgtg gacattgatg tttactgtat atgtagggac 900
aattggaagg gctctaacag acottggatg agaatcaaca acgagactat actggaacaa 960
ggatatgtat gtagtaaat tcactcagac acccccaggc cagctgaccc ttcaataatg 1020
tcatgtgact ccccaagcaa tgtcaatgga ggacccggag tgaaggggtt tggtttcaa 1080
gctggcaatg atgtatggtt aggtagaaca gtgtcaacta gtggtagatc gggctttgaa 1140
attatcaaag ttacagaagg gtggatcaac tctcctaacc atgtcaaatc aattacacaa 1200
acactagtgt ccaacaatga ctggtcaggc tattcaggtg gcttcattgt caaagccaag 1260

```



-continued

---

gactgttttc agccctgttt ttatgttgag cttatacgag ggaggcccaa caagaatgat	1320
gacgtctctt ggacaagtaa tagtatagtt actttctgtg gactagacaa tgaacctgga	1380
tcgggaaatt ggccagatgg ttctaacatt ggggtttatgc ccaagtaata gaaaaaagca	1440
ccttgtttct act	1453

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 1733

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 25

agcgaaagca ggggatacaa aatgaatact caaattttgg cattcattgc ttgtatgctg	60
attggaacta aaggagacaa aatatgtctt gggcaccatg ctgtggcaaa tgggacaaaa	120
gtgaacacac taacagagag gggaattgaa gtagtcaatg ccacggagac ggtggaaact	180
gtaaattaa agaaaatatg cactcaagga aaaaggccaa cagatctggg acaatgtgga	240
cttctaggaa ccctaatagg acctcccaa tgcgatcaat ttctggagtt tgacgcta	300
ttgataattg aacgaagaga aggaaccgat gtgtgctatc ccgggaagtt cacaaatgaa	360
gaatcactga ggcagatcct tcgagggta ggaggaattg ataaagagtc aatgggtttc	420
acctatagtg gaataagaac caatggggcg acgagtgcct gcagaagatc aggttcttct	480
ttctatgcgg agatgaagtg gttactgtcg aattcagaca atgcggcatt tccccaaatg	540
actaagtcgt ataggaatcc caggaacaaa ccagctctga taatctgggg agtgcacac	600
tctggatcag ctactgagca gaccaaactc tatggaagtg gaaacaagtt gataacagta	660
ggaagctcga aataccagca atcattcact ccaagtccgg gagcacggcc acaagtgaat	720
ggacaatcag gaaggattga ttttcattgg ctactccttg accccaatga cacagtgacc	780
ttcactttca atggggcatt catagccctt gacagggcaa gtttcttttag aggagaaatc	840
ctaggagtcc agagtgatgt tcctttggat tctggttggtg aaggggattg ctccacagt	900
gggggtacga tagtcagttc cctgccatcc caaaacatca accctagaac agtggggaaa	960
tgccctcgat atgtcaaaca gacaagcctc cttttggcta caggaatgag aaacgtccca	1020
gagaacccca agaccagagg cctttttgga gcgattgctg gattcataga gaatggatgg	1080
gaaggtctca tcgatggatg gtatggttcc agacatcaa atgcacaagg agaaggaact	1140
gcagctgact acaaaagcac ccaatctgca atagatcaga tcacaggcaa attgaatcgt	1200
ctgattgaca aaacaaacca gcagtttgaa ctgatagaca atgaattcag tgagatagaa	1260
caacaaatcg ggaatgtcat taactggaca cgagactcaa tgactgaggt atggtcgtat	1320
aatgctgagc tgttggtggc aatggagaat cagcatacaa tagatcttgc agactcagaa	1380
atgaacaaac ttacgaacg cgtcagaaaa caactaaggg aaaatgctga agaagatgga	1440
actggatgct ttgagatatt ccataagtgt gatgatcagt gtatggagag cataaggaac	1500
aaacttatg accataccca atacaggaca gagtcattgc agaatagaat acagatagac	1560
ccagtgaat tgagttagtg atacaaagac ataattctat ggtttagctt cggggcatca	1620
tggtttcttc ttctagccat tgcaatggga ttggttttca ttgcataaa gaatggaaac	1680
atcggtgca ctatttgat atagtttgag aaaaaaacac ccttgtttct act	1733

-continued

<210> SEQ ID NO 26  
 <211> LENGTH: 1453  
 <212> TYPE: DNA  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 26

```

agcaaaagca ggtgcgagat gaatccgaat cagaagataa taacaatcgg ggtagtgaat      60
accactctgt caacaatagc cttcttcatt ggagtgggaa acttagtttt caacacagtc     120
atacatgaga aaataggaga ccatcaaata gtgacccatc caacaataat gacccctgaa     180
gtaccgaact gcagtgcac tataataaca tacaataaca ctgttataaa caacataaca     240
acaacaataa taactgaagc agaaaggcct ttcaagtctc cactaccgct gtgcccttc      300
agaggattct tcccttttca caaggacaat gcaatacgac tgggtgaaaa caaagacgtc     360
atagtcacaa gggagcctta tgtagctgc gataatgaca actgctggtc ctttgctctc     420
gcacaaggag cattgctagg gactaaacat agcaatggga ccattaaaga cagaacacca     480
tataggcttc taattcgttt cccaatagga acagctccag tactaggaaa ttacaaagag     540
atatgcattg ctgtgcgag cagcagttgc tttgacggga aagagtggat gcatgtgtgc     600
atgacagggg atgataatga tgcaagtgcc cagataatat atggaggaag aatgacagac     660
tccattaaat catggaggaa ggacatacta agaaccagg agtctgaatg tcaatgcatt     720
gacgggactt gtgtgtgtgc tgtcacagat ggccctgctg ctaatagtgc agatcacagg     780
gtttactgga tacgggaggg aagaataata aagtatgaaa atgttcccaa aacaaagata     840
caacacttag aagaatgttc ctgctatgtg gacattgatg tttactgtat atgtagggac     900
aattggaagg gctctaacag accttgatg agaatcaaca acgagactat actggaacaa     960
ggatatgtat gtagtaaatt tcaactcagc acccccaggc cagctgaccc ttcaataatg    1020
tcatgtgact ccccaagcaa tgtcaatgga ggacccggag tgaaggggtt tggtttcaaa    1080
gctggcaatg atgtatgggt aggtagaaca gtgtcaacta gtggtagatc gggctttgaa    1140
attatcaaag ttacagaagg gtggatcaac tctcctaacc atgtcaaatc aattacacaa    1200
acactagtgt ccaacaatga ctggtcaggc tattcaggta gcttcattgt caaagccaag    1260
gactgttttc agccctgttt ttatgttgag cttatacgag ggaggcccaa caagaatgat    1320
gacgtctctt ggacaagtaa tagtatagtt actttctgtg gactagacaa tgaacctgga    1380
tcgggaaatt ggccagatgg ttctaacatt gggtttatgc ccaagtaata gaaaaaagca    1440
cttgtttct act                                     1453

```

<210> SEQ ID NO 27  
 <211> LENGTH: 562  
 <212> TYPE: PRT  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 27

```

Met Asn Thr Gln Ile Leu Val Phe Ala Leu Val Ala Ser Ile Pro Thr
1           5           10           15

Asn Ala Asp Lys Ile Cys Leu Gly His His Ala Val Ser Asn Gly Thr
          20           25           30

Lys Val Asn Thr Leu Thr Glu Arg Gly Val Glu Val Val Asn Ala Thr
          35           40           45

Glu Thr Val Glu Arg Thr Asn Val Pro Arg Ile Cys Ser Lys Gly Lys
          50           55           60

```

-continued

---

Arg	Thr	Val	Asp	Leu	Gly	Gln	Cys	Gly	Leu	Leu	Gly	Thr	Ile	Thr	Gly
65					70					75					80
Pro	Pro	Gln	Cys	Asp	Gln	Phe	Leu	Glu	Phe	Ser	Ala	Asp	Leu	Ile	Ile
			85						90					95	
Glu	Arg	Arg	Glu	Gly	Ser	Asp	Val	Cys	Tyr	Pro	Gly	Lys	Phe	Val	Asn
			100					105					110		
Glu	Glu	Ala	Leu	Arg	Gln	Ile	Leu	Arg	Glu	Ser	Gly	Gly	Ile	Asp	Lys
		115					120					125			
Glu	Thr	Met	Gly	Phe	Thr	Tyr	Ser	Gly	Ile	Arg	Thr	Asn	Gly	Thr	Thr
	130						135					140			
Ser	Ala	Cys	Arg	Arg	Ser	Gly	Ser	Ser	Phe	Tyr	Ala	Glu	Met	Lys	Trp
145					150					155					160
Leu	Leu	Ser	Asn	Thr	Asp	Asn	Ala	Ala	Phe	Pro	Gln	Met	Thr	Lys	Ser
			165						170					175	
Tyr	Lys	Asn	Thr	Arg	Lys	Asp	Pro	Ala	Leu	Ile	Ile	Trp	Gly	Ile	His
		180						185					190		
His	Ser	Gly	Ser	Thr	Thr	Glu	Gln	Thr	Lys	Leu	Tyr	Gly	Ser	Gly	Asn
	195						200					205			
Lys	Leu	Ile	Thr	Val	Gly	Ser	Ser	Asn	Tyr	Gln	Gln	Ser	Phe	Val	Pro
	210					215					220				
Ser	Pro	Gly	Ala	Arg	Pro	Gln	Val	Asn	Gly	Gln	Ser	Gly	Arg	Ile	Asp
225					230					235					240
Phe	His	Trp	Leu	Ile	Leu	Asn	Pro	Asn	Asp	Thr	Val	Thr	Phe	Ser	Phe
			245						250					255	
Asn	Gly	Ala	Phe	Ile	Ala	Pro	Asp	Arg	Ala	Ser	Phe	Leu	Arg	Gly	Lys
		260					265						270		
Ser	Met	Gly	Ile	Gln	Ser	Glu	Val	Gln	Val	Asp	Ala	Asn	Cys	Glu	Gly
	275					280						285			
Asp	Cys	Tyr	His	Ser	Gly	Gly	Thr	Ile	Ile	Ser	Asn	Leu	Pro	Phe	Gln
	290					295					300				
Asn	Ile	Asn	Ser	Arg	Ala	Val	Gly	Lys	Cys	Pro	Arg	Tyr	Val	Lys	Gln
305				310						315					320
Glu	Ser	Leu	Leu	Leu	Ala	Thr	Gly	Met	Lys	Asn	Val	Pro	Glu	Ile	Pro
			325						330					335	
Lys	Arg	Arg	Arg	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	Phe	Ile	Glu
			340				345						350		
Asn	Gly	Trp	Glu	Gly	Leu	Ile	Asp	Gly	Trp	Tyr	Gly	Phe	Arg	His	Gln
	355					360						365			
Asn	Ala	Gln	Gly	Glu	Gly	Thr	Ala	Ala	Asp	Tyr	Lys	Ser	Thr	Gln	Ser
	370					375					380				
Ala	Ile	Asp	Gln	Ile	Thr	Gly	Lys	Leu	Asn	Arg	Leu	Ile	Glu	Lys	Thr
385					390					395					400
Asn	Gln	Gln	Phe	Glu	Leu	Ile	Asp	Asn	Glu	Phe	Thr	Glu	Val	Glu	Arg
			405						410					415	
Gln	Ile	Gly	Asn	Val	Ile	Asn	Trp	Thr	Arg	Asp	Ser	Met	Thr	Glu	Val
			420					425					430		
Trp	Ser	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Ala	Met	Glu	Asn	Gln	His	Thr
	435					440						445			
Ile	Asp	Leu	Ala	Asp	Ser	Glu	Met	Asn	Lys	Leu	Tyr	Glu	Arg	Val	Lys
	450					455					460				

-continued

---

Arg	Gln	Leu	Arg	Glu	Asn	Ala	Glu	Glu	Asp	Gly	Thr	Gly	Cys	Phe	Glu
465					470					475					480
Ile	Phe	His	Lys	Cys	Asp	Asp	Asp	Cys	Met	Ala	Ser	Ile	Arg	Asn	Asn
			485						490					495	
Thr	Tyr	Asp	His	Ser	Lys	Tyr	Arg	Glu	Glu	Ala	Ile	Gln	Asn	Arg	Ile
			500					505					510		
Gln	Ile	Asp	Pro	Val	Lys	Leu	Ser	Ser	Gly	Tyr	Lys	Asp	Val	Ile	Leu
		515					520					525			
Trp	Phe	Ser	Phe	Gly	Ala	Ser	Cys	Phe	Ile	Leu	Leu	Ala	Ile	Ala	Met
	530					535						540			
Gly	Leu	Val	Phe	Ile	Cys	Val	Lys	Asn	Gly	Asn	Met	Arg	Cys	Thr	Ile
545					550					555					560

Cys Ile

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 471

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 28

Met	Asn	Pro	Asn	Gln	Lys	Leu	Phe	Ala	Leu	Ser	Gly	Val	Ala	Ile	Ala
1			5						10					15	
Leu	Ser	Val	Leu	Asn	Leu	Leu	Ile	Gly	Ile	Ser	Asn	Val	Gly	Leu	Asn
			20				25						30		
Val	Ser	Leu	His	Leu	Lys	Glu	Lys	Gly	Pro	Lys	Gln	Glu	Glu	Asn	Leu
		35				40						45			
Thr	Cys	Thr	Thr	Ile	Asn	Gln	Asn	Asn	Thr	Thr	Val	Val	Glu	Asn	Thr
	50				55						60				
Tyr	Val	Asn	Asn	Thr	Thr	Ile	Ile	Thr	Lys	Gly	Thr	Asp	Leu	Lys	Thr
65				70						75					80
Pro	Ser	Tyr	Leu	Leu	Asn	Lys	Ser	Leu	Cys	Asn	Val	Glu	Gly	Trp	
			85					90					95		
Val	Val	Ile	Ala	Lys	Asp	Asn	Ala	Val	Arg	Phe	Gly	Glu	Ser	Glu	Gln
			100				105						110		
Ile	Ile	Val	Thr	Arg	Glu	Pro	Tyr	Val	Ser	Cys	Asp	Pro	Thr	Gly	Cys
		115				120						125			
Lys	Met	Tyr	Ala	Leu	His	Gln	Gly	Thr	Thr	Ile	Arg	Asn	Lys	His	Ser
	130				135						140				
Asn	Gly	Thr	Ile	His	Asp	Arg	Thr	Ala	Phe	Arg	Gly	Leu	Ile	Ser	Thr
145				150						155					160
Pro	Leu	Gly	Thr	Pro	Pro	Thr	Val	Ser	Asn	Ser	Asp	Phe	Met	Cys	Val
			165					170						175	
Gly	Trp	Ser	Ser	Thr	Thr	Cys	His	Asp	Gly	Ile	Ala	Arg	Met	Thr	Ile
		180				185							190		
Cys	Ile	Gln	Gly	Asn	Asn	Asp	Asn	Ala	Thr	Ala	Thr	Val	Tyr	Tyr	Asn
	195					200						205			
Arg	Arg	Leu	Thr	Thr	Thr	Ile	Lys	Thr	Trp	Ala	Arg	Asn	Ile	Leu	Arg
	210					215					220				
Thr	Gln	Glu	Ser	Glu	Cys	Val	Cys	His	Asn	Gly	Thr	Cys	Ala	Val	Val
225				230						235					240
Met	Thr	Asp	Gly	Ser	Ala	Ser	Ser	Gln	Ala	Tyr	Thr	Lys	Val	Met	Tyr
				245				250						255	

-continued

---

Phe His Lys Gly Leu Val Val Lys Glu Glu Glu Leu Arg Gly Ser Ala  
                   260                  265                  270  
 Arg His Ile Glu Glu Cys Ser Cys Tyr Gly His Asn Gln Lys Val Thr  
                   275                  280                  285  
 Cys Val Cys Arg Asp Asn Trp Gln Gly Ala Asn Arg Pro Ile Ile Glu  
                   290                  295                  300  
 Ile Asp Met Ser Thr Leu Glu His Thr Ser Arg Tyr Val Cys Thr Gly  
 305                  310                  315                  320  
 Ile Leu Thr Asp Thr Ser Arg Pro Gly Asp Lys Ser Ser Gly Asp Cys  
                   325                  330                  335  
 Ser Asn Pro Ile Thr Gly Ser Pro Gly Val Pro Gly Val Lys Gly Phe  
                   340                  345                  350  
 Gly Phe Leu Asn Gly Asp Asn Thr Trp Leu Gly Arg Thr Ile Ser Pro  
                   355                  360                  365  
 Arg Ser Arg Ser Gly Phe Glu Met Leu Lys Ile Pro Asn Ala Gly Thr  
                   370                  375                  380  
 Asp Pro Asn Ser Arg Ile Ala Glu Arg Gln Glu Ile Val Asp Asn Asn  
 385                  390                  395                  400  
 Asn Trp Ser Gly Tyr Ser Gly Ser Phe Ile Asp Tyr Trp Asn Asp Asn  
                   405                  410                  415  
 Ser Glu Cys Tyr Asn Pro Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg  
                   420                  425                  430  
 Pro Glu Glu Ala Lys Tyr Val Trp Trp Ala Ser Asn Ser Leu Ile Ala  
                   435                  440                  445  
 Leu Cys Gly Ser Pro Phe Pro Val Gly Ser Gly Ser Phe Pro Asp Gly  
                   450                  455                  460  
 Ala Gln Ile Gln Tyr Phe Ser  
 465                  470

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 567

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 29

Met Asn Thr Gln Ile Leu Ala Phe Ile Ala Cys Met Leu Ile Gly Thr  
 1                  5                  10                  15  
 Lys Gly Asp Lys Ile Cys Leu Gly His His Ala Val Ala Asn Gly Thr  
                   20                  25                  30  
 Lys Val Asn Thr Leu Thr Glu Arg Gly Ile Glu Val Val Asn Ala Thr  
                   35                  40                  45  
 Glu Thr Val Glu Thr Val Asn Ile Lys Lys Ile Cys Thr Gln Gly Lys  
                   50                  55                  60  
 Arg Pro Thr Asp Leu Gly Gln Cys Gly Leu Leu Gly Thr Leu Ile Gly  
 65                  70                  75                  80  
 Pro Pro Gln Cys Asp Gln Phe Leu Glu Phe Asp Ala Asn Leu Ile Ile  
                   85                  90                  95  
 Glu Arg Arg Glu Gly Thr Asp Val Cys Tyr Pro Gly Lys Phe Thr Asn  
                   100                  105                  110  
 Glu Glu Ser Leu Arg Gln Ile Leu Arg Gly Ser Gly Gly Ile Asp Lys  
                   115                  120                  125  
 Glu Ser Met Gly Phe Thr Tyr Ser Gly Ile Arg Thr Asn Gly Ala Thr  
                   130                  135                  140

-continued

---

Ser	Ala	Cys	Arg	Arg	Ser	Gly	Ser	Ser	Phe	Tyr	Ala	Glu	Met	Lys	Trp
145					150					155					160
Leu	Leu	Ser	Asn	Ser	Asp	Asn	Ala	Ala	Phe	Pro	Gln	Met	Thr	Lys	Ser
			165						170					175	
Tyr	Arg	Asn	Pro	Arg	Asn	Lys	Pro	Ala	Leu	Ile	Ile	Trp	Gly	Val	His
		180						185					190		
His	Ser	Gly	Ser	Ala	Thr	Glu	Gln	Thr	Lys	Leu	Tyr	Gly	Ser	Gly	Asn
	195					200						205			
Lys	Leu	Ile	Thr	Val	Gly	Ser	Ser	Lys	Tyr	Gln	Gln	Ser	Phe	Thr	Pro
210					215					220					
Ser	Pro	Gly	Ala	Arg	Pro	Gln	Val	Asn	Gly	Gln	Ser	Gly	Arg	Ile	Asp
225					230				235					240	
Phe	His	Trp	Leu	Leu	Leu	Asp	Pro	Asn	Asp	Thr	Val	Thr	Phe	Thr	Phe
			245					250					255		
Asn	Gly	Ala	Phe	Ile	Ala	Pro	Asp	Arg	Ala	Ser	Phe	Phe	Arg	Gly	Glu
		260					265						270		
Ser	Leu	Gly	Val	Gln	Ser	Asp	Val	Pro	Leu	Asp	Ser	Gly	Cys	Glu	Gly
	275					280						285			
Asp	Cys	Phe	His	Ser	Gly	Gly	Thr	Ile	Val	Ser	Ser	Leu	Pro	Phe	Gln
290					295					300					
Asn	Ile	Asn	Pro	Arg	Thr	Val	Gly	Lys	Cys	Pro	Arg	Tyr	Val	Lys	Gln
305				310					315					320	
Thr	Ser	Leu	Leu	Leu	Ala	Thr	Gly	Met	Arg	Asn	Val	Pro	Glu	Asn	Pro
		325						330						335	
Lys	Gln	Ala	Tyr	Arg	Lys	Arg	Met	Thr	Arg	Gly	Leu	Phe	Gly	Ala	Ile
		340					345						350		
Ala	Gly	Phe	Ile	Glu	Asn	Gly	Trp	Glu	Gly	Leu	Ile	Asp	Gly	Trp	Tyr
	355					360						365			
Gly	Phe	Arg	His	Gln	Asn	Ala	Gln	Gly	Glu	Gly	Thr	Ala	Ala	Asp	Tyr
370					375						380				
Lys	Ser	Thr	Gln	Ser	Ala	Ile	Asp	Gln	Ile	Thr	Gly	Lys	Leu	Asn	Arg
385				390					395					400	
Leu	Ile	Asp	Lys	Thr	Asn	Gln	Gln	Phe	Glu	Leu	Ile	Asp	Asn	Glu	Phe
			405					410					415		
Ser	Glu	Ile	Glu	Gln	Gln	Ile	Gly	Asn	Val	Ile	Asn	Trp	Thr	Arg	Asp
	420						425						430		
Ser	Met	Thr	Glu	Val	Trp	Ser	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Ala	Met
	435					440					445				
Glu	Asn	Gln	His	Thr	Ile	Asp	Leu	Ala	Asp	Ser	Glu	Met	Asn	Lys	Leu
450					455						460				
Tyr	Glu	Arg	Val	Arg	Lys	Gln	Leu	Arg	Glu	Asn	Ala	Glu	Glu	Asp	Gly
465				470					475					480	
Thr	Gly	Cys	Phe	Glu	Ile	Phe	His	Lys	Cys	Asp	Asp	Gln	Cys	Met	Glu
		485						490					495		
Ser	Ile	Arg	Asn	Asn	Thr	Tyr	Asp	His	Thr	Gln	Tyr	Arg	Thr	Glu	Ser
	500						505						510		
Leu	Gln	Asn	Arg	Ile	Gln	Ile	Asp	Pro	Val	Lys	Leu	Ser	Ser	Gly	Tyr
	515					520					525				
Lys	Asp	Ile	Ile	Leu	Trp	Phe	Ser	Phe	Gly	Ala	Ser	Cys	Phe	Leu	Leu
530					535						540				

-continued

---

Leu Ala Ile Ala Met Gly Leu Val Phe Ile Cys Ile Lys Asn Gly Asn  
545 550 555 560

Met Arg Cys Thr Ile Cys Ile  
565

<210> SEQ ID NO 30

<211> LENGTH: 469

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 30

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Val Val Asn Thr Thr  
1 5 10 15

Leu Ser Thr Ile Ala Leu Leu Ile Gly Val Gly Asn Leu Val Phe Asn  
20 25 30

Thr Val Ile His Glu Lys Ile Gly Asp His Gln Ile Val Thr His Pro  
35 40 45

Thr Ile Met Thr Pro Glu Val Pro Asn Cys Ser Asp Thr Ile Ile Thr  
50 55 60

Tyr Asn Asn Thr Val Ile Asn Asn Ile Thr Thr Thr Ile Ile Thr Glu  
65 70 75 80

Ala Glu Arg Pro Phe Lys Ser Pro Leu Pro Leu Cys Pro Phe Arg Gly  
85 90 95

Phe Phe Pro Phe His Lys Asp Asn Ala Ile Arg Leu Gly Glu Asn Lys  
100 105 110

Asp Val Ile Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Asn Asp Asn  
115 120 125

Cys Trp Ser Phe Ala Leu Ala Gln Gly Ala Leu Leu Gly Thr Lys His  
130 135 140

Ser Asn Gly Thr Ile Lys Asp Arg Thr Pro Tyr Arg Ser Leu Ile Arg  
145 150 155 160

Phe Pro Ile Gly Thr Ala Pro Val Leu Gly Asn Tyr Lys Glu Ile Cys  
165 170 175

Ile Ala Trp Ser Ser Ser Ser Cys Phe Asp Gly Lys Glu Trp Met His  
180 185 190

Val Cys Met Thr Gly Asn Asp Asn Asp Ala Ser Ala Gln Ile Ile Tyr  
195 200 205

Gly Gly Arg Met Thr Asp Ser Ile Lys Ser Trp Arg Lys Asp Ile Leu  
210 215 220

Arg Thr Gln Glu Ser Glu Cys Gln Cys Ile Asp Gly Thr Cys Val Val  
225 230 235 240

Ala Val Thr Asp Gly Pro Ala Ala Asn Ser Ala Asp His Arg Val Tyr  
245 250 255

Trp Ile Arg Glu Gly Arg Ile Ile Lys Tyr Glu Asn Val Pro Lys Thr  
260 265 270

Lys Ile Gln His Leu Glu Glu Cys Ser Cys Tyr Val Asp Ile Asp Val  
275 280 285

Tyr Cys Ile Cys Arg Asp Asn Trp Lys Gly Ser Asn Arg Pro Trp Met  
290 295 300

Arg Ile Asn Asn Glu Thr Ile Leu Glu Thr Gly Tyr Val Cys Ser Lys  
305 310 315 320

Phe His Ser Asp Thr Pro Arg Pro Ala Asp Pro Ser Ile Met Ser Cys  
325 330 335

-continued

---

```

Asp Ser Pro Ser Asn Val Asn Gly Gly Pro Gly Val Lys Gly Phe Gly
    340                      345                      350

Phe Lys Ala Gly Asn Asp Val Trp Leu Gly Arg Thr Val Ser Thr Ser
    355                      360                      365

Gly Arg Ser Gly Phe Glu Ile Ile Lys Val Thr Glu Gly Trp Ile Asn
    370                      375                      380

Ser Pro Asn His Val Lys Ser Ile Thr Gln Thr Leu Val Ser Asn Asn
    385                      390                      395                      400

Asp Trp Ser Gly Tyr Ser Gly Ser Phe Ile Val Lys Ala Lys Asp Cys
    405                      410                      415

Phe Gln Pro Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg Pro Asn Lys
    420                      425                      430

Asn Asp Asp Val Ser Trp Thr Ser Asn Ser Ile Val Thr Phe Cys Gly
    435                      440                      445

Leu Asp Asn Glu Pro Gly Ser Gly Asn Trp Pro Asp Gly Ser Asn Ile
    450                      455                      460

Gly Phe Met Pro Lys
465

```

```

<210> SEQ ID NO 31
<211> LENGTH: 560
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 31

```

```

Met Asn Thr Gln Ile Leu Ala Phe Ile Ala Cys Met Leu Ile Gly Thr
1          5          10          15

Lys Gly Asp Lys Ile Cys Leu Gly His His Ala Val Ala Asn Gly Thr
20         25         30

Lys Val Asn Thr Leu Thr Glu Arg Gly Ile Glu Val Val Asn Ala Thr
35         40         45

Glu Thr Val Glu Thr Val Asn Ile Lys Lys Ile Cys Thr Gln Gly Lys
50         55         60

Arg Pro Thr Asp Leu Gly Gln Cys Gly Leu Leu Gly Thr Leu Ile Gly
65         70         75         80

Pro Pro Gln Cys Asp Gln Phe Leu Glu Phe Asp Ala Asn Leu Ile Ile
85         90         95

Glu Arg Arg Glu Gly Thr Asp Val Cys Tyr Pro Gly Lys Phe Thr Asn
100        105        110

Glu Glu Ser Leu Arg Gln Ile Leu Arg Gly Ser Gly Gly Ile Asp Lys
115        120        125

Glu Ser Met Gly Phe Thr Tyr Ser Gly Ile Arg Thr Asn Gly Ala Thr
130        135        140

Ser Ala Cys Arg Arg Ser Gly Ser Ser Phe Tyr Ala Glu Met Lys Trp
145        150        155        160

Leu Leu Ser Asn Ser Asp Asn Ala Ala Phe Pro Gln Met Thr Lys Ser
165        170        175

Tyr Arg Asn Pro Arg Asn Lys Pro Ala Leu Ile Ile Trp Gly Val His
180        185        190

His Ser Gly Ser Ala Thr Glu Gln Thr Lys Leu Tyr Gly Ser Gly Asn
195        200        205

Lys Leu Ile Thr Val Gly Ser Ser Lys Tyr Gln Gln Ser Phe Thr Pro

```



-continued

210	215	220
Ser Pro Gly Ala Arg	Pro Gln Val Asn Gly	Gln Ser Gly Arg Ile Asp
225	230	235 240
Phe His Trp Leu Leu	Leu Asp Pro Asn Asp	Thr Val Thr Phe Thr Phe
	245	250 255
Asn Gly Ala Phe Ile	Ala Pro Asp Arg Ala	Ser Phe Phe Arg Gly Glu
	260	265 270
Ser Leu Gly Val Gln	Ser Asp Val Pro Leu	Asp Ser Gly Cys Glu Gly
	275	280 285
Asp Cys Phe His Ser	Gly Gly Thr Ile Val	Ser Ser Leu Pro Phe Gln
	290	295 300
Asn Ile Asn Pro Arg	Thr Val Gly Lys Cys	Pro Arg Tyr Val Lys Gln
	305	310 315 320
Thr Ser Leu Leu Leu	Ala Thr Gly Met Arg	Asn Val Pro Glu Asn Pro
	325	330 335
Lys Thr Arg Gly Leu	Phe Gly Ala Ile Ala	Gly Phe Ile Glu Asn Gly
	340	345 350
Trp Glu Gly Leu Ile	Asp Gly Trp Tyr Gly	Phe Arg His Gln Asn Ala
	355	360 365
Gln Gly Glu Gly Thr	Ala Ala Asp Tyr Lys	Ser Thr Gln Ser Ala Ile
	370	375 380
Asp Gln Ile Thr Gly	Lys Leu Asn Arg Leu	Ile Asp Lys Thr Asn Gln
	385	390 395 400
Gln Phe Glu Leu Ile	Asp Asn Glu Phe Ser	Glu Ile Glu Gln Gln Ile
	405	410 415
Gly Asn Val Ile Asn	Trp Thr Arg Asp Ser	Met Thr Glu Val Trp Ser
	420	425 430
Tyr Asn Ala Glu Leu	Leu Val Ala Met Glu	Asn Gln His Thr Ile Asp
	435	440 445
Leu Ala Asp Ser Glu	Met Asn Lys Leu Tyr	Glu Arg Val Arg Lys Gln
	450	455 460
Leu Arg Glu Asn Ala	Glu Glu Asp Gly Thr	Gly Cys Phe Glu Ile Phe
	465	470 475 480
His Lys Cys Asp Asp	Gln Cys Met Glu Ser	Ile Arg Asn Asn Thr Tyr
	485	490 495
Asp His Thr Gln Tyr	Arg Thr Glu Ser Leu	Gln Asn Arg Ile Gln Ile
	500	505 510
Asp Pro Val Lys Leu	Ser Ser Gly Tyr Lys	Asp Ile Ile Leu Trp Phe
	515	520 525
Ser Phe Gly Ala Ser	Cys Phe Leu Leu Leu	Ala Ile Ala Met Gly Leu
	530	535 540
Val Phe Ile Cys Ile	Lys Asn Gly Asn Met	Arg Cys Thr Ile Cys Ile
	545	550 555 560

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 469

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 32

Met Asn Pro Asn Gln	Lys Ile Ile Thr	Ile Gly Val Val	Asn Thr Thr
1	5	10	15

Leu	Ser	Thr	Ile	Ala	Leu	Leu	Ile	Gly	Val	Gly	Asn	Leu	Val	Phe	Asn
			20				25				30				
Thr	Val	Ile	His	Glu	Lys	Ile	Gly	Asp	His	Gln	Ile	Val	Thr	His	Pro
			35				40				45				
Thr	Ile	Met	Thr	Pro	Glu	Val	Pro	Asn	Cys	Ser	Asp	Thr	Ile	Ile	Thr
			50				55				60				
Tyr	Asn	Asn	Thr	Val	Ile	Asn	Asn	Ile	Thr	Thr	Thr	Ile	Ile	Thr	Glu
			65				70				75				
Ala	Glu	Arg	Pro	Phe	Lys	Ser	Pro	Leu	Pro	Leu	Cys	Pro	Phe	Arg	Gly
			85				90				95				
Phe	Phe	Pro	Phe	His	Lys	Asp	Asn	Ala	Ile	Arg	Leu	Gly	Glu	Asn	Lys
			100				105				110				
Asp	Val	Ile	Val	Thr	Arg	Glu	Pro	Tyr	Val	Ser	Cys	Asp	Asn	Asp	Asn
			115				120				125				
Cys	Trp	Ser	Phe	Ala	Leu	Ala	Gln	Gly	Ala	Leu	Leu	Gly	Thr	Lys	His
			130				135				140				
Ser	Asn	Gly	Thr	Ile	Lys	Asp	Arg	Thr	Pro	Tyr	Arg	Ser	Leu	Ile	Arg
			145				150				155				
Phe	Pro	Ile	Gly	Thr	Ala	Pro	Val	Leu	Gly	Asn	Tyr	Lys	Glu	Ile	Cys
			165				170				175				
Ile	Ala	Trp	Ser	Ser	Ser	Ser	Cys	Phe	Asp	Gly	Lys	Glu	Trp	Met	His
			180				185				190				
Val	Cys	Met	Thr	Gly	Asn	Asp	Asn	Asp	Ala	Ser	Ala	Gln	Ile	Ile	Tyr
			195				200				205				
Gly	Gly	Arg	Met	Thr	Asp	Ser	Ile	Lys	Ser	Trp	Arg	Lys	Asp	Ile	Leu
			210				215				220				
Arg	Thr	Gln	Glu	Ser	Glu	Cys	Gln	Cys	Ile	Asp	Gly	Thr	Cys	Val	Val
			225				230				235				
Ala	Val	Thr	Asp	Gly	Pro	Ala	Ala	Asn	Ser	Ala	Asp	His	Arg	Val	Tyr
			245				250				255				
Trp	Ile	Arg	Glu	Gly	Arg	Ile	Ile	Lys	Tyr	Glu	Asn	Val	Pro	Lys	Thr
			260				265				270				
Lys	Ile	Gln	His	Leu	Glu	Glu	Cys	Ser	Cys	Tyr	Val	Asp	Ile	Asp	Val
			275				280				285				
Tyr	Cys	Ile	Cys	Arg	Asp	Asn	Trp	Lys	Gly	Ser	Asn	Arg	Pro	Trp	Met
			290				295				300				
Arg	Ile	Asn	Asn	Glu	Thr	Ile	Leu	Glu	Thr	Gly	Tyr	Val	Cys	Ser	Lys
			305				310				315				
Phe	His	Ser	Asp	Thr	Pro	Arg	Pro	Ala	Asp	Pro	Ser	Ile	Met	Ser	Cys
			325				330				335				
Asp	Ser	Pro	Ser	Asn	Val	Asn	Gly	Gly	Pro	Gly	Val	Lys	Gly	Phe	Gly
			340				345				350				
Phe	Lys	Ala	Gly	Asn	Asp	Val	Trp	Leu	Gly	Arg	Thr	Val	Ser	Thr	Ser
			355				360				365				
Gly	Arg	Ser	Gly	Phe	Glu	Ile	Ile	Lys	Val	Thr	Glu	Gly	Trp	Ile	Asn
			370				375				380				
Ser	Pro	Asn	His	Val	Lys	Ser	Ile	Thr	Gln	Thr	Leu	Val	Ser	Asn	Asn
			385				390				395				
Asp	Trp	Ser	Gly	Tyr	Ser	Gly	Ser	Phe	Ile	Val	Lys	Ala	Lys	Asp	Cys
			405				410				415				
Phe	Gln	Pro	Cys	Phe	Tyr	Val	Glu	Leu	Ile	Arg	Gly	Arg	Pro	Asn	Lys

-continued

420	425	430	
Asn Asp Asp Val Ser Trp Thr Ser Asn Ser Ile Val Thr Phe Cys Gly			
435	440	445	
Leu Asp Asn Glu Pro Gly Ser Gly Asn Trp Pro Asp Gly Ser Asn Ile			
450	455	460	
Gly Phe Met Pro Lys			
465			
 <210> SEQ ID NO 33			
<211> LENGTH: 1743			
<212> TYPE: DNA			
<213> ORGANISM: Influenza A virus			
 <400> SEQUENCE: 33			
agcaaaagca ggggaaaatg attgcagtca ttataatagc ggtactggca acggccggaa	60		
aatcagacaa gatctgcatt gggatatcatg ccaacaattc aacaacacaa gtggatacga	120		
tacttgagaa gaatgtaacc gtcacacact cagttgaatt gctggagaac caaaaagaag	180		
aaagattctg caagatcttg aacaaggccc ctctcgattt aagaggatgt accatagagg	240		
gttggatctt ggggaatccc caatgcgacc tattgcttgg tgatcaaagc tggtcataata	300		
tagtggaag acctacagct caaaatggga tctgctaccc aggaattttg aatgaagtag	360		
aaagaactgaa ggcaacttatt ggatcaggag aaagagtgga gagatttgaa atgtttccca	420		
aaagtacatg ggcaggagta gacaccagca gtggggtaac aaaggcttgc ccttatacta	480		
gtggttcgtc tttctacaga aacctctat ggataataaa aaccaagtcc gcagcatatc	540		
cagtaattaa gggaacctac aataacactg gaagccagcc aatcctctat ttctggggtg	600		
tgcaccatcc tcttgacacc aatgagcaaa acactttgta tggctctggt gatcgatatg	660		
tcaggatggg aactgaaagc atgaattttg ccaagagccc agaaattgcg gcaaggcctg	720		
ctgtgaatgg tcaaagaggc agaattgatt attactggtc tgttttaag ccgggggaaa	780		
ccttgaatgt ggaatctaataa ggaatctaa tgcgcccttg gtatgcatac aaatttgtca	840		
gcaccaatag taaaggagcc gtcttcaagt caaatttacc aatcgagaac tgtgatgcca	900		
catgccagac tattgcagga gtcttaagaa ccaataaaac atttcagaat gtaagccctc	960		
tgtggatagg agaatgcccc aaatatgtga aaagtgaag tttgaggctt gcaactggac	1020		
taagaaatat tccacagatt gagactagag gacttttcgg agctatcgca gggtttattg	1080		
aaggaggatg gactggaatg atagatgggt ggtatggcta tcaccatgaa aattctcaag	1140		
gctcagggta tgcggcagac agagaaagca ctcaaagggc tatagacgga attacaata	1200		
agggtcaattc cattatagac aaaatgaaca cacaattcga agctatagac cacgaattct	1260		
caaatattga gagaagaatt gacagtctga acaaaagaat ggaagatgga tttctggacg	1320		
tttgacata caatgctgaa ctgttggttc ttcttgaaaa cgaaaggaca ctgacacctac	1380		
atgacgcgaa tgtgaagaac ctgtatgaaa aggtcaaatc acaactacgg gacaatgcta	1440		
atgatctagg aaatggatgc tttgaatttt ggcataagtg tgacaatgaa tgcatagagt	1500		
ctgtcaaaaa tggtaacctat gactatccca aatatcagga tgaaagcaaa ttgaacaggc	1560		
aggaaataga atcgggtgaag ctggagaacc ttggtgtgta tcaaatctc gccatttata	1620		
gtacggatc gagcagtcta gtcttggtgagg ggtgattat agcaatgggt ctttgatgt	1680		
gttcaaatgg ttcaatgcaa tgcaggatat gtatataatt aagaaaaaca cccttgttct	1740		

---

-continued

---

act 1743

<210> SEQ ID NO 34  
<211> LENGTH: 1460  
<212> TYPE: DNA  
<213> ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 34

agcaaaagca gggtaagat gaatccaaat cagaagattc tatgcacatc tgctactgcc 60  
attgcaatag gcacaattgc tgtattaata ggaatagcaa acctggggtt gaacatagga 120  
ctacacctga aaccgagctg caactgctcc aacctctctc ctgaaacaac aaatgtaagc 180  
caacaataa taaacaatta ctacaatgaa acaaatgtta cccaaataag taacacaaac 240  
attcaacata tggggggaac cgaaaaggac ttcaacaatc tgactaaagg gctctgcaca 300  
ataaattcat ggcataatatt cggaaggac aatgctataa gaatagggga gaactctgat 360  
gttttagtca caagagagcc atatgtttct tgtgatccag atgaatgcag attctatgct 420  
ctcagccaag gaacaacaat acggggaaag cactcaaatg gaacaatata cgatagatcc 480  
caataccgtg cttagtgag ctggccttta tcatcaccac ccactgtgta caataccaga 540  
gtagaatgca ttggatggtc cagtacaagc tgccatgatg ggaaagcacg aatgtctata 600  
tgtgtctcag gtcccaacaa caatgcata gcagtgattt ggtacaaagg gcggcctatc 660  
acggaaatca atacgtgggc ccgaacata ttgagaaccc aagaatctga gtgtgtatgc 720  
cacaatggaa tatgtccagt agtgttctact gacggttctg ccaccggctc agcagaaact 780  
aggatatact atttcaaaga ggggaaaatc ctcaaattgg agccactaac tggaaccgcc 840  
aagcacattg aagaatgctc ttgctatggg aaagactcag aaataacgtg cacatgtaga 900  
gacaattggc aaggctcgaa tagaccagta atacaataa accccacaat gatgactcac 960  
actagtcaat acatagtcag ccctgtctc acagacaatc cagcccccaa tgacccacg 1020  
gtaggcaagt gtaatgatcc ttatccagga aacaacaata atggagtcaa aggattctca 1080  
tatttagatg gtgacaatac atggctagga agaacgataa gcacagcctc taggtctggg 1140  
tatgaaatgc tgaaagtgc taatgcattg acagatgata gatcaaaacc tactcaaggt 1200  
cagacaattg tattaacac agactggagt gggtacagtg ggtcttctat tgattactgg 1260  
gcaaaagggg agtgctatag agcatgcttc tacgttgagc tgatccgtgg aaggccaaaa 1320  
gaggacaaag tgtgtgggac cagtaatagt atagtgtcga tgtgttcag cacagagttc 1380  
cttgacaat ggaactggcc agatggggct aaaatagagt acttcctcta agatgtagaa 1440  
aaaagaccct tgtttctact 1460

<210> SEQ ID NO 35  
<211> LENGTH: 1747  
<212> TYPE: DNA  
<213> ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 35

agcaaaagca ggggaaatg attgcaatca ttgtaatagc aatactggca gcagccggaa 60  
aatcagacaa gatctgcatt gggatatcat ccaacaattc aacaacacag gtagatacga 120  
tacttgagaa gaatgtgact gtcacacact caattgaatt gctggaaaat cagaaggaag 180  
aaagattctg caagatattg aacaaggccc ctctcgactt aagggaatgt accatagagg 240

-continued

---

```

gttggatcctt ggggaatccc caatgcgacc tattgcttgg tgatcaaagc tggtcataca    300
ttgtggaaaag acctactgct caaaacggga tctgctaccc aggaacctta aatgaggtag    360
aagaactgag ggcacttatt ggatcaggag aaagggtaga gagatttgag atgtttcccc    420
aaagcacctg gcaaggagtt gacaccaaca gtggaacaac aagatcctgc ccttattcta    480
ctggtgatcc gtctttctac agaaaacctcc tatggataat aaaaaccaag acagcagaat    540
atccagtaat taaggaattt tacaacaaca ctggaacca gccaatcctc tatttctggg    600
gtgtgcatca tcctcctaac accgacgagc aagatactct gtatggctct ggtgatcgat    660
acgttagaat ggggaactgaa agcatgaatt ttgccaagag tccggaaatt gcggcaaggc    720
ctgctgtgaa tggacaaaga ggcagaattg attattattg gtcggtttta aaaccagggg    780
aaaccttgaa tgtggaatct aatggaaatc taatcgcccc ttggtatgca taaaaatttg    840
tcaacacaaa tagtaaagga gccgtcttca ggtcagattt accaatcgag aactgcgatg    900
ccacatgcca gactattgca ggggttctaa ggaccaataa aacatttcag aatgtgagtc    960
ccctgtggat aggagaatgt cccaaatacg tgaagagtgaa aagtctgagg cttgcaactg   1020
gactaagaaa tgttcacagc attgaaacta gaggactctt cggagctatt gcagggttta   1080
ttgaaggagg atggactggg atgatagatg ggtgggatgg ctatcaccat gaaaattctc   1140
aagggtcagg atatgcagcg gacagagaaa gcactcaaaa ggctgtaaac agaattacaa   1200
ataaggtcaa ttccatcatc aacaaaatga acacacaatt tgaagctgtc gatcacgaat   1260
tttcaaatct ggagaggaga atcgacaatc tgaacaaaag aatgcaagat ggatttcttg   1320
atgtttggac atacaatgct gaactgttgg ttcttcttga aaacgaaaga aactagaca   1380
tgcattgacg aaatgtgaag aacctacatg aaaaggtcaa atcacaaacta agggacaatg   1440
ctaacgatct agggaatggt tgctttgaat tttggcataa gtgtgacaat gaatgcatag   1500
agtctgtcaa aaatggtaca tatgactatc ccaaatacca gactgaaagc aaattaaaca   1560
ggctaaaaat agaatcagta aagctagaga accttggtgt gtatcaaatt cttgccattt   1620
atagtacggt atcgagcagc ctagtgttgg tagggctgat catggcaatg ggtctttgga   1680
tgtgttcaaa tggttcaatg cagtgcattg tgtgtatatg attaagaaaa acacccttgt   1740
ttctact                                           1747

```

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 1401

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 36

```

agcaaaaagca ggagttaaac atgaatccaa atcagaagat aataaccatt gggtaaatct    60
gtatggtagt tggaaataac agcttgatgt taaaaattgg aaacataata tcaatatggg    120
ttagccacat aattcagact gggcatccaa accagcctgg gccatgcaat caaagcatca    180
atttttacac tgagcaggct gcagcttcag tgacattagc gggtaattcc tctctctgcc    240
ctattagtgg atgggtcata tacagtaaaag acaatagtat aagaattggt tccaaagggg    300
atgtgtttgt tatgagagaa ccattcgttt catgctccca tttggaatgc agaacctttt    360
tcttgactca aggagcccta ttgaatgaca agcattctaa tgggaccgtt aaagacagaa    420
gccctatag aactttaatg agctgtcctg ttggtgaggc tccttcccca tacaactcaa    480

```

-continued

---

```

ggtttgagtc tgttgcttgg tcagcaagtg cttgccatga tggcattagt tggctaacaa 540
ttggaatttc cgggccggat aatggggctg tggctgtgtt gaaatacaat ggcataataa 600
cagacaccat caagagttag aggaacaaca tactgaggac acaagagtct gaatgtgcat 660
gtgtgaatgg ttcttgtttt actgtaatga cagatggacc gagtaatgaa caggcctcat 720
acaagatttt caagatagag aaggggaaa tagtcaaac agttgagttg aacgccccta 780
attatcatta cgaggaatgc tcctgttata ctgatgctgg cgaaatcaca tgtgtgtgca 840
gggataattg gcatggctcg aaccgacctg ggggtgtctt caatcagaat ctggagtatc 900
aaataggata tatatgcagt ggggttttcg gagacagtcc acgcccacat gatggaacag 960
gcagttgctg tccagtgtct cttaacggag agtatggagt aaaaggggtt tcatttaagt 1020
acggtgatgg tgtttggatc gggagaacca aaagcactag ttccaggagc gggtttgaaa 1080
tgatttggga tccaaatggg tggaccgaaa cagatagtaa cttctcattg aagcaagaca 1140
tcatagcaat aactgattgg tcaggataca gcgggagttt tgtccaacat ccagaactga 1200
caggattaaa ttgcatgagg ccttgcttct ggggtgaact aatcagaggg aggcccaaag 1260
agaaaacaat ctggactagt gggagcagta tatctttctg tgggtgaaat agtgacactg 1320
tgggttggtc ttggccagac ggtgctgagg tgccattcac cattgacaag tagtttggtc 1380
aaaaaactcc ttgtttctac t 1401

```

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 1745

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 37

```

agcaaaagca ggggaaaatg attgcaatca taatacttgc aatagtggtc tctaccagca 60
agtcagacag gatctgcatt ggttaccatg caaacaactc gacaacacaa gtggacacaa 120
tattagagaa gaatgtgaca gtgacacact cagtggagct cctagaaaac cagaaggaga 180
atagattctg cagagtcttg aataaagcgc cactggatct aatggactgc accactgagg 240
gttggatcct tggaaacccc cgatgtgata acttactcgg tgatcaaagt tggtcataca 300
tagtagagag gcctgatgcc caaaatggga tatgttacc aggggtattg aaggagacgg 360
aagagctgaa agcactcatt gggctctatg atagcataca aagatttgaa atgtttccca 420
agagcacgtg gaccggggta gatactaata gcggagttac gagcgcttgc cctacaatg 480
gtgaatcttc cttttacagg aatctgttgt ggataataaa aataagatct gatccgtact 540
cattgatcaa ggggacatat accaatacag gctctcagcc aatcttatat ttctggggtg 600
tgcaccatcc tccagatgaa gttgagcaag ctaacttgta tggaaattgg acccggtatg 660
ttaggatggg aactgaaagt atgaattttg ccaaaggctc tgaatatgca ggcagaccac 720
ctgcgaatgg gcaacaggga agaattgatt attattggtc tgtgttgaag ccaggagaaa 780
ccttgaatgt ggaatccaat ggaaatttaa tagctccttg gtatgcttac aagttcacta 840
gttcagaaaa caaggagct attttcaaat cagaccttcc aattgagaat tgtgatgctg 900
tctgtcaaac tttagctgga gcaataaata caaacaaaac cttccaaaat attagtccag 960
tctggattgg agaatgcccc aaatatgtta aaagtaagag cctaaaaacta gcaactggtc 1020
tgagaaatgt tccacaggca gaaacaagag gattgttttg agcaatagct gggtttatag 1080

```

-continued

---

```

aaggaggatg gacaggtatg gtagacggat ggtacggata ccaccatgaa aattcacagg 1140
ggtctgggta tgcagcagat aaagaaagca ctacagaaagc aatagacggg atcaccaata 1200
aagtcaattc aatcattgac aaaatgaaca cacaatttga ggcagtagag catgagttct 1260
caagtctcga aaggagaata ggcaatctga acaaaagaat ggaagatgga tttttagacg 1320
tgtggacata caatgctgaa cttctgggtc tactggaaaa tgagaggact ttggacatgc 1380
atgatgctaa tgtaagaat ctacatgaaa aggtgaaatc acaattaagg gataatgcaa 1440
aggatttggg taatgggtgt tttgaatttt ggcacaaatg cgacaatgaa tgcatacaact 1500
cagttaaaaa tggcacatat gactacccaa agtaccagga agagagcaga cttaaataggc 1560
aggaaataaa atcagtgatg ctggaaaatc tgggagtata ccaaatcctt gctatttata 1620
gtacggtatc gagcagtcgt gttttgggtg gactgatcat tgccatgggt ctttggatgt 1680
gctcaaatgg ctcaatgcaa tgcaagatat gtatataatt agaaaaaac acccttgttt 1740
ctact 1745

```

```

<210> SEQ ID NO 38
<211> LENGTH: 1467
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 38

```

```

agcaaaaagca ggagtgaata tgaatccaaa tcagaggata ataacaattg gatccgtctc 60
tctaactatt gcaacagtggt gtttcctcat gcagattgcc atcctagcaa cgactgtgac 120
actgcatttc aaacaaaatg aatgcagcat tcccgcaaac aaccaagtaa cgccatgtga 180
accaatagta atagagagga acataacaga gatagtgtat ttgaataata ctaccataga 240
aaaagagatt tgtcctgaag tagtagaata caggaattgg tcaaaaccgc aatgtcaaat 300
tacagggttt gtcctttctt ccaaggacaa ctcaattcgg ctttctgctg gtggggacat 360
ttggataaca agagaacctt atgtgtcatg cgaccccgat aatgtttatc aatttgcact 420
cgggcagggg accacgctgg acaacaaaca ctcaaatggc acaatacatg atagaatccc 480
tcacgagacc cttttgatga atgaattggg tgtccggtt catttgggaa ccaacaagt 540
gtgcatagca tgggtccagc caagctgtca tgatgggaaa gcatggttgc acgtttgtgt 600
cactggggat gatagaaatg caactgctag tttcatttat gatgggatgc ttattgacag 660
tattggttcc tgggtctcaa atatcctcag gactcaggag tcagaatgcg tttgtatcag 720
tggaacttgt acagtagtaa tgactgatg aagtgcacga ggaagggcag aactagaat 780
actattcatt agagagggga aaattgtcca cattagtcca ttgtcaggaa gtgctcagca 840
ttagagaggaa tgttcttgtt atccccggta cccaaacgtc agatgtgtct gcagagacaa 900
ctggaagggc tctaataaggc ccgttataga tataaatatg gcagattata gcattgactc 960
aagttatgtg tgctcaggac ttgttgaga cacaccaagg aacgatgata gctctagcag 1020
cagcaactgc agggatccta ataatgagag agggaaccca ggagtgaag ggtgggcctt 1080
tgataatgga aatgatgtgt ggatgggaag aacaatcagt aaagattcgc gctcaggcta 1140
tgagaccttc aaggtcattg gtggttgggc cattgctaata tccaagtcac agaccaatag 1200
acaagtcata gttgataata acaactggtc tggttattct ggtattttct ctgttgaaag 1260
caaaggctgc atcaataggt gtttttatgt ggagttgata agagggaaggc cacaggagac 1320

```

-continued

---

```

tagagtatgg tggacctcaa acagtattgt cgtatattgt ggcacttcag ggacatatgg 1380
aacaggctca tggcctgatg gggcgaatat cgatttcacg cctatataag ctttcgcaat 1440
tttagaaaaa aactccttgt ttctact 1467

```

```

<210> SEQ ID NO 39
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 39

```

```

Met  Ile  Ala  Val  Ile  Ile  Ile  Ala  Val  Leu  Ala  Thr  Ala  Gly  Lys  Ser
 1          5          10          15
Asp  Lys  Ile  Cys  Ile  Gly  Tyr  His  Ala  Asn  Asn  Ser  Thr  Thr  Gln  Val
          20          25          30
Asp  Thr  Ile  Leu  Glu  Lys  Asn  Val  Thr  Val  Thr  His  Ser  Val  Glu  Leu
          35          40          45
Leu  Glu  Asn  Gln  Lys  Glu  Glu  Arg  Phe  Cys  Lys  Ile  Leu  Asn  Lys  Ala
          50          55          60
Pro  Leu  Asp  Leu  Arg  Gly  Cys  Thr  Ile  Glu  Gly  Trp  Ile  Leu  Gly  Asn
        65          70          75          80
Pro  Gln  Cys  Asp  Leu  Leu  Leu  Gly  Asp  Gln  Ser  Trp  Ser  Tyr  Ile  Val
          85          90          95
Glu  Arg  Pro  Thr  Ala  Gln  Asn  Gly  Ile  Cys  Tyr  Pro  Gly  Ile  Leu  Asn
          100         105         110
Glu  Val  Glu  Glu  Leu  Lys  Ala  Leu  Ile  Gly  Ser  Gly  Glu  Arg  Val  Glu
          115         120         125
Arg  Phe  Glu  Met  Phe  Pro  Lys  Ser  Thr  Trp  Ala  Gly  Val  Asp  Thr  Ser
          130         135         140
Ser  Gly  Val  Thr  Lys  Ala  Cys  Pro  Tyr  Thr  Ser  Gly  Ser  Ser  Phe  Tyr
          145         150         155         160
Arg  Asn  Leu  Leu  Trp  Ile  Ile  Lys  Thr  Lys  Ser  Ala  Ala  Tyr  Pro  Val
          165         170         175
Ile  Lys  Gly  Thr  Tyr  Asn  Asn  Thr  Gly  Ser  Gln  Pro  Ile  Leu  Tyr  Phe
          180         185         190
Trp  Gly  Val  His  His  Pro  Pro  Asp  Thr  Asn  Glu  Gln  Asn  Thr  Leu  Tyr
          195         200         205
Gly  Ser  Gly  Asp  Arg  Tyr  Val  Arg  Met  Gly  Thr  Glu  Ser  Met  Asn  Phe
          210         215         220
Ala  Lys  Ser  Pro  Glu  Ile  Ala  Ala  Arg  Pro  Ala  Val  Asn  Gly  Gln  Arg
          225         230         235         240
Gly  Arg  Ile  Asp  Tyr  Tyr  Trp  Ser  Val  Leu  Lys  Pro  Gly  Glu  Thr  Leu
          245         250         255
Asn  Val  Glu  Ser  Asn  Gly  Asn  Leu  Ile  Ala  Pro  Trp  Tyr  Ala  Tyr  Lys
          260         265         270
Phe  Val  Ser  Thr  Asn  Ser  Lys  Gly  Ala  Val  Phe  Lys  Ser  Asn  Leu  Pro
          275         280         285
Ile  Glu  Asn  Cys  Asp  Ala  Thr  Cys  Gln  Thr  Ile  Ala  Gly  Val  Leu  Arg
          290         295         300
Thr  Asn  Lys  Thr  Phe  Gln  Asn  Val  Ser  Pro  Leu  Trp  Ile  Gly  Glu  Cys
          305         310         315         320
Pro  Lys  Tyr  Val  Lys  Ser  Glu  Ser  Leu  Arg  Leu  Ala  Thr  Gly  Leu  Arg

```



-continued

325					330					335					
Asn	Ile	Pro	Gln	Ile	Glu	Thr	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly
			340					345					350		
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr
		355					360					365			
His	His	Glu	Asn	Ser	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Arg	Glu	Ser
	370					375					380				
Thr	Gln	Arg	Ala	Ile	Asp	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Ile	Ile
385					390					395					400
Asp	Lys	Met	Asn	Thr	Gln	Phe	Glu	Ala	Ile	Asp	His	Glu	Phe	Ser	Asn
			405						410					415	
Leu	Glu	Arg	Arg	Ile	Asp	Ser	Leu	Asn	Lys	Arg	Met	Glu	Asp	Gly	Phe
		420						425					430		
Leu	Asp	Val	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
	435						440					445			
Glu	Arg	Thr	Leu	Asp	Leu	His	Asp	Ala	Asn	Val	Lys	Asn	Leu	Tyr	Glu
	450					455					460				
Lys	Val	Lys	Ser	Gln	Leu	Arg	Asp	Asn	Ala	Asn	Asp	Leu	Gly	Asn	Gly
465					470					475					480
Cys	Phe	Glu	Phe	Trp	His	Lys	Cys	Asp	Asn	Glu	Cys	Ile	Glu	Ser	Val
			485					490						495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Gln	Asp	Glu	Ser	Lys	Leu
		500						505					510		
Asn	Arg	Gln	Glu	Ile	Glu	Ser	Val	Lys	Leu	Glu	Asn	Leu	Gly	Val	Tyr
		515					520					525			
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ser	Ser	Ser	Leu	Val	Leu	Val
	530					535					540				
Gly	Leu	Ile	Ile	Ala	Met	Gly	Leu	Trp	Met	Cys	Ser	Asn	Gly	Ser	Met
545					550				555						560
Gln	Cys	Arg	Ile	Cys	Ile										
			565												

&lt;210&gt; SEQ ID NO 40

&lt;211&gt; LENGTH: 470

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 40

Met	Asn	Pro	Asn	Gln	Lys	Ile	Leu	Cys	Thr	Ser	Ala	Thr	Ala	Ile	Ala
1			5						10					15	
Ile	Gly	Thr	Ile	Ala	Val	Leu	Ile	Gly	Ile	Ala	Asn	Leu	Gly	Leu	Asn
		20					25					30			
Ile	Gly	Leu	His	Leu	Lys	Pro	Ser	Cys	Asn	Cys	Ser	Asn	Pro	Pro	Pro
	35					40					45				
Glu	Thr	Thr	Asn	Val	Ser	Gln	Thr	Ile	Ile	Asn	Asn	Tyr	Tyr	Asn	Glu
	50					55				60					
Thr	Asn	Val	Thr	Gln	Ile	Ser	Asn	Thr	Asn	Ile	Gln	His	Met	Gly	Gly
65				70					75					80	
Thr	Glu	Lys	Asp	Phe	Asn	Asn	Leu	Thr	Lys	Gly	Leu	Cys	Thr	Ile	Asn
		85						90					95		
Ser	Trp	His	Ile	Phe	Gly	Lys	Asp	Asn	Ala	Ile	Arg	Ile	Gly	Glu	Asn
		100					105						110		

-continued

---

```

Ser Asp Val Leu Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Asp
    115                      120                      125

Glu Cys Arg Phe Tyr Ala Leu Ser Gln Gly Thr Thr Ile Arg Gly Lys
    130                      135                      140

His Ser Asn Gly Thr Ile His Asp Arg Ser Gln Tyr Arg Ala Leu Val
    145                      150                      155                      160

Ser Trp Pro Leu Ser Ser Pro Pro Thr Val Tyr Asn Thr Arg Val Glu
    165                      170                      175

Cys Ile Gly Trp Ser Ser Thr Ser Cys His Asp Gly Lys Ala Arg Met
    180                      185                      190

Ser Ile Cys Val Ser Gly Pro Asn Asn Asn Ala Ser Ala Val Ile Trp
    195                      200                      205

Tyr Lys Gly Arg Pro Ile Thr Glu Ile Asn Thr Trp Ala Arg Asn Ile
    210                      215                      220

Leu Arg Thr Gln Glu Ser Glu Cys Val Cys His Asn Gly Ile Cys Pro
    225                      230                      235                      240

Val Val Phe Thr Asp Gly Ser Ala Thr Gly Pro Ala Glu Thr Arg Ile
    245                      250                      255

Tyr Tyr Phe Lys Glu Gly Lys Ile Leu Lys Trp Glu Pro Leu Thr Gly
    260                      265                      270

Thr Ala Lys His Ile Glu Glu Cys Ser Cys Tyr Gly Lys Asp Ser Glu
    275                      280                      285

Ile Thr Cys Thr Cys Arg Asp Asn Trp Gln Gly Ser Asn Arg Pro Val
    290                      295                      300

Ile Gln Ile Asn Pro Thr Met Met Thr His Thr Ser Gln Tyr Ile Cys
    305                      310                      315                      320

Ser Pro Val Leu Thr Asp Asn Pro Arg Pro Asn Asp Pro Thr Val Gly
    325                      330                      335

Lys Cys Asn Asp Pro Tyr Pro Gly Asn Asn Asn Asn Gly Val Lys Gly
    340                      345                      350

Phe Ser Tyr Leu Asp Gly Asp Asn Thr Trp Leu Gly Arg Thr Ile Ser
    355                      360                      365

Thr Ala Ser Arg Ser Gly Tyr Glu Met Leu Lys Val Pro Asn Ala Leu
    370                      375                      380

Thr Asp Asp Arg Ser Lys Pro Thr Gln Gly Gln Thr Ile Val Leu Asn
    385                      390                      395                      400

Thr Asp Trp Ser Gly Tyr Ser Gly Ser Phe Ile Asp Tyr Trp Ala Lys
    405                      410                      415

Gly Glu Cys Tyr Arg Ala Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg
    420                      425                      430

Pro Lys Glu Asp Lys Val Trp Trp Thr Ser Asn Ser Ile Val Ser Met
    435                      440                      445

Cys Ser Ser Thr Glu Phe Leu Gly Gln Trp Asn Trp Pro Asp Gly Ala
    450                      455                      460

Lys Ile Glu Tyr Phe Leu
    465                      470

```

&lt;210&gt; SEQ ID NO 41

&lt;211&gt; LENGTH: 567

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 41

-continued

---

Met	Ile	Ala	Ile	Ile	Val	Ile	Ala	Ile	Leu	Ala	Ala	Ala	Gly	Lys	Ser
1			5						10					15	
Asp	Lys	Ile	Cys	Ile	Gly	Tyr	His	Ala	Asn	Asn	Ser	Thr	Thr	Gln	Val
	20						25						30		
Asp	Thr	Ile	Leu	Glu	Lys	Asn	Val	Thr	Val	Thr	His	Ser	Ile	Glu	Leu
	35					40					45				
Leu	Glu	Asn	Gln	Lys	Glu	Glu	Arg	Phe	Cys	Lys	Ile	Leu	Asn	Lys	Ala
	50				55					60					
Pro	Leu	Asp	Leu	Arg	Glu	Cys	Thr	Ile	Glu	Gly	Trp	Ile	Leu	Gly	Asn
65				70					75					80	
Pro	Gln	Cys	Asp	Leu	Leu	Leu	Gly	Asp	Gln	Ser	Trp	Ser	Tyr	Ile	Val
		85					90						95		
Glu	Arg	Pro	Thr	Ala	Gln	Asn	Gly	Ile	Cys	Tyr	Pro	Gly	Thr	Leu	Asn
	100					105						110			
Glu	Val	Glu	Glu	Leu	Arg	Ala	Leu	Ile	Gly	Ser	Gly	Glu	Arg	Val	Glu
	115				120						125				
Arg	Phe	Glu	Met	Phe	Pro	Gln	Ser	Thr	Trp	Gln	Gly	Val	Asp	Thr	Asn
	130				135					140					
Ser	Gly	Thr	Thr	Arg	Ser	Cys	Pro	Tyr	Ser	Thr	Gly	Asp	Pro	Ser	Phe
145				150					155					160	
Tyr	Arg	Asn	Leu	Leu	Trp	Ile	Ile	Lys	Thr	Lys	Thr	Ala	Glu	Tyr	Pro
		165						170					175		
Val	Ile	Lys	Gly	Ile	Tyr	Asn	Asn	Thr	Gly	Thr	Gln	Pro	Ile	Leu	Tyr
	180					185						190			
Phe	Trp	Gly	Val	His	His	Pro	Pro	Asn	Thr	Asp	Glu	Gln	Asp	Thr	Leu
	195					200					205				
Tyr	Gly	Ser	Gly	Asp	Arg	Tyr	Val	Arg	Met	Gly	Thr	Glu	Ser	Met	Asn
	210				215					220					
Phe	Ala	Lys	Ser	Pro	Glu	Ile	Ala	Ala	Arg	Pro	Ala	Val	Asn	Gly	Gln
225				230					235					240	
Arg	Gly	Arg	Ile	Asp	Tyr	Tyr	Trp	Ser	Val	Leu	Lys	Pro	Gly	Glu	Thr
		245						250					255		
Leu	Asn	Val	Glu	Ser	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Tyr
	260					265						270			
Lys	Phe	Val	Asn	Thr	Asn	Ser	Lys	Gly	Ala	Val	Phe	Arg	Ser	Asp	Leu
	275					280				285					
Pro	Ile	Glu	Asn	Cys	Asp	Ala	Thr	Cys	Gln	Thr	Ile	Ala	Gly	Val	Leu
	290				295					300					
Arg	Thr	Asn	Lys	Thr	Phe	Gln	Asn	Val	Ser	Pro	Leu	Trp	Ile	Gly	Glu
305				310					315					320	
Cys	Pro	Lys	Tyr	Val	Lys	Ser	Glu	Ser	Leu	Arg	Leu	Ala	Thr	Gly	Leu
		325						330					335		
Arg	Asn	Val	Pro	Gln	Ile	Glu	Thr	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala
	340					345						350			
Gly	Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly
	355					360					365				
Tyr	His	His	Glu	Asn	Ser	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Arg	Glu
	370				375					380					
Ser	Thr	Gln	Lys	Ala	Val	Asn	Arg	Ile	Thr	Asn	Lys	Val	Asn	Ser	Ile
385				390					395					400	

-continued

---

```

Ile Asn Lys Met Asn Thr Gln Phe Glu Ala Val Asp His Glu Phe Ser
      405                      410                      415

Asn Leu Glu Arg Arg Ile Asp Asn Leu Asn Lys Arg Met Gln Asp Gly
      420                      425                      430

Phe Leu Asp Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu
      435                      440                      445

Asn Glu Arg Thr Leu Asp Met His Asp Ala Asn Val Lys Asn Leu His
      450                      455                      460

Glu Lys Val Lys Ser Gln Leu Arg Asp Asn Ala Asn Asp Leu Gly Asn
      465                      470                      475                      480

Gly Cys Phe Glu Phe Trp His Lys Cys Asp Asn Glu Cys Ile Glu Ser
      485                      490                      495

Val Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Gln Thr Glu Ser Lys
      500                      505                      510

Leu Asn Arg Leu Lys Ile Glu Ser Val Lys Leu Glu Asn Leu Gly Val
      515                      520                      525

Tyr Gln Ile Leu Ala Ile Tyr Ser Thr Val Ser Ser Ser Leu Val Leu
      530                      535                      540

Val Gly Leu Ile Met Ala Met Gly Leu Trp Met Cys Ser Asn Gly Ser
      545                      550                      555                      560

Met Gln Cys Asn Val Cys Ile
      565

```

&lt;210&gt; SEQ ID NO 42

&lt;211&gt; LENGTH: 450

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 42

```

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Cys Met Val
1      5      10      15

Val Gly Ile Ile Ser Leu Met Leu Gln Ile Gly Asn Ile Ile Ser Ile
      20      25      30

Trp Val Ser His Ile Ile Gln Thr Gly His Pro Asn Gln Pro Gly Pro
      35      40      45

Cys Asn Gln Ser Ile Asn Phe Tyr Thr Glu Gln Ala Ala Ala Ser Val
      50      55      60

Thr Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Ser Gly Trp Ala Ile
      65      70      75      80

Tyr Ser Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly Asp Val Phe
      85      90      95

Val Met Arg Glu Pro Phe Val Ser Cys Ser His Leu Glu Cys Arg Thr
      100     105     110

Phe Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly
      115     120     125

Thr Val Lys Asp Arg Ser Pro Tyr Arg Thr Leu Met Ser Cys Pro Val
      130     135     140

Gly Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp
      145     150     155     160

Ser Ala Ser Ala Cys His Asp Gly Ile Ser Trp Leu Thr Ile Gly Ile
      165     170     175

Ser Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr Asn Gly Ile
      180     185     190

```

-continued

---

```

Ile Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Ile Leu Arg Thr Gln
 195                200                205

Glu Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr Val Met Thr
 210                215                220

Asp Gly Pro Ser Asn Glu Gln Ala Ser Tyr Lys Ile Phe Lys Ile Glu
 225                230                235                240

Lys Gly Lys Val Val Lys Ser Val Glu Leu Asn Ala Pro Asn Tyr His
 245                250                255

Tyr Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile Thr Cys Val
 260                265                270

Cys Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val Ser Phe Asn
 275                280                285

Gln Asn Leu Glu Tyr Gln Ile Gly Tyr Ile Cys Ser Gly Val Phe Gly
 290                295                300

Asp Ser Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly Pro Val Ser
 305                310                315                320

Leu Asn Gly Glu Tyr Gly Val Lys Gly Phe Ser Phe Lys Tyr Gly Asp
 325                330                335

Gly Val Trp Ile Gly Arg Thr Lys Ser Thr Ser Ser Arg Ser Gly Phe
 340                345                350

Glu Met Ile Trp Asp Pro Asn Gly Trp Thr Glu Thr Asp Ser Asn Phe
 355                360                365

Ser Leu Lys Gln Asp Ile Ile Ala Ile Thr Asp Trp Ser Gly Tyr Ser
 370                375                380

Gly Ser Phe Val Gln His Pro Glu Leu Thr Gly Leu Asn Cys Met Arg
 385                390                395                400

Pro Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys Glu Lys Thr
 405                410                415

Ile Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val Asn Ser Asp
 420                425                430

Thr Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Val Pro Phe Thr Ile
 435                440                445

Asp Lys
 450

```

&lt;210&gt; SEQ ID NO 43

&lt;211&gt; LENGTH: 566

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Influenza A virus

&lt;400&gt; SEQUENCE: 43

```

Met Ile Ala Ile Ile Ile Leu Ala Ile Val Val Ser Thr Ser Lys Ser
 1             5             10             15

Asp Arg Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Thr Gln Val
 20             25             30

Asp Thr Ile Leu Glu Lys Asn Val Thr Val Thr His Ser Val Glu Leu
 35             40             45

Leu Glu Asn Gln Lys Glu Asn Arg Phe Cys Arg Val Leu Asn Lys Ala
 50             55             60

Pro Leu Asp Leu Met Asp Cys Thr Thr Glu Gly Trp Ile Leu Gly Asn
 65             70             75             80

Pro Arg Cys Asp Asn Leu Leu Gly Asp Gln Ser Trp Ser Tyr Ile Val

```

-continued

85								90					95				
Glu	Arg	Pro	Asp	Ala	Gln	Asn	Gly	Ile	Cys	Tyr	Pro	Gly	Val	Leu	Lys		
			100					105					110				
Glu	Thr	Glu	Glu	Leu	Lys	Ala	Leu	Ile	Gly	Ser	Ile	Asp	Ser	Ile	Gln		
		115					120					125					
Arg	Phe	Glu	Met	Phe	Pro	Lys	Ser	Thr	Trp	Thr	Gly	Val	Asp	Thr	Asn		
	130					135					140						
Ser	Gly	Val	Thr	Ser	Ala	Cys	Pro	Tyr	Asn	Gly	Glu	Ser	Ser	Phe	Tyr		
145					150					155					160		
Arg	Asn	Leu	Leu	Trp	Ile	Ile	Lys	Ile	Arg	Ser	Asp	Pro	Tyr	Ser	Leu		
			165						170					175			
Ile	Lys	Gly	Thr	Tyr	Thr	Asn	Thr	Gly	Ser	Gln	Pro	Ile	Leu	Tyr	Phe		
		180						185					190				
Trp	Gly	Val	His	His	Pro	Pro	Asp	Glu	Val	Glu	Gln	Ala	Asn	Leu	Tyr		
		195					200					205					
Gly	Ile	Gly	Thr	Arg	Tyr	Val	Arg	Met	Gly	Thr	Glu	Ser	Met	Asn	Phe		
	210					215					220						
Ala	Lys	Gly	Pro	Glu	Ile	Ala	Gly	Arg	Pro	Pro	Ala	Asn	Gly	Gln	Arg		
225					230					235					240		
Gly	Arg	Ile	Asp	Tyr	Tyr	Trp	Ser	Val	Leu	Lys	Pro	Gly	Glu	Thr	Leu		
			245						250					255			
Asn	Val	Glu	Ser	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Tyr	Lys		
		260						265					270				
Phe	Thr	Ser	Ser	Arg	Asn	Lys	Gly	Ala	Ile	Phe	Lys	Ser	Asp	Leu	Pro		
		275					280					285					
Ile	Glu	Asn	Cys	Asp	Ala	Val	Cys	Gln	Thr	Leu	Ala	Gly	Ala	Ile	Asn		
	290					295					300						
Thr	Asn	Lys	Thr	Phe	Gln	Asn	Ile	Ser	Pro	Val	Trp	Ile	Gly	Glu	Cys		
305					310					315					320		
Pro	Lys	Tyr	Val	Lys	Ser	Lys	Ser	Leu	Lys	Leu	Ala	Thr	Gly	Leu	Arg		
			325						330					335			
Asn	Val	Pro	Gln	Ala	Glu	Thr	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly		
		340						345				350					
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	Tyr		
	355						360					365					
His	His	Glu	Asn	Ser	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Lys	Glu	Ser		
	370					375					380						
Thr	Gln	Lys	Ala	Ile	Asp	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Ile	Ile		
385					390					395					400		
Asp	Lys	Met	Asn	Thr	Gln	Phe	Glu	Ala	Val	Glu	His	Glu	Phe	Ser	Ser		
			405						410					415			
Leu	Glu	Arg	Arg	Ile	Gly	Asn	Leu	Asn	Lys	Arg	Met	Glu	Asp	Gly	Phe		
		420						425					430				
Leu	Asp	Val	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn		
	435						440					445					
Glu	Arg	Thr	Leu	Asp	Met	His	Asp	Ala	Asn	Val	Lys	Asn	Leu	His	Glu		
	450					455					460						
Lys	Val	Lys	Ser	Gln	Leu	Arg	Asp	Asn	Ala	Lys	Asp	Leu	Gly	Asn	Gly		
465					470					475					480		
Cys	Phe	Glu	Phe	Trp	His	Lys	Cys	Asp	Asn	Glu	Cys	Ile	Asn	Ser	Val		
			485						490					495			

-continued

---

Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Gln Glu Glu Ser Arg Leu  
500 505 510

Asn Arg Gln Glu Ile Lys Ser Val Met Leu Glu Asn Leu Gly Val Tyr  
515 520 525

Gln Ile Leu Ala Ile Tyr Ser Thr Val Ser Ser Ser Leu Val Leu Val  
530 535 540

Gly Leu Ile Ile Ala Met Gly Leu Trp Met Cys Ser Asn Gly Ser Met  
545 550 555 560

Gln Cys Lys Ile Cys Ile  
565

<210> SEQ ID NO 44  
<211> LENGTH: 469  
<212> TYPE: PRT  
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 44

Met Asn Pro Asn Gln Arg Ile Ile Thr Ile Gly Ser Val Ser Leu Thr  
1 5 10 15

Ile Ala Thr Val Cys Phe Leu Met Gln Ile Ala Ile Leu Ala Thr Thr  
20 25 30

Val Thr Leu His Phe Lys Gln Asn Glu Cys Ser Ile Pro Ala Asn Asn  
35 40 45

Gln Val Thr Pro Cys Glu Pro Ile Val Ile Glu Arg Asn Ile Thr Glu  
50 55 60

Ile Val Tyr Leu Asn Asn Thr Thr Ile Glu Lys Glu Ile Cys Pro Glu  
65 70 75 80

Val Val Glu Tyr Arg Asn Trp Ser Lys Pro Gln Cys Gln Ile Thr Gly  
85 90 95

Phe Ala Pro Phe Ser Lys Asp Asn Ser Ile Arg Leu Ser Ala Gly Gly  
100 105 110

Asp Ile Trp Ile Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Ser Lys  
115 120 125

Cys Tyr Gln Phe Ala Leu Gly Gln Gly Thr Thr Leu Asp Asn Lys His  
130 135 140

Ser Asn Gly Thr Ile His Asp Arg Ile Pro His Arg Thr Leu Leu Met  
145 150 155 160

Asn Glu Leu Gly Val Pro Phe His Leu Gly Thr Lys Gln Val Cys Ile  
165 170 175

Ala Trp Ser Ser Ser Ser Cys His Asp Gly Lys Ala Trp Leu His Val  
180 185 190

Cys Val Thr Gly Asp Asp Arg Asn Ala Thr Ala Ser Phe Ile Tyr Asp  
195 200 205

Gly Met Leu Ile Asp Ser Ile Gly Ser Trp Ser Gln Asn Ile Leu Arg  
210 215 220

Thr Gln Glu Ser Glu Cys Val Cys Ile Ser Gly Thr Cys Thr Val Val  
225 230 235 240

Met Thr Asp Gly Ser Ala Ser Gly Arg Ala Asp Thr Arg Ile Leu Phe  
245 250 255

Ile Arg Glu Gly Lys Ile Val His Ile Ser Pro Leu Ser Gly Ser Ala  
260 265 270

Gln His Val Glu Glu Cys Ser Cys Tyr Pro Arg Tyr Pro Asn Val Arg

-continued

275	280	285
Cys Val Cys Arg Asp Asn Trp Lys Gly Ser Asn Arg Pro Val Ile Asp 290 295 300		
Ile Asn Met Ala Asp Tyr Ser Ile Asp Ser Ser Tyr Val Cys Ser Gly 305 310 315 320		
Leu Val Gly Asp Thr Pro Arg Asn Asp Asp Ser Ser Ser Ser Asn 325 330 335		
Cys Arg Asp Pro Asn Asn Glu Arg Gly Asn Pro Gly Val Lys Gly Trp 340 345 350		
Ala Phe Asp Asn Gly Asn Asp Val Trp Met Gly Arg Thr Ile Ser Lys 355 360 365		
Asp Ser Arg Ser Gly Tyr Glu Thr Phe Lys Val Ile Gly Gly Trp Ala 370 375 380		
Ile Ala Asn Ser Lys Ser Gln Thr Asn Arg Gln Val Ile Val Asp Asn 385 390 395 400		
Asn Asn Trp Ser Gly Tyr Ser Gly Ile Phe Ser Val Glu Ser Lys Gly 405 410 415		
Cys Ile Asn Arg Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg Pro Gln 420 425 430		
Glu Thr Arg Val Trp Trp Thr Ser Asn Ser Ile Val Val Phe Cys Gly 435 440 445		
Thr Ser Gly Thr Tyr Gly Thr Gly Ser Trp Pro Asp Gly Ala Asn Ile 450 455 460		
Asp Phe Met Pro Ile 465		

<210> SEQ ID NO 45  
 <211> LENGTH: 1464  
 <212> TYPE: DNA  
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 45

agcaaaagca gggatgatcga gaataaatcc aaatcagaaa ctatttgcat tatctggagt	60
ggcaatagca cttagtgtac tgaacttatt gataggaatc tcaaacgtcg gattgaacgt	120
atctctacat ctaaaggaaa aaggacccaa acaggaggag aatttaacat gcacgaccat	180
taatcaaaac aacactactg tagtagaaaa cacatatgta aataatacaa caataattac	240
caaggggaact gatttgaaaa caccaagcta tctgctgttg aacaagagcc tgtgcaatgt	300
tgaagggtgg gtcgtgatag caaaagacaa tgcagtaaga tttggggaaa gtgaacaaat	360
cattgttacc agggagccat atgtatcatg cgacccaaca ggatgcaaaa tgtatgcctt	420
gcaccaaggg actaccatta ggaacaaaca ttcaaatgga acgattcatg acagaacagc	480
tttcagaggt ctcattccca ctccattggg cactccacca accgtaagta acagtgactt	540
tatgtgtgtt ggatggtcaa gcacaacttg ccatgatggg attgctagga tgactatctg	600
tatacaagga aataatgaca atgctacagc aacggtttat tacaacagaa ggctgaccac	660
taccattaag acctgggcca gaaacattct gaggactcaa gaatcagaat gtgtgtgcca	720
caatggcaca tgtgcagttg taatgaccga cggatcggct agtagtcaag cctatacaaa	780
agtaatgtat ttccacaagg gattagtagt taaggaggag gagttaaggg gatcagccag	840
acatatgtag gaatgctcct gttatggaca caatcaaaag gtgacctgtg tgtgcagaga	900



-continued

taactggcag ggagcaaaca ggcctattat agaaattgat atgagcacat tggagcacac	960
aagtagatac gtgtgcactg gaattctcac agacaccagc agacctgggg acaaatctag	1020
tggtgattgt tccaatccaa taactgggag tcccggcgtt cggggagtga agggattcgg	1080
gtttctaaat ggggataaca catggcttgg taggaccatc agccccagat caagaagtgg	1140
attcgaaatg ttgaaaatac ctaatgcagg tactgatccc aattctagaa tagcagaacg	1200
acaggaaatt gtcgacaata acaattggtc aggctattcc ggaagcttta ttgactattg	1260
gaatgataac agtgaatgct acaatccatg cttttacgta gagttaatta gaggaagacc	1320
cgaagaggct aaatacgtat ggtgggcaag taacagtcta attgcctat gtggaagccc	1380
attcccagtt gggctcgtt ccttcccga tggggcacia atccaatact tttcgtaaaa	1440
tgcaaaaaca ccctgtttc tact	1464

What is claimed is:

1. An isolated polypeptide, wherein said polypeptide is selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence encoded by the nucleotide sequence as shown in any one of SEQ ID NOS:21-26 or 33-38 or 45;
- b) a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS:27-32 or 39-44;
- c) the mature form of a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44;
- d) a polypeptide comprising an amino acid sequence encoded by a polynucleotide which hybridizes under highly stringent conditions to a polynucleotide comprising a nucleotide sequence encoding (a) (b) or (c); and
- e) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the polypeptide of (b).

2. An immunogenic composition comprising an immunologically effective amount of at least one polypeptide of claim 1.

3. An isolated antibody that specifically binds the polypeptide of claim 1.

4. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the polypeptide of claim 1 in a physiologically acceptable carrier.

5. A recombinant influenza virus comprising the polypeptide of claim 1.

6. An immunogenic composition comprising an immunologically effective amount of the recombinant influenza virus of claim 5.

7. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the recombinant influenza virus of claim 5 in a physiologically acceptable carrier.

8. An isolated polynucleotide, wherein said polynucleotide is selected from the group consisting of:

- a) a polynucleotide comprising the nucleotide sequence as shown in any one of SEQ ID NOS: 21-26 or 33-38 or 45, or a complementary sequence thereof;
- b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44, or a complementary nucleotide sequence thereof;
- c) a polynucleotide which hybridizes under highly stringent conditions over substantially the entire length of the polynucleotide of (a); and
- d) a polynucleotide comprising a nucleotide sequence having at least 98% sequence identity to the polynucleotide of (a).

9. An immunogenic composition comprising at least one polynucleotide of claim 8.

10. A cell comprising at least one polynucleotide of claim 8.

11. A vector comprising the polynucleotide of claim 8.

12. The vector of claim 11, wherein the vector is a plasmid, a cosmid, a phage, a virus, or a fragment of a virus.

13. The vector of claim 12, wherein the vector is an expression vector.

14. A cell comprising the vector of claim 13.

15. An influenza virus comprising one or more polynucleotides of claim 8.

16. The virus of claim 15, wherein the virus is a reassortant virus.

17. A 6:2 reassortant influenza virus, wherein said virus comprises 6 internal genome segments from A/Ann Arbor/6/60 and 2 genome segments that encode an HA and/or a NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS:27-32, and 39-44.

18. A method of producing a reassortant influenza virus, the method comprising: culturing the cell of claim 14 in a suitable culture medium under conditions permitting expression of said polynucleotide; and, isolating the reassortant influenza virus from a cell population comprising said cell or the medium.

19. An immunogenic composition comprising an immunologically effective amount of the reassortant influenza virus of claim 17.

20. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the reassortant influenza virus of claim 17 in a physiologically effective carrier.

21. A method of producing an isolated or recombinant polypeptide, the method comprising: culturing the cell of claim 10 in a suitable culture medium under conditions permitting expression of said polynucleotide; and, isolating the polypeptide from the cell or the medium.

22. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject, the virus of claim 17 in an amount effective to produce an immunogenic response against the viral infection.

23. The method of claim 22, wherein the subject is a human.

24. The immunogenic composition of claim 19, wherein the hemagglutinin comprises a modified polybasic cleavage site.

25. A live attenuated influenza vaccine comprising the composition of claim 19.

26. A split virus or killed virus vaccine comprising the composition of claim 19.

27. A live attenuated influenza vaccine comprising the composition of claim 24.

28. A split virus or killed virus vaccine comprising the composition of claim 24.

29. A method for producing an influenza virus in cell culture, the method comprising:

- i) introducing into a population of host cells, which population of host cells is capable of supporting replication of influenza virus, a plurality of vectors comprising nucleotide sequences corresponding to at least 6 internal genome segments of A/Ann Arbor/6/60; and, at least one genome segment comprising a polynucleotide encoding an HA and/or a NA polypeptide selected from the group consisting of: the polypeptides of SEQ ID NOS:27-32, and 39-44,
- ii) culturing the population of host cells at a temperature less than or equal to 35° C.; and,
- iii) recovering an influenza virus.

30. The method of claim 29, wherein the polynucleotide encoding the HA and/or NA polypeptide is selected from the group consisting of:

a) a polynucleotide comprising the nucleotide sequence of any one of SEQ ID NOS:21, 23-26 or 33-38, or 45, or a complementary nucleotide sequence thereof;

b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as shown in any one of SEQ ID NOS: 27-32 or 39-44, or a complementary nucleotide sequence thereof;

c) a polynucleotide which hybridizes under highly stringent conditions over substantially the entire length of the polynucleotide of (a); and

d) a polynucleotide comprising a nucleotide sequence having at least 98% sequence identity to the polynucleotide of (a).

31. An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of claim 29.

32. An immunogenic composition comprising an immunologically effective amount of the influenza virus produced by the method of claim 30.

33. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus produced by the method of claim 29 in a physiologically effective carrier.

34. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual an immunologically effective amount of the influenza virus produced by the method of claim 30 in a physiologically effective carrier.

35. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of claim 31.

36. A method for stimulating the immune system of an individual to produce a protective immune response against influenza virus, the method comprising administering to the individual the immunogenic composition of claim 32.

37. A live attenuated influenza vaccine comprising the immunogenic composition of claim 31.

38. A split virus or killed virus vaccine comprising the immunogenic composition of claim 32.

\* \* \* \* \*