

A New Empirical Constraint on the Prevalence of Technological Species in the Universe

A. Frank¹ and W.T. Sullivan III²

Abstract

In this article, we address the cosmic frequency of technological species. Recent advances in exoplanet studies provide strong constraints on all astrophysical terms in the Drake equation. Using these and modifying the form and intent of the Drake equation, we set a firm lower bound on the probability that one or more technological species have evolved anywhere and at any time in the history of the observable Universe. We find that as long as the probability that a habitable zone planet develops a technological species is larger than $\sim 10^{-24}$, humanity is not the only time technological intelligence has evolved. This constraint has important scientific and philosophical consequences. Key Words: Life—Intelligence—Extraterrestrial life. Astrobiology 2016, xxx–xxx.

1. Introduction

THE HISTORY of physics has shown that fundamental insights into a problem can sometimes be acquired by setting order-of-magnitude estimates of scales, limits, and boundaries via combinations of constants or parameters. This approach can be particularly effective for problems in which few empirically derived constraints are available, such as the derivation of the quantum gravitational Planck length, that is, $l_p = (\hbar G/c^3)^{1/2}$, or early studies using the cosmic density parameter ($\Omega = \rho/\rho_c$). Here we adopt such an approach to address an overarching goal of astrobiology: understanding if humanity is “alone” (Sullivan and Baross, 2007; Impey *et al.*, 2012). One way of framing this question is to ask if other technology-building species exist now, or have ever existed at any time, in the Universe. Given that we have only one known example of a planet where such evolution has occurred (and also only one example of where even microbial life has evolved), there seems to be little hope in determining the fecundity of the Universe in producing technological species. This conclusion, however, ignores the rapid and substantial progress astrobiology has made in the last two decades. In particular, the empirical determination of exoplanet statistics has radically changed the nature and quality of constraints astrobiologists now have at their disposal when considering the prevalence of life in the Universe. In this article, we employ these new constraints to set a lower limit on the probability that technological species have ever evolved anywhere other than on Earth.

2. Method

Our approach asks a very different question from the usual treatment of the subject. Standard astrobiological discussions of intelligent life focus on how many technological species *currently* exist with which we might communicate (Vakoch and Dowd, 2015). But rather than asking whether we are *now* alone, we ask whether we are the only technological species that has *ever* arisen. Such an approach allows us to set limits on what might be called the “cosmic archaeological question”: How often in the history of the Universe has evolution ever led to a technological species, whether short- or long-lived? As we shall show, providing constraints on an answer to this question has profound philosophical and practical implications.

We first modify the Drake equation in order to address how many technological species have formed over the history of the observable Universe. We call this number A (for archaeology) and use it to investigate the probability that humanity is unique (*i.e.*, $A = 1$). Note that we are explicitly *not* concerned with the average lifetime ($\langle L \rangle$) of such species or if they still exist such that we could receive their signals or signal them. This is in contrast to the usual Drake equation formulation, which calculates N_c , the number of technological species *now* existing (hence its concern with $\langle L \rangle$). Given this approach, effects of time are removed where they normally appear in the form of star-formation rates, stellar lifetimes, technological species lifetimes, and distribution of epochs of arising. As usual for the Drake equation,

¹Department of Physics and Astronomy, University of Rochester, Rochester, New York.

²Department of Astronomy and Astrobiology Program, University of Washington, Seattle, Washington.

we assume that technology is associated with planets and their host stars.

We define the “A-form” of the Drake equation, which describes the total number of technological species that have ever evolved anywhere in the currently observable Universe:

$$A = [N_* f_p n_p] [f_i f_t] \quad (1)$$

$$= N_{\text{ast}} f_{\text{bt}}$$

where N_* is the total number of stars, f_p is the fraction of those stars that form planets, n_p is the average number of planets in the habitable zone of a star with planets, f_i is the probability that a habitable zone planet develops life, f_t is the probability that a planet with life develops intelligence, and f_{bt} is the probability that a planet with intelligent life develops technology (of the “energy intensive” kind such as that of our own civilization).

The second version of Eq. 1 reduces the right-hand side to two factors, the first of which, N_{ast} , includes all factors involving astrophysics and represents the total number of habitable zone planets. The second factor, f_{bt} , gathers the three factors involved with biology, evolution, and “planetary sociology,” and represents the total “biotechnical” probability that a given habitable zone planet has ever evolved a technological species. The factor $f_{\text{bt}} = f_i f_t$ is extremely uncertain because (a) we have no theory to guide any estimates and (b) we have only one known example of the occurrence and history of life, intelligence, and technology. In what follows, we leave f_{bt} as statistically unknown at this time and examine the consequences of its taking on various values depending on one’s pessimism or optimism.

The key development that enables our new approach is that observations now provide accepted or statistical well-determined values for all factors contributing to N_{ast} . The importance of this accomplishment cannot be overstated—until recently only one factor (N_*) was known, and it was entirely possible that habitable zone planets might have been extremely rare ($N_{\text{ast}}/N_* \ll 1$). But the combination of radial velocity, transit, and microlensing based methods now yields statistically well-constrained values for both f_p and n_p .¹

We adopt the values $f_p \sim 1.0$ (Cassan *et al.*, 2012) and $n_p \sim 0.2$ (Petigura *et al.*, 2013); therefore $N_{\text{ast}}/N_* = 0.2$. Note that the number of stars N_* is also an observable quantity, which depends on the size of the region being considered.

3. Results

We now turn to the specific question, “Has even one other technological species ever existed in the observable Universe?” We take $N_* = 2 \times 10^{22}$ for the total number of stars in the observable Universe (Silburt *et al.*, 2015) To address our question, A is set to a conservative value en-

suring that Earth is the only location in the history of the cosmos where a technological civilization has ever evolved. Adopting $A = 0.01$ means that in a statistical sense were we to rerun the history of the Universe 100 times, only once would a lone technological species occur. A lower bound \bar{f}_{bt} on the probability is then

$$\bar{f}_{\text{bt}} = A/N_{\text{ast}} = 0.01/4 \times 10^{21} = 2.5 \times 10^{-24}$$

Thus only for values of the product f_{bt} lower than 2.5×10^{-24} are we likely to be alone and singular in the history of the observable Universe. This limiting value of f_{bt} can be considered to define a “pessimism line” in discussions of the prevalence of technological civilizations on a cosmic scale. On the other hand, if evolutionary processes lead to higher values, we can be assured that we are not the only instance in which the Universe has hosted a technological species.

We can generalize these results for structural scales (of size R_s) within the Universe: galaxies, clusters of galaxies, and superclusters of galaxies (Fukugita and Peebles, 2004). Table 1 provides typical values for these entities and lists the corresponding values of f_{bt} for each of them, as well as for the entire Universe. For instance, for our own galaxy f_{bt} is 1.7×10^{-11} , meaning that another technical species has likely occurred in the history of the Milky Way if the probability of a technological species arising on a *given planet in a habitable zone* is greater than one in 60 billion. Figure 1 presents these results for given values of f_{bt} . The figure allows one to see the corresponding number of technological species that have ever arisen on various scales for various assumptions of the difficulty of biology and technology to evolve.

4. Discussion and Conclusions

These results have wide implications. First, the long history of debate over extraterrestrial intelligence can be characterized as one of pessimists versus optimists relative to choices for the various factors in the Drake equation (Vakoch and Dowd, 2015). These debates have, however, been unconstrained in terms of how pessimistic one can be. With our approach we have, for the first time, provided a quantitative and empirically constrained limit on what it means to be pessimistic about the likelihood of another technological species ever having arisen in the history of the Universe. We have done so by segregating newly measured astrophysical factors from the fully unconstrained biotechnical ones, and by shifting the focus toward a question of “cosmic archaeology” and away from technological species lifetimes. Our constraint addresses an issue that is of particular scientific and philosophical consequence: the question “Have they ever existed?” rather than the usual narrower concern of the Drake equation, “Do they exist now?” Perhaps in the long term we should contemplate undertaking a field survey in cosmic archaeology, seeking possible evidence for such past technical species (Stevens *et al.*, 2015).

Secondly, we note that sample sizes A of order 100 or 1000 have a particular importance in current discussions over efforts to create a sustainable, energy-intensive, high-technology civilization here on Earth (Frank and Sullivan, 2014). Given the challenges human society faces from

¹Rather than f_p and n_p , some authors focus on η_E , defined as the occurrence rate of Earth-sized planets (1–2 Earth radii) in the habitable zones of Sun-like stars. For example, one recent study based on Kepler mission transiting planets finds a value $\eta_E \sim 0.06$ (Silburt *et al.*, 2015). For the purposes of our order-of-magnitude estimates, which also include non-Sun-like stars, we stay with the factors in the conventional Drake equation.

TABLE 1. LIMITS ON THE NUMBER OF TECHNOLOGICAL SPECIES OCCURRING ON DIFFERENT SCALE LENGTHS

| Scale length | Size R_s (M lt-yr) | No. galaxies | N_{ast} | f_{bt} for $A=1$ |
|---------------------|----------------------|--------------------|--------------------|-----------------------|
| Galaxy | 0.1 | 1 | 6×10^{10} | 1.7×10^{-11} |
| Galaxy cluster | 5 | 300 | 2×10^{13} | 5×10^{-14} |
| Supercluster | 300 | 3000 | 2×10^{14} | 5×10^{-15} |
| Observable Universe | 13,700 | 7×10^{10} | 4×10^{21} | 2.5×10^{-22} |

climate change, resource allocation, and loss of species diversity, it is not clear if the kind of long-term global culture we hope to build is even sustainable. However $A > 100$ in our formulation implies that the evolution of technological species has occurred enough times that the ensemble of their histories (or trajectories in a suitably defined phase space) is statistically meaningful. In particular, for large-enough values of f_{bt} , the average longevity $\langle L \rangle$ of such a sample of technological species does, in principle, exist. Thus if f_{bt} is such that $A \geq 1000$ (the statistically relevant population), it is reasonable to consider that different versions of humanity's current technological experiment in the alteration of our planetary system have occurred before. Note that many aspects of the feedback between an energy-intensive, technological species and its host planet would depend solely on physical processes and constraints (atmospheric chemistry

changes, hydrological cycle disruptions, etc.; Frank and Sullivan, 2014). Thus modeling ensembles of such species to understand the broad classifications of their planetary feedback histories is theoretically well-grounded for large-enough values of f_{bt} . Such a project would be an attempt to understand both average properties like $\langle L \rangle$ as well as what led some trajectories to collapse and others to long-term sustainability.

It is also possible to make the connection between our A form of the Drake equation and traditional SETI studies that ask about N_c , the number of technological species existing in our galaxy *now*. The traditional form of the Drake equation can be written

$$N_c = [N_{ast}] [f_{bt}] \frac{\langle L \rangle}{L_*}$$

Thus $N_c = A \frac{\langle L \rangle}{L_*}$. As we have shown in Table 1, for each astrophysical scale of interest s , we can define a minimum biotechnical probability via $A_{\min}^s = [N_{ast}^s] [f_{bt, \min}^s] = 1$. Using this, we can then ask at each scale how much does f_{bt}^s have to be increased above $f_{bt, \min}^s$ to get some desired value A^s above A_{\min}^s . Thus we can write

$$N_c^s = [N_{ast}^s] [f_{bt, \min}^s] A^s \frac{\langle L \rangle}{L_*}$$

Taking L_* , the lifetime of a star, to be $\sim 10^{10}$ yr and using Table 1 for a single galaxy (our own), one has

$$N_c^g = [6 \times 10^{10}] [1.7 \times 10^{-11} A^g] \frac{\langle L \rangle}{10^{10} \text{ yr}} = 10^{-10} A^g \langle L \rangle$$

In order for there to be one other partner technological species in the Galaxy with which to make contact ($N_c^g = 2$), the product $A^g \langle L \rangle$ must be equal to 2×10^{10} yr. For an optimistic upper limit of $\langle L \rangle = 10^6$ yr, A^g must be at least 2×10^4 (i.e., there must have been at least 20,000 technological species spawned across the lifetime of the Galaxy). For a perhaps more realistic guess (based on our own civilization) of $\langle L \rangle = 10^4$ yr, $A^g = 2 \times 10^6$.

Casting these results in a different way, one can imagine conducting a general Milky Way SETI search that might hope for one in every million stars to host a technical species today, which means $N_c^g = 3 \times 10^5$. To achieve this number, if we very optimistically adopt $f_{bt} = 1$, then $\langle L \rangle$ must be 5×10^4 yr. More realistic lower values of f_{bt} require proportionately higher values for $\langle L \rangle$ to reach the desired number of potential finds. Note that in all cases the number of technological species that have ever existed in our galaxy would be $6 \times 10^{10} f_{bt}$.




FIG. 1. The number A of technological species that have ever occurred over the history of the Universe versus the scale size R_s of different hierarchical structures in which their host stars are found: galaxies, clusters of galaxies, superclusters of galaxies, and the entire Universe. Reference values of $A=0.01$, 1, and 100 are marked. The four curves represent different values of f_{bt} , the probability of a technological species arising on a given habitable zone planet. Only if f_{bt} falls as low as 2.5×10^{-24} to 2.5×10^{-22} is it likely that no other technological species has ever arisen in the entire Universe. Color images available online at www.liebertpub.com/ast

In conclusion, we have shown that recent advances in astrobiology and exoplanet studies mean that an empirically derived lower limit can now be placed on the probabilities that even one other technological species has ever evolved in the Universe or in our galaxy. This limit provides a framework for discussions of both life in its cosmic context and questions about trajectories of technological species relevant to our own issues of global sustainability.

Acknowledgments

We thank Dan Watson and David Catling for comments on the text, and Matt McQuinn, Caleb Scharf, and Gavin Schmidt for discussion.

References

- Cassan, A., Kubas, D., Beaulieu, J.-P., Dominik, M., Horne, K., Greenhill, J., Wambsganss, J., Menzies, J., Williams, A., Jørgensen, U.G., Udalski, A., Bennett, D.P., Albrow, M.D., Batista, V., Brillant, S., Caldwell, J.A.R., Cole, A., Coutures, Ch., Cook, K.H., Dieters, S., Prester, D.D., Donatowicz, J., Fouqué, P., Hill, K., Kains, N., Kane, S., Marquette, J.-B., Martin, R., Pollard, K.R., Sahu, K.C., Vinter, C., Warren, D., Watson, B., Zub, M., Sumi, T., Szymański, M.K., Kubiak, M., Poleski, R., Soszynski, I., Ulaczyk, K., Pietrzyński, G., and Wyrzykowski, Ł. (2012) One or more bound planets per Milky Way star from microlensing observations. *Nature* 481:167–169.
- Frank, A. and Sullivan, W. (2014) Sustainability and the astrobiological perspective: framing human futures in a planetary context. *Anthropocene* 5:32–41.
- Fukugita, M. and Peebles, P.J.E. (2004) The cosmic energy inventory. *Astrophys J* 616:643–668.
- Impey, C., Lunine, J., and Funes, J., editors. (2012) *Frontiers of Astrobiology*, Cambridge University Press, Cambridge, UK.
- Petigura, E.A., Howard, A.W., and Marcy, G.W. (2013) Prevalence of Earth-size planets orbiting Sun-like stars. *Proc Natl Acad Sci USA* 110:19273–19278.
- Silburt, A., Gaidos, E., and Wu, Y. (2015) Statistical reconstruction of the planet population around Kepler solar-type stars. *Astrophys J* 799, doi:10.1088/0004-637X/799/2/180.
- Stevens, A., Forgan, D., and O'Malley James, J. (2015) Observational signatures of self-destructive civilisations. arXiv 1507.08530
- Sullivan, W.T. and Baross, J.A., editors. (2007) *Planets and Life: The Emerging Science of Astrobiology*, Cambridge University Press, Cambridge, UK.
- Vakoch, D.A. and Dowd, M.F., editors. (2015) *The Drake Equation: Estimating the Prevalence of Extraterrestrial Life through the Ages*, Cambridge University Press, Cambridge, UK.

Address correspondence to:

A. Frank
Department of Physics and Astronomy
University of Rochester
Rochester, NY 14620

E-mail: afrank@pas.rochester.edu

Submitted 5 October 2015

Accepted 16 February 2016